



Glomerular Filtration Rate (GFR)

You are scheduled for a glomerular filtration rate (GFR) study. This test measures how well your kidneys work to filter waste products from your body. It uses a small amount of radioactive material (less than that of a chest x-ray). This material helps measure the kidney's filtering ability. The test will take place in the Nuclear Medicine Department.

Preparation:

- You will be asked not to take any medications that may interfere with this test. Your nurse or doctor will inform you which medications should not be taken.
- I.V. (intravenous) lines will be inserted: one in each arm. One will be used to give you the radioactive drug and the other will be used to take blood samples. The line from which blood samples will be taken is called a heparin lock. This device allows blood samples to be taken without repeated needle sticks.
- Your weight will be measured before the test.

Procedure:

- A small amount of radioactive material will be injected into your blood through one of the I.V. lines already in place.
- Two blood specimens will be drawn: 1 hour after the injection and 3 hours after the injection.
- After the second blood sample is drawn, the I.V. lines will be removed.

After the Procedure:

Most patients feel the same before and after the test. They may return to their rooms or go home (if they are outpatients).


If you have questions about the procedure, please ask. Your nurse and doctor are ready to assist you at all times.

Special Instructions:

- Because it uses radioactivity, this test is not performed in pregnant women. *If you are pregnant or think you might be pregnant, please*

inform your doctor immediately so that a decision can be made about this test.

- *Also, please inform your doctor immediately if you are breast-feeding. Some tests can be performed in breast-feeding women if they are willing to stop breast-feeding for a while.*



3/00

This information is prepared specifically for patients participating in clinical research at the Warren Grant Magnuson Clinical Center at the National Institutes of Health and is not necessarily applicable to individuals who are patients elsewhere. If you have questions about the information presented here, talk to a member of your healthcare team.

Questions about the Clinical Center? OCCC@cc.nih.gov

Where applicable, brand names of commercial products are provided only as illustrative examples of acceptable products, and do not imply endorsement by NIH; nor does the fact that a particular brand name product is not identified imply that such product is unsatisfactory.

National Institutes of Health
Warren Grant Magnuson Clinical Center
Bethesda, Maryland 20892