

**Table 29. Tuberculosis Cases by Initial Drug Regimen: 59 Reporting Areas, 2000**

Reporting Area	Total Cases	Cases Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percent of Cases with Initial Drug Regimen <sup>1,2</sup>		
			No.	%	IR	IRZ	IRZ,E/S
<b>United States</b>	<b>16,377</b>	<b>15,947</b>	<b>15,874</b>	<b>99.5</b>	<b>1.9</b>	<b>10.3</b>	<b>78.2</b>
Alabama	310	296	296	100.0	0.3	68.9	27.4
Alaska	108	108	108	100.0	0.0	20.4	75.9
Arizona	261	253	253	100.0	2.4	9.5	75.9
Arkansas	199	188	186	98.9	29.6	49.5	14.0
California	3,297	3,242	3,240	99.9	1.1	5.5	86.3
Colorado	97	93	92	98.9	1.1	6.5	82.6
Connecticut	105	102	101	99.0	3.0	11.9	70.3
Delaware	28	27	27	100.0	7.4	11.1	77.8
District of Columbia	85	80	78	97.5	0.0	2.6	85.9
Florida	1,171	1,123	1,123	100.0	0.8	8.4	83.5
Georgia	703	679	675	99.4	1.2	12.0	74.8
Hawaii	136	134	134	100.0	3.7	14.2	73.9
Idaho	16	16	16	100.0	0.0	18.8	81.3
Illinois	743	728	726	99.7	2.1	7.7	80.6
Indiana	145	138	138	100.0	2.9	18.8	68.8
Iowa	40	40	40	100.0	0.0	30.0	65.0
Kansas	77	76	76	100.0	1.3	18.4	73.7
Kentucky	147	142	142	100.0	1.4	15.5	78.2
Louisiana	331	313	303	96.8	3.3	13.9	75.2
Maine	24	24	24	100.0	0.0	33.3	66.7
Maryland	282	277	276	99.6	0.4	2.9	92.8
Massachusetts	285	281	279	99.3	1.4	4.3	85.7
Michigan	287	277	277	100.0	7.9	28.9	56.7
Minnesota	178	178	178	100.0	0.6	10.7	73.0
Mississippi	173	171	171	100.0	0.0	15.2	83.0
Missouri	211	204	204	100.0	0.5	3.9	62.7
Montana	21	21	21	100.0	0.0	19.0	71.4
Nebraska	24	23	23	100.0	0.0	34.8	52.2
Nevada	96	91	90	98.9	0.0	8.9	84.4
New Hampshire	22	22	22	100.0	0.0	13.6	86.4
New Jersey	565	544	538	98.9	1.9	9.5	75.5
New Mexico	46	44	44	100.0	2.3	47.7	45.5
New York State <sup>3</sup>	412	405	405	100.0	1.2	7.7	85.7
New York City	1,332	1,308	1,308	100.0	1.3	4.2	81.5
North Carolina	447	430	430	100.0	0.5	4.2	88.4
North Dakota	5	5	5	100.0	0.0	20.0	80.0
Ohio	340	330	330	100.0	4.5	20.9	62.1
Oklahoma	154	149	148	99.3	9.5	14.9	56.1
Oregon	119	117	117	100.0	0.0	4.3	85.5
Pennsylvania	383	375	365	97.3	0.3	5.8	75.6
Rhode Island	49	49	49	100.0	0.0	0.0	87.8
South Carolina	286	269	269	100.0	2.2	6.7	73.2
South Dakota	16	14	14	100.0	0.0	42.9	57.1
Tennessee	383	369	369	100.0	3.3	15.4	75.6
Texas	1,506	1,476	1,450	98.2	1.7	7.7	79.0
Utah	49	48	48	100.0	0.0	6.3	85.4
Vermont	4	4	4	100.0	0.0	25.0	75.0
Virginia	292	285	283	99.3	1.4	6.7	85.9
Washington	258	254	254	100.0	2.8	3.9	88.6
West Virginia	33	32	32	100.0	3.1	28.1	53.1
Wisconsin	92	89	89	100.0	2.2	11.2	79.8
Wyoming	4	4	4	100.0	0.0	25.0	50.0
American Samoa <sup>4</sup>	...	...	...	...	...	...	...
Fed. States of Micronesia <sup>4</sup>	...	...	...	...	...	...	...
Guam <sup>4</sup>	54	53	53	100.0	0.0	0.0	90.6
N. Mariana Islands <sup>4</sup>	75	75	75	100.0	0.0	1.3	94.7
Puerto Rico <sup>4</sup>	174	151	151	100.0	2.0	2.6	90.7
Republic of Palau <sup>4</sup>	...	...	...	...	...	...	...
U.S. Virgin Islands <sup>4</sup>	...	...	...	...	...	...	...

1. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information reported for  $\geq 75\%$  of cases.

2. I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin.

3. Excludes New York City.

4. Not included in U.S. totals.

Ellipses indicate data not available.

Note: Excluding cases with no information on initial drug regimen, 171 (1.1%) were not started on any drugs,

25 (0.2%) were started on one drug, and 1,314 (8.3%) had an initial multidrug regimen other than IR, IRZ, and IRZ,E/S.