Table 10. Percentage of Reported TB Cases by Initial Drug Regimen, Use of Directly Observed Therapy, and Completion of Therapy (COT): United States, 1993-2001

	Initial Drug Regimen ^{1,2}			Directly C	Observed Therapy ³	Therapy ≤1 Year Indicated⁴	
Year	IR	IRZ	IRZ,E/S	DOT Only	Both DOT and Self-Administered	COT <u><</u> 1 Year	СОТ
1993	13.0	31.2	40.9	21.7	14.4	63.6	87.5
1994	7.0	23.3	56.3	28.1	20.5	68.5	87.9
1995	5.2	20.3	63.3	37.2	21.5	72.9	89.6
1996	4.2	17.5	67.9	42.5	22.4	75.6	90.3
1997	3.2	15.1	72.4	46.9	23.8	77.7	91.2
1998	2.6	12.9	74.7	47.7	26.5	80.2	92.4
1999	2.2	11.2	77.2	49.5	27.5	79.9	92.0
2000	2.0	10.4	78.6				
2001	1.6	9.4	78.8	•••	•••		

¹Includes cases in persons alive at diagnosis.

Note: Data for all years updated through April 10, 2002.

See Surveillance Slides #24 and #25.

Table 11. Number and Percentage of Reported TB Cases with HIV Test Results and with HIV Coinfection by Age Group: United States, 1993-2000

	25-44 Years Old				All Ages			
	HIV Test Results ¹		HIV Positive ²		HIV Test Results ¹		HIV Positive ²	
Year	No.	%	No.	%	No.	%	No.	%
1993	4,376	46	2,787	29	7,456	30	3,680	15
1994	4,439	49	2,664	29	7,878	33	3,595	15
1995	4,271	52	2,170	26	8,174	36	3,036	13
1996	4,347	57	1,856	25	8,799	41	2,615	12
1997	4,137	60	1,472	21	8,759	44	2,091	11
1998	3,857	61	1,239	20	8,277	45	1,831	10
1999	3,808	63	1,176	19	8,408	48	1,726	10
2000	3,498	63	952	17	8,042	49	1,457	9

¹Includes cases with positive, negative, or indeterminate HIV test results and cases from California also reported with AIDS. Rhode Island reported HIV test results in 1998, 1999, and 2000. HIV test results were not reported from California. However, California provided HIV status for TB cases reported during 1993-2000 in persons with AIDS (i.e., HIV-positive). Percentages based on all reported TB cases.

Note: Data for all years updated through April 10, 2002.

See Surveillance Slides #22 and #23.

²I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin. Excluding cases with no information on initial drug regimen, 1% were not started on any drugs, less than 1% were started on one drug, and approximately 10% had an initial multidrug regimen other than IR, IRZ, or IRZ,E/S.

³Includes cases in persons alive at diagnosis with initial drug regimen of one or more drugs prescribed.

⁴Includes cases in persons alive at diagnosis, with initial drug regimen of one or more drugs prescribed, who did not die during therapy. Excludes persons with initial isolate resistant to rifampin and pediatric (aged <15) cases with meningeal, bone or joint, or miliary disease. See Technical Notes (Appendix A) for description of COT calculation. Ellipses indicate data not available.

²Includes cases with HIV-positive test results and California cases also reported with AIDS. Percentages based on all reported TB cases.