

Table 31. Tuberculosis Cases by Initial Drug Regimen: 59 Reporting Areas, 2002

Reporting Area	Total Cases	Cases Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percent of Cases with Initial Drug Regimen ^{1,2}		
			No.	%	IR	IRZ	IRZ,E/S
United States	15,075	14,742	14,637	99.3	1.7	8.8	80.2
Alabama	233	224	224	100.0	2.2	60.7	32.1
Alaska	49	49	41	83.7	0.0	2.4	87.8
Arizona	263	253	252	99.6	1.6	6.3	85.3
Arkansas	136	129	127	98.4	20.5	51.2	25.2
California	3,169	3,107	3,104	99.9	1.6	4.2	87.2
Colorado	104	102	102	100.0	0.0	1.0	86.3
Connecticut	104	102	101	99.0	2.0	7.9	82.2
Delaware	25	25	25	100.0	0.0	4.0	88.0
District of Columbia	82	80	79	98.8	0.0	1.3	74.7
Florida	1,086	1,058	1,058	100.0	0.9	9.3	80.0
Georgia	524	512	509	99.4	1.4	5.5	82.1
Hawaii	148	148	138	93.2	6.5	8.7	71.0
Idaho	14	13	12	92.3	0.0	33.3	66.7
Illinois	680	673	673	100.0	1.0	7.4	75.6
Indiana	128	128	128	100.0	0.8	10.2	81.3
Iowa	34	33	33	100.0	3.0	18.2	72.7
Kansas	89	86	86	100.0	3.5	8.1	74.4
Kentucky	146	142	142	100.0	2.1	7.0	83.8
Louisiana	230	227	225	99.1	0.9	12.4	79.6
Maine	23	22	21	95.5	0.0	14.3	76.2
Maryland	306	291	291	100.0	0.0	6.2	89.3
Massachusetts	271	269	267	99.3	0.0	4.9	84.6
Michigan	315	301	301	100.0	4.3	28.2	59.8
Minnesota	237	234	234	100.0	0.0	9.0	81.6
Mississippi	134	129	128	99.2	0.0	17.2	78.1
Missouri	136	130	130	100.0	3.8	6.9	76.2
Montana	12	12	12	100.0	0.0	50.0	50.0
Nebraska	28	27	25	92.6	0.0	28.0	56.0
Nevada	85	79	78	98.7	0.0	1.3	92.3
New Hampshire	19	18	18	100.0	0.0	5.6	94.4
New Jersey	530	524	515	98.3	1.6	7.2	77.7
New Mexico	57	56	52	92.9	1.9	25.0	63.5
New York State ³	350	345	345	100.0	1.4	5.5	86.1
New York City	1,084	1,069	1,069	100.0	0.8	4.3	84.0
North Carolina	434	420	420	100.0	0.0	4.0	86.0
North Dakota	6	6	6	100.0	0.0	0.0	100.0
Ohio	257	252	252	100.0	2.0	21.0	67.9
Oklahoma	190	187	187	100.0	13.9	21.4	47.1
Oregon	111	108	108	100.0	0.0	0.9	93.5
Pennsylvania	353	342	329	96.2	1.2	5.8	81.5
Rhode Island	49	48	47	97.9	0.0	2.1	91.5
South Carolina	256	250	250	100.0	1.2	12.4	77.2
South Dakota	13	12	12	100.0	0.0	58.3	41.7
Tennessee	308	296	296	100.0	1.4	10.1	83.4
Texas	1,550	1,519	1,485	97.8	1.8	7.7	78.4
Utah	31	31	31	100.0	0.0	12.9	83.9
Vermont	8	8	8	100.0	0.0	37.5	62.5
Virginia	315	310	308	99.4	1.3	7.5	86.4
Washington	252	249	248	99.6	0.8	4.4	91.1
West Virginia	30	29	28	96.6	10.7	25.0	57.1
Wisconsin	78	75	74	98.7	1.4	10.8	79.7
Wyoming	3	3	3	100.0	0.0	66.7	33.3
American Samoa ⁴
Fed. States of Micronesia ⁴
Guam ⁴	65	63	62	98.4	0.0	0.0	85.5
N. Mariana Islands ⁴	53	52	52	100.0	0.0	0.0	96.2
Puerto Rico ⁴	129	117	117	100.0	0.0	0.9	94.0
Republic of Palau ⁴
U.S. Virgin Islands ⁴

¹Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information reported for $\geq 75\%$ of cases.

²I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin.

³Excludes New York City.

⁴Not included in U.S. totals.

Ellipses indicate data not available.

Note: Excluding cases with no information on initial drug regimen, 493 (3.3%) were not started on any drugs, 12 (0.08%) were started on one drug, and 1,127 (7.5%) had an initial multidrug regimen other than IR, IRZ, or IRZ,E/S.