

Table 8. Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with No Previous TB by Origin: United States, 1993-2002

Year	Resistant to Isoniazid ¹						Resistant to Isoniazid and Rifampin ¹					
	Total Cases ²		U. S.-born		Foreign-born ³		Total Cases ²		U. S.-born		Foreign-born ³	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1993	1,401	8.4	805	6.8	579	12.4	410	2.5	302	2.6	105	2.3
1994	1,355	8.3	709	6.4	632	12.1	352	2.2	238	2.2	109	2.1
1995	1,172	7.3	554	5.4	617	11.0	252	1.6	168	1.6	84	1.5
1996	1,136	7.4	494	5.2	639	11.3	206	1.3	104	1.1	101	1.8
1997	1,080	7.5	437	5.0	639	11.2	155	1.1	76	0.9	79	1.4
1998	1,012	7.5	366	4.7	644	11.3	131	1.0	55	0.7	75	1.3
1999	904	7.1	284	4.0	618	11.0	128	1.0	39	0.6	89	1.6
2000	893	7.5	270	4.3	620	11.0	121	1.0	38	0.6	83	1.5
2001	806	7.1	245	4.4	560	9.6	116	1.0	34	0.6	82	1.4
2002	778	7.5	197	4.0	574	10.6	121	1.2	37	0.8	83	1.5

¹Isolates may be resistant to other drugs.

²Includes persons of unknown country of birth.

³Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, U.S. Virgin Islands, and U.S. minor and outlying Pacific islands.

Note: Data for all years updated through March 28, 2003.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.

See Surveillance Slides #18, #19, #20, and #21.

Table 9. Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with Previous TB by Origin: United States, 1993-2002

Year	Resistant to Isoniazid ¹						Resistant to Isoniazid and Rifampin ¹					
	Total Cases ²		U. S.-born		Foreign-born ³		Total Cases ²		U. S.-born		Foreign-born ³	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1993	164	16.6	85	12.7	76	25.0	75	7.7	30	4.5	45	15.0
1994	177	17.1	81	11.7	95	28.1	75	7.3	35	5.1	39	11.6
1995	168	17.6	77	13.0	91	25.1	70	7.3	28	4.7	42	11.6
1996	142	16.5	67	12.0	74	24.4	43	5.0	20	3.6	22	7.3
1997	109	14.7	35	7.7	74	25.9	44	5.9	12	2.6	32	11.2
1998	98	13.0	38	7.8	60	22.8	23	3.1	6	1.2	17	6.5
1999	82	12.2	25	6.5	55	19.4	28	4.2	6	1.6	22	7.8
2000	82	13.0	22	6.1	60	22.2	24	3.8	2	0.6	22	8.2
2001	83	13.4	26	8.0	57	19.2	30	4.8	6	1.9	24	8.1
2002	73	13.4	22	7.5	51	20.8	25	4.6	3	1.0	22	8.9

¹Isolates may be resistant to other drugs.

²Includes persons of unknown country of birth.

³Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, U.S. Virgin Islands, and U.S. minor and outlying Pacific islands.

Note: Data for all years updated through March 28, 2003.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.