

**Table 35. Completion of Tuberculosis Therapy (COT): 59 Reporting Areas, 1997**

Reporting Area	Total Cases	Therapy $\leq$ 1 Year Indicated <sup>1</sup>			Therapy $>$ 1 Year Indicated <sup>2</sup>		Overall	
		No. <sup>3</sup>	COT $\leq$ 1 Year(%)	COT(%)	No. <sup>3</sup>	COT(%)	No. <sup>3</sup>	COT(%)
<b>United States</b>	<b>19,753</b>	<b>16,884</b>	<b>77.2</b>	<b>90.0</b>	<b>297</b>	<b>73.7</b>	<b>17,181</b>	<b>89.7</b>
Alabama	405	338	82.5	94.1	4	75.0	342	93.9
Alaska	78	72	91.7	98.6	0	...	72	98.6
Arizona	296	245	75.5	88.6	6	66.7	251	88.0
Arkansas	200	170	86.5	91.8	1	100.0	171	91.8
California	4,044	3,559	76.5	90.8	64	79.7	3,623	90.6
Colorado	93	78	84.6	89.7	4	75.0	82	89.0
Connecticut	128	110	89.1	95.5	1	100.0	111	95.5
Delaware	39	34	85.3	91.2	0	...	34	91.2
District of Columbia	110	88	65.9	85.2	2	50.0	90	84.4
Florida	1,392	1,152	77.7	90.7	22	86.4	1,174	90.6
Georgia	695	597	78.9	93.6	12	66.7	609	93.1
Hawaii	165	146	89.7	93.8	0	...	146	93.8
Idaho	15	15	--	--	0	...	15	--
Illinois	966	806	76.8	88.7	21	81.0	827	88.5
Indiana	168	138	82.6	88.4	0	...	138	88.4
Iowa	74	69	76.8	81.2	0	...	69	81.2
Kansas	78	67	71.6	79.1	1	100.0	68	79.4
Kentucky	184	154	85.1	96.1	0	...	154	96.1
Louisiana	406	341	--	--	6	--	347	--
Maine	21	19	78.9	84.2	0	...	19	84.2
Maryland	339	291	85.2	93.5	1	100.0	292	93.5
Massachusetts	268	242	79.8	90.9	8	100.0	250	91.2
Michigan	372	311	78.1	89.7	2	50.0	313	89.5
Minnesota	161	146	85.6	95.9	2	100.0	148	95.9
Mississippi	245	212	89.6	95.3	1	100.0	213	95.3
Missouri	248	204	84.8	93.6	3	100.0	207	93.7
Montana	18	17	88.2	100.0	0	...	17	100.0
Nebraska	22	17	64.7	76.5	0	...	17	76.5
Nevada	112	90	68.9	78.9	1	100.0	91	79.1
New Hampshire	17	16	75.0	87.5	0	...	16	87.5
New Jersey	718	635	71.3	89.4	14	64.3	649	88.9
New Mexico	72	64	87.5	98.4	0	...	64	98.4
New York State <sup>4</sup>	526	458	69.9	86.9	7	71.4	465	86.7
New York City	1,704	1,403	74.3	86.8	50	40.0	1,453	85.2
North Carolina	463	394	87.1	95.4	6	83.3	400	95.3
North Dakota	12	7	85.7	100.0	0	...	7	100.0
Ohio	286	236	80.9	92.8	1	100.0	237	92.8
Oklahoma	211	174	81.0	92.5	4	100.0	178	92.7
Oregon	161	149	83.9	89.3	2	50.0	151	88.7
Pennsylvania	524	431	74.7	89.3	3	66.7	434	89.2
Rhode Island	40	36	83.3	91.7	0	...	36	91.7
South Carolina	328	275	62.5	86.2	1	100.0	276	86.2
South Dakota	19	18	88.9	100.0	0	...	18	100.0
Tennessee	468	375	83.7	96.3	4	100.0	379	96.3
Texas	1,980	1,711	75.2	89.1	32	84.4	1,743	89.0
Utah	36	33	72.7	97.0	0	...	33	97.0
Vermont	6	5	80.0	100.0	0	...	5	100.0
Virginia	349	301	76.1	86.4	4	50.0	305	85.9
Washington	305	272	82.4	90.8	5	80.0	277	90.6
West Virginia	54	43	72.1	79.1	0	...	43	79.1
Wisconsin	130	118	86.4	97.5	2	100.0	120	97.5
Wyoming	2	2	100.0	100.0	0	...	2	100.0
American Samoa <sup>5</sup>	...	...	...	...	...	...	...	...
Fed. States of Micronesia <sup>5</sup>	...	...	...	...	...	...	...	...
Guam <sup>5</sup>	...	...	...	...	...	...	...	...
N. Mariana Islands <sup>5</sup>	94	88	71.6	71.6	4	25.0	92	69.6
Puerto Rico <sup>5</sup>	257	183	87.4	94.0	5	40.0	188	92.6
Republic of Palau <sup>5</sup>	17	16	--	--	0	...	16	--
U.S. Virgin Islands <sup>5</sup>	...	...	...	...	...	...	...	...

1. Initial isolate susceptible to rifampin (n=12,776) or susceptibility unknown (n=4,802); culture-negative (n=2,673); culture status unknown (n=954); age unknown (n=1).

2. Initial isolate rifampin resistant, or pediatric (aged  $<$ 15) case with meningeal, bone or joint, or miliary disease.

3. Number of cases in persons alive at diagnosis, with initial drug regimen of one or more drugs prescribed, who did not die during therapy. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information on reason therapy stopped for  $\geq$ 90% of cases.

4. Excludes New York City.

5. Not included in U.S. totals.

Ellipses indicate data not available.

**Note:** See Technical Notes (Appendix A) for description of COT calculation.