## Table 7. Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with No Previous TB by Origin: United States, 1993-1999

	Resistant to Isoniazid <sup>1</sup>							Resistant to Isoniazid and Rifampin <sup>1</sup>							
	Total Cases <sup>2</sup>		U. Sborn		Foreign-born <sup>3</sup>		Total C		Cases <sup>2</sup>	U. Sborn		Foreign-born <sup>3</sup>			
Year	No.	%	No.	%	No.	%		No.	%	No.	%	No.	%		
1993	1,400	8.4	804	6.8	579	12.4		410	2.5	302	2.6	105	2.3		
1994	1,355	8.3	709	6.5	632	12.1		352	2.2	238	2.2	109	2.1		
1995	1,171	7.3	554	5.4	615	11.0		252	1.6	168	1.6	84	1.5		
1996	1,134	7.4	492	5.2	638	11.3		205	1.3	103	1.1	101	1.8		
1997	1,076	7.5	429	5.0	640	11.3		157	1.1	75	0.9	81	1.4		
1998	1,012	7.6	366	4.8	644	11.4		127	1.0	52	0.7	74	1.3		
1999	865	7.2	271	4.1	590	11.1		127	1.1	38	0.6	88	1.7		

1. Isolates may be resistant to other drugs.

2. Includes persons of unknown country of birth.

3. Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, U.S. Minor Outlying Islands, U.S. Miscellaneous Pacific Islands, and the U.S. Virgin Islands.

Note: Data for all years updated through May 3, 2000.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.

See Surveillance Slides #14 and #15.

## Table 8. Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with Previous TB by Origin: United States, 1993-1999

	Resistant to Isoniazid <sup>1</sup>							Resistant to Isoniazid and Rifampin <sup>1</sup>						
	Total Cases <sup>2</sup>		U. Sborn		Foreign-born <sup>3</sup>		Total Cas		Cases <sup>2</sup>	U. S.	U. Sborn		Foreign-born <sup>3</sup>	
Year	No.	%	No.	%	No.	%		No.	%	No.	%	No.	%	
1993	164	16.6	85	12.6	76	25.2		75	7.6	30	4.5	45	15.0	
1994	177	17.1	81	11.7	95	28.1		75	7.3	35	5.1	39	11.6	
1995	168	17.6	77	13.0	91	25.1		70	7.3	28	4.7	42	11.6	
1996	141	16.4	67	12.0	73	24.2		42	4.9	20	3.6	21	7.0	
1997	108	14.9	35	8.0	73	25.6		42	5.8	11	2.5	31	10.9	
1998	97	12.8	37	7.5	60	22.6		23	3.0	6	1.2	17	6.4	
1999	80	12.2	27	7.2	51	18.3		25	3.8	6	1.6	19	6.8	

1. Isolates may be resistant to other drugs.

2. Includes persons of unknown country of birth.

3. Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, U.S. Minor Outlying Islands, U.S. Miscellaneous Pacific Islands, and the U.S. Virgin Islands.

Note: Data for all years updated through May 3, 2000.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.