Table 9. Percentage of Reported TB Cases by Initial Drug Regimen, Use of Directly Observed Therapy, and Completion of Therapy (COT): United States, 1993-1999

				Directly (Observed Therapy ³	Therapy	
	Initial Drug Regimen ¹				Both DOT and	≤1 Year Indicated ⁴	
Year	IR	IRZ	IRZ,E/S ²	DOT Only	Self-Administered	COT <1 Year	COT
1993	13.0	31.2	40.9	21.7	14.4	63.5	87.5
1994	7.0	23.3	56.3	28.1	20.5	68.4	87.8
1995	5.2	20.3	63.4	37.2	21.4	72.8	89.5
1996	4.1	17.5	67.9	42.4	22.4	75.5	90.2
1997	3.2	15.1	72.4	46.9	23.7	77.2	90.0
1998	2.6	12.9	74.6		•••	•••	
1999	2.2	11.2	76.8		•••		

^{1.} Includes cases in persons alive at diagnosis.

Note: Data for all years updated through May 3, 2000.

See Surveillance Slides #17 and #18.

Table 10. Number and Percentage of Reported TB Cases with HIV Test Results and with HIV Coinfection by Age Group: United States, 1993-1998

	25-44 Years Old					All Ages			
	HIV Test Results ¹		HIV Positive ²			HIV Test Results ¹		HIV Positive ²	
Year	No.	%	No.	%	No.	%	No.	%	
1993	4,372	46	2,786	29	7,447	30	3,678	15	
1994	4,432	49	2,659	29	7,869	33	3,588	15	
1995	4,266	52	2,167	26	8,168	36	3,031	13	
1996	4,328	57	1,856	25	8,757	41	2,615	12	
1997	4,106	60	1,466	21	8,715	44	2,084	11	
1998	3,837	60	1,239	20	8,229	45	1,824	10	

^{1.} Rhode Island reported HIV test results during 1998. HIV test results were not reported from California. California did provide HIV status for TB cases reported during 1993-1998 in persons also reported with AIDS (i.e., HIV-positive). Includes cases with positive, negative, or indeterminate HIV test results and California cases also reported with AIDS. Percentages based on all reported TB cases.

Note: Data for all years updated through May 3, 2000.

See Surveillance Slide #16.

^{2.} I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin. Excluding cases with no information on initial drug regimen, 1% were not started on any drugs, less than 1% were started on one drug, and approximately 10% had an initial multidrug regimen other than IR, IRZ, and IRZ,E/S.

^{3.} Includes cases in persons alive at diagnosis with initial drug regimen of one or more drugs prescribed.

^{4.} Includes cases in persons alive at diagnosis, with initial drug regimen of one or more drugs prescribed, who did not die during therapy. Excludes persons with initial isolate resistant to rifampin and pediatric (aged <15) cases with meningeal, bone or joint, or miliary disease. See Technical Notes (Appendix A) for description of COT calculation. Ellipses indicate data not available.

^{2.} Includes cases with HIV-positive test results and California cases also reported with AIDS. Percentages based on all reported TB cases.