

Table 10. Number and percent distribution of outpatient department visits with corresponding standard errors, by selected primary diagnosis groups: United States, 2000

Primary diagnosis group and ICD-9-CM code(s) ¹	Number of visits in thousands	Standard error in thousands	Percent distribution	Standard error of percent
All visits	83,289	7,018	100.0	...
Essential hypertension	401	3,591	552	4.3
Routine infant or child health check	V20.2	3,201	456	3.8
Acute upper respiratory infection, excluding pharyngitis	460–461,463–466	2,980	433	3.6
Diabetes mellitus	250	2,633	318	3.2
Malignant neoplasms	140–208,230–234	2,420	441	2.9
Normal pregnancy	V22	2,291	353	2.8
Arthropathies and related disorders	710–719	2,125	409	2.6
Otitis media and eustachian tube disorders	381–382	1,806	278	2.2
Spinal disorders	720–724	1,695	297	2.0
General medical examination	V70	1,654	281	2.0
Follow-up examination	V67	1,478	242	1.8
Rheumatism, excluding back	725–729	1,412	214	1.7
Chronic sinusitis	473	1,239	236	1.5
Heart disease, excluding ischemic	391–392,0,393–398,402,404,415–416,420–429	1,225	301	1.5
Asthma	493	1,036	161	1.2
Psychoses, excluding major depressive disorder	290–295,296.0–296.1,296.4–299	1,026	246	1.2
Acute pharyngitis	462	1,017	165	1.2
Complications of pregnancy, childbirth, and the puerperium	630–677	991	212	1.2
Ischemic heart disease	410–414.9	942	204	1.1
Abdominal pain	789	814	125	1.0
All other diagnoses	47,713	3,969	57.3	1.0

... Category not applicable.

¹Based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) (10). However, certain codes have been combined in this table to describe the utilization of ambulatory care services.

NOTE: Numbers may not add to totals because of rounding.