UNITED STATES SECTION

List of Tables and Figures

United States Section

Asbestosis Mortality

- Figure 1-1. Asbestosis: Number of deaths, crude and age-adjusted mortality rates, U.S. residents age 15 and over, 1968-1992
- Figure 1-2. Asbestosis: Crude mortality rates by state, U.S. residents age 15 and over, 1991-1992
- Figure 1-3. Asbestosis: Median age at death, U. S. residents age 15 and over, 1968-1992
- Table 1-1. Asbestosis: Number of deaths by sex, race, and age, U.S. residents age 15 and over, 1991-1992
- Table 1-2. Asbestosis: Number of deaths by state, U.S. residents age 15 and over, 1968-1992
- Table 1-3. Asbestosis: Mortality rates (per 1,000,000 population) by race and sex, U.S. residents age 15 and over, 1991-992
- Table 1-4. Asbestosis: Years of potential life lost by race and sex, U.S. residents age 15 and over, 1991-1992
- Table 1-5. Asbestosis: Total number of deaths, crude and age-adjusted mortality rates (per 1,000,000 population), and total years of potential life lost (YPLL) to life expectancy, by state, U.S. residents age 15 and over, 1988-1992
- Table 1-6. Asbestosis: Most frequently recorded occupations on death certificate, selected states, U.S. residents age 15 and over, 1991-1992
- Table 1-7. Asbestosis: Most frequently recorded industries on death certificate, selected states, U.S. residents age 15 and over, 1991-1992
- Table 1-8. Asbestosis: Proportionate mortality ratio (PMR) by usual occupation, selected states and years, U.S. residents age 15 and over, 1985-1992
- Table 1-9. Asbestosis: Proportionate mortality ratios (PMR) by usual industry, selected states and years, U.S. residents age 15 and over, 1985-1992
- Figure 1-4. Asbestosis: Total number of deaths by county, U.S. residents age 15 and over, 1983-1992
- Figure 1-5. Asbestosis: Age-adjusted mortality rates by county, U.S. residents age 15 and over, 1983-1992

Asbestos Exposure

- Figure 1-6. Asbestos: Number of MSHA and OSHA inspector samples and percent exceeding the permissible exposure limit (PEL), 1974-1994
- Figure 1-7. Asbestos: Number of MSHA and OSHA inspector samples and average severity levels, 1974-1994
- Table 1-10. Asbestos: Number of MSHA and OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity level, by industry, 1993-1994
- Table 1-11. Asbestos: Number of MSHA and OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity level, by industry, 1993-1994
- Table 1-12. Asbestos: Number of MSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity levels (Avg. Sev.), by state, 1974-1994
- Figure 1-8. Asbestos: Number of MSHA inspector samples collected by state, 1993-1994
- Figure 1-9. Asbestos: Percent of MSHA inspector samples that exceeded the permissible exposure limit (PEL), by state, 1993-1994
- Figure 1-10. Asbestos: Average severity level by state for MSHA inspector samples, 1993-1994
- Figure 1-11. Asbestos: Average severity level by state for MSHA inspector samples that exceeded the permissible exposure limit (PEL), 1993-1994
- Figure 1-12. Asbestos: Number of MSHA inspector samples collected, by county, 1993-1994

- Table 1-13. Asbestos: Number of OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity levels (Avg. Sev.), by state, 1979-1994
- Figure 1-13. Asbestos: Number of OSHA inspector samples collected by state, 1993-1994
- Figure 1-14. Asbestos: Percent of OSHA inspector samples that exceeded the permissible exposure limit (PEL), by state, 1993-1994
- Figure 1-15. Asbestos: Average severity level by state for OSHA inspector samples, 1993-1994
- Figure 1-16. Asbestos: Average severity level by state for OSHA inspector samples that exceeded the permissible exposure limit (PEL), 1993-1994
- Figure 1-17. Asbestos: Number of OSHA inspector samples collected by county, 1993-1994

Coal Workers' Pneumoconiosis Mortality

All tables and figures are analogous to those listed under Asbestosis Mortality.

Coal and Coal Mine Dust Exposure

All tables and figures are analogous to those listed under Asbestos Exposure.

Silicosis Mortality

All tables and figures are analogous to those listed under Asbestosis Mortality.

Silica Exposure

All tables and figures are analogous to those listed under Asbestos Exposure.

Byssinosis Mortality

All tables and figures are analogous to those listed under Asbestosis Mortality.

Cotton Dust Exposure

- Figure 4-6. Cotton dust: Number of OSHA inspector samples and percent exceeding the permissible exposure limit (PEL), 1979-1994
- Figure 4-7. Cotton dust: Number of OSHA inspector samples and average severity levels, 1979-1994
- Table 4-10. Cotton dust: Number of OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity level, by industry, 1993-1994
- Table 4-11. Cotton dust: Number of OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity level, by industry, 1993-1994
- Table 4-12. Cotton dust: Number of OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity levels (Avg. Sev.), by state, 1979-1994
- Figure 4-8. Cotton dust: Number of OSHA inspector samples collected by state, 1993-1994
- Figure 4-9. Cotton dust: Percent of OSHA inspector samples that exceeded the permissible exposure limit (PEL), by state, 1993-1994
- Figure 4-10. Cotton dust: Average severity level by state for OSHA inspector samples, 1993-1994
- Figure 4-11. Cotton dust: Average severity level by state for OSHA inspector samples that exceeded the permissible exposure limit (PEL), 1993-1994
- Figure 4-12. Cotton dust: Number of OSHA inspector samples collected by county, 1993-1994

Unspecified/Other Pneumoconioses Mortality

All tables and figures are analogous to those listed under Asbestosis Mortality.

List of Tables and Figures

All Pneumoconioses Mortality

- Figure 6-1. All pneumoconioses: Number of deaths, crude and age-adjusted mortality rates, U.S. residents age 15 and over, 1968-1992
- Figure 6-2. All pneumoconioses: Crude mortality rates by state, U.S. residents age 15 and over, 1991-1992
- Table 6-1. All pneumoconioses: Number of deaths by state, U.S. residents age 15 and over, 1968-1992
- Table 6-2. All pneumoconioses: Percent of deaths by condition and state, U.S. residents age 15 and over, 1968-1992
- Table 6-3. All pneumoconioses: Total number of deaths, crude and age-adjusted mortality rates (per 1,000,000 population), and total years of potential life lost (YPLL) to life expectancy, by state, U.S. residents age 15 and over, 1988-1992
- Table 6-4. All pneumoconioses: Proportionate mortality ratio (PMR) by usual occupation, selected states and years, U.S. residents age 15 and over, 1985-1992
- Table 6-5. All pneumoconioses: Proportionate mortality ratios (PMR) by usual industry, selected states and years, U.S. residents age 15 and over, 1985-1992
- Figure 6-3. All pneumoconioses: Total number of deaths by county, U.S. residents age 15 and over, 1983-1992
- Figure 6-4. All pneumoconioses: Age-adjusted mortality rates by county, U.S. residents age 15 and over, 1983-1992

Pneumoconiotic Agents Exposure

- Figure 6-5. Pneumoconiotic agents: Number of MSHA and OSHA inspector samples and percent exceeding the permissible exposure limit (PEL), 1970-1994
- Figure 6-6. Pneumoconiotic agents: Number of MSHA and OSHA inspector samples and average severity levels, 1970-1994
- Table 6-6. Pneumoconiotic agents: Number of MSHA and OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity level, by industry, 1993-1994
- Table 6-7. Pneumoconiotic agents: Number of MSHA and OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity level, by industry, 1993-1994
- Table 6-8. Pneumoconiotic agents: Number of MSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity levels (Avg. Sev.), by state, 1970 -1994
- Figure 6-7. Pneumoconiotic agents: Number of MSHA inspector samples collected by state, 1993-1994
- Figure 6-8. Pneumoconiotic agents: Percent of MSHA inspector samples that exceeded the permissible exposure limit (PEL), by state, 1993-1994
- Figure 6-9. Pneumoconiotic agents: Average severity level by state for MSHA inspector samples, 1993-1994
- Figure 6-10. Pneumoconiotic agents: Average severity level by state for MSHA inspector samples that exceeded the permissible exposure limit (PEL), 1993-1994
- Figure 6-11. Pneumoconiotic agents: Number of MSHA inspector samples collected by county, 1993-1994
- Table 6-9. Pneumoconiotic agents: Number of OSHA inspector samples, percent exceeding the permissible exposure limit (PEL) and average severity levels (Avg. Sev.), by state, 1979-1994
- Figure 6-12. Pneumoconiotic agents: Number of OSHA inspector samples collected by state, 1993-1994
- Figure 6-13. Pneumoconiotic agents: Percent of OSHA inspector samples that exceeded the permissible exposure limit (PEL), by state, 1993-1994
- Figure 6-14. Pneumoconiotic agents: Average severity level by state for OSHA inspector samples, 1993-1994
- Figure 6-15. Pneumoconiotic agents: Average severity level by state for OSHA inspector samples that exceeded the permissible exposure limit (PEL), 1993-1994
- Figure 6-16. Pneumoconiotic agents: Number of OSHA inspector samples collected by county, 1993-1994