

Dating of young ground water with CFCs, SF₆, ³H, and ³H/³He:

Examples from the Blue Ridge and Valley and Ridge of parts of Virginia and Pennsylvania

L.N. Plummer, USGS, Reston, VA


“Owing to dispersion, the “age” of a groundwater sample corresponds generally to a time distribution of many elementary flows. Thus, except in the theoretical case of a pure piston flow system, or of stationary waters entrapped in a geological formation, the concept of groundwater age has little significance”

J.-Ch. Fontes (1983) *in* Guidebook on Nuclear Techniques in Hydrology, IAEA



Selected Environmental Tracers 0-50 Year Timescale

tracer1b.grf (JK Böhlke) 4/22/03



Cook and Bohlke, 2000

- ^3H , $^3\text{H}/^3\text{He}$
- ^{85}Kr
- CFC-11, CFC-12, CFC-113
- SF₆
- Event Markers: ^3H , ^{36}Cl , ^{14}C
- Age: time elapsed since recharge


Why measure environmental tracers in ground water?

- Estimate fractions of young water and the mean age of the young fraction in mixtures.
- Evaluate vulnerability to contamination.
- Estimate recharge rates.
- Calibrate models of groundwater flow.
- Estimate rates of geochemical and microbiological processes.
- Retrieve historical records of contaminant loading to aquifers.
- Estimate remediation times.



Approach to “Dating” Young GW

- Collect water samples without contacting air.
- Minimize mixing effects by sampling monitoring wells with narrow screens.
- Analyze with high precision for CFCs, SF₆, ³H, ³H/³He, and others (multi-tracer approach).
- **Age interpretation.** Evaluate multiple tracer data in context of models of groundwater flow.
- **Age is model dependent.**



Age Interpretation


- Comparison of simulated and observed tracer concentrations. **Lumped-parameter models.** (1) multiple tracers from the source, (2) a time series from the source, or (3) multiple tracers from multiple sources in the system. *Choose a model based on hydrogeology.*
- Tracer plots. Method of comparing simulated and observed multiple tracer data; recognizing cases of possible piston flow and binary mixing (dilution); elimination of some mixing models.
- Flow-model calibration and simulation of age.
- All “ages”, regardless of method, are model dependent.

Tritium/Helium-3



- Half-life 12.3 years; decays to ^3He .
- Atmospheric thermonuclear weapons testing from the 1950's, and especially in the period 1962-1963
- Initial ^3H measured
- Terrigenous He sources
- Dispersion around bomb peak
- Confinement of ^3He



Chlorofluorocarbons



New bottle method of collection.
See <http://water.usgs.gov/lab/cfc>

- CFC-12 (CF_2Cl_2), 1930
- CFC-11 (CFCl_3), 1936
- CFC-113 ($\text{C}_2\text{F}_3\text{Cl}_3$), 1944
- Input smooth, increasing until 1990s (dating range ~1950 to early 1990s).
- Stable in aerobic environments.
- In future- dual ages.
- Use of ratios.
- Can detect post 1940's water.
- Collection/analysis not overly labor intensive.
- Problems with contamination, degradation (anoxic), sorption?

Sulfur Hexafluoride




- **Electrical insulator in high voltage switches**
- **First produced 1953**
- **Very low solubility in water.**
- **Does not degrade**
- **Terrigenous source**
- **Dating range: 1970-modern.**
- **Smooth input, increasing in air at 6%/yr; 5 pptv today.**

Dating with CFCs and SF₆

- Henry's law solubility $C_i = K_H \times p_i$

- Requirements:

- Gas-water equilibrium at recharge
- Recharge temperature
- Barometric pressure at recharge
- Knowledge of atmospheric history of the gas



Limitations

$^3\text{H}/^3\text{He}$

- Cost
- Terrigenous He
- Bubbles, Gas-stripping, confinement
- Mixing

SF_6

- Terrigenous SF_6
- Mixing

CFCs


- Contamination
- Degradation
- Mixing

^{85}Kr

- Very difficult collection and analysis
- No labs available to us

There is no perfect tracer. Multi-tracer approach recommended.

Tracer Method of dating the young fraction in binary mixtures of young and old



- CFC pptv ratio defines age of young fraction.
 - % young water $(\text{pptv}_{(\text{measured})} / \text{pptv}_{(\text{ratio year})}) \times 100$
-
- Cannot date outside range for ratio.
 - Cannot use if one of the CFCs in the selected ratio is “contaminated”, even if ratio is “in range”.
 - Ratio-based age must be less than (younger than) apparent (piston flow) model ages for both CFCs in the ratio.

$^3\text{H}/^3\text{He}$ Age Applies to that of the Young Fraction in Simple Binary Mixtures

$^3\text{H}/^3\text{He}$ Age is based on an isotope ratio.

$$t = \frac{1}{\lambda} \cdot \ln \left(1 + \frac{^3\text{He}_{tri}}{^3\text{H}_m} \right)$$

$\lambda = \text{decay constant} = \ln 2 / t_{1/2} = 0.05635 \text{ year}^{-1}$

$t_{1/2} = 12.3 \text{ years}$

Blue Ridge Mountains of Virginia





Plummer, L.N., Busenberg, E., Böhlke, J.K., Nelms, D.L., Michel, R.L., and Schlosser, P., 2001, Groundwater residence times in Shenandoah National Park, Blue Ridge Mountains, Virginia, USA: A multi-tracer approach. *Chemical Geology*, v. 179/1-4, p. 93-111.

- 800 km²; elev. 170-1230 m
- Annual precipitation averaged 114 cm; mean annual temperature, 7.8 °C in high altitude, central part.
- Precambrian to Cambrian fractured crystalline rocks, metabasalts with thin cover of colluvium and residuum.
- Two largest springs have maximum discharges of 20 and 11 l/sec. Most with max. discharge of < 2 l/sec.
- In drought, sprs. about 10% of max.
- wells produce < 6 l/sec, and typically < 1/sec.
- 34 springs, 15 wells: 1996 (wet season), 1997 (dry season).
- Shallow recharge through residuum and colluvium recharge fracture system which has low storage.
- CFCs, ³H/³He, SF₆, ³⁵S, stable isotopes, dissolved N₂, Ar.

SHENANDOAH NATIONAL PARK BIG MEADOWS AREA

Ground-Water-Flow System







Lewis Mountain Spring





Sampling from a Typical Spring Box and Weir














Two ^3H input functions:


- Wash. DC (upper)
- Wash. DC scaled to SNP (lower)

Also 0-2 cc/L of excess air



Average Seasonal
 $\Delta \delta^{18}\text{O} = 0.12$ per mil
Average Age ~ 5 years

- ▲ 3H/3He (Springs)
- SF6 (Springs)
- ◆ CFC-12 (Springs)
- ▲ 3H/3He (Wells)
- SF5 (Wells)
- ◆ CFC-12 (Wells)



Conclusions from Blue Ridge (SNP)

- $d^{18}O$ data: Mean residence time of 5 years. (Range 0-10 yrs.) Age includes uz travel time.
- Ages based on CFCs and SF_6 young; generally consistent with stable isotope data; do not include uz travel time.
- Ages based on $^3H/^3He$ biased young (0-2 yrs, most 0).
- Water from wells have ages of 0 to 25 yrs. $^3H/^3He$ dating works well for these (and applies to the young fraction).
- Some wells discharge mixtures; ratios of CFCs can define age of young fraction and percent young water in mixture.
- Excess CFCs-- Anthropogenic sources? Evidence of shallow recharge?




Chesapeake Bay Watershed

Region of 165,000 km² over parts of NY, PA, MD, DE, VA, and WV.

Evaluate residence times and nitrate transport in groundwater discharging to streams in the watershed.

Assess lag time between changes at the land surface and the response in the base-flow component of groundwater discharge to the bay.

Lindsey and others, 2003, USGS WRIR 03-4035, <http://pa.water.usgs.gov/reports/wrir03-4035.pdf>



The Chesapeake Bay Watershed

64,000 Square Miles of Land, Water, and People

"A Better Bay Through Better Science"


1997

Produced by the USGS from a mosaic of Landsat satellite imagery acquired from 1980-1984




Layered fracture density model (Gburek, Folmar, and Urban (1998))

Depth (ft)




Burton et al. 2002

CFC-113 Concentration in pptv




East Piezometer Transect

--Preferred CFC-12 ages NE of stream are older...
 ..than those underneath and SW of stream
 ..although these ages are really mixture ages!




West Piezometer Transect


--Ages are even more mixed, but % of young water (2nd #) shows older waters generally prevailing to NW
 ...and younger to SE





USGS Chesapeake Bay watershed nutrient study

- SF_6 vs ^3H , CFCs
- Monitoring wells, springs from watersheds











Hydrogeology of Carbonate Terranes of the Valley & Ridge Province



-  Soil & Regolith
-  Alluvium
-  Limestone

-  Younger Ground Water
-  Older Ground Water
-  Mixture of Younger and Older Ground Water



Modified from Brahana and others, 1986


NOT TO SCALE



Tritium vs CFC-12

- Young, piston flow
- Binary mixtures of young and old
- Few CFC-12 contamination






- Valley and Ridge Carbonates**
- △ C-Bay Sprs.
 - Muddy Creek Sprs.
 - Muddy Creek Dom. wells
 - Muddy Creek Mon. wells
 - ◇ Shenandoah Sprs.
 - ◇ Shenandoah wells
 - VAS Sprs.
 - VAS wells

- 2 samples beyond dating range of $^3\text{H}/^3\text{He}$.
- 9 samples may be unmixed.
- 13 samples look like mixtures.





- Valley and Ridge Carbonates**
- △ C-Bay Sprs.
 - Muddy Creek Sprs.
 - Muddy Creek Dom. wells
 - Muddy Creek Mon. wells
 - ◇ Shenandoah Sprs.
 - ◇ Shenandoah wells
 - VAS Sprs.
 - VAS wells


Ages and mixing fractions determined from CFC-113/CFC-12

5-22 years and 10-100% young water in mixture.

Some are inconsistent with ³H (CFC contam.)



Tritium in Precipitation in TU



- Valley and Ridge Carbonates**
- △ C-Bay Sprs.
 - Muddy Creek Sprs.
 - Muddy Creek Dom. wells
 - Muddy Creek Mon. wells
 - ◇ Shenandoah Sprs.
 - ◇ Shenandoah wells
 - VAS Sprs.
 - VAS wells

**$^3\text{H} + ^3\text{He}$ (tritogenic)
as function of $^3\text{H}/^3\text{He}$
age.**

**Numbers are % of
Young fraction in
Mixture.**



Bear Lithia Spring (9/2/99)

Piston-Flow Ages

CFC-11 27.2 yrs

CFC-12 27.2 yrs

CFC-113 >Modern (100 pptv)

$^3\text{H}/^3\text{He}$ 30.0 yrs

Agreement in ages suggests
Piston flow (30 yrs in pipe flow).

Major Contradiction:


Tritium = 1.2 ± 0.2 TU

1969.6 water would contained
about 130 TU; decays to 25 TU



A. B. C. Expectorant warranted purely vegetable—no narcotics.

NATURE'S OWN REMEDY!



Cures KIDNEY and BLADDER Troubles,
URIC ACID, GOUT, and RHEUMATISM,
PHOSPHORIC DEPOSITS,

INFLAMMATION OF THE BLADDER,
DROPSICAL AFFECTIONS,
BRICK-DUST DEPOSITS,

And all forms of DYSPEPSIA which
Arise from a Non-Assimilation of Food,
Such as INDIGESTION, HEARTBURN,

FLATULENCE, SICK HEADACHE
And a Sense of Fullness After Eating.


For further particulars, address

BEAR LITHIA WATER COMPANY,
ELKTON, VIRGINIA.

Price, \$3.50 per Case of one dozen half-gallon bottles, f. o. b., at
Elkton Depot, S. V. R. R., Rockingham county, Va.

AGENTS.—Polk Müller & Co., Richmond; N. Wyatt & Bro.,
Staunton; E. P. Mertz, Washington, D. C.; P. Schever & Co.,
New York; W. H. Douglass, Brooklyn, N. Y.

Routt's Emulsion is used and prescribed by the best physicians.



- Piston Flow
- + Recharge Date
- - Binary Mixing
- . - Exponential Mixing
- Residence Time, Yrs.
- △ C-Bay Springs
- Muddy Creek Sprs.
- Muddy Creek Dom. wells
- Muddy Creek Mon. wells
- ◇ Shenandoah Sprs.
- ◇ Shenandoah Wells
- VAS Sprs.
- VAS Wells

Without multiple Tracers, we would not know this is a Mixture.





Bear Lithia Spring (9/2/99)

Correcting for assumed pre-bomb tritiogenic ^3He in the old fraction, young fraction varies from 5% at 30 yrs to 12% at 0 yrs age.

From CFC-11: 33.0 % 0 yrs water.
From CFC-12: 32.8 % 0 yrs water.

CFCs Imply ^3H should have been around 3 TU instead of 1.2 TU.


CONCLUSION: Discharge 67-88 % pre-bomb water mixed with approx. 12-33% modern water. $^3\text{H}/^3\text{He}$ and CFC age of young fraction is very uncertain (0-30 yrs).




GC-ECD Chromatograms

- Analysis by GC-ECD: Purge and trap gas chromatography with electron-capture detector.
- Detects halogenated VOCs (examples: CFCs, CCl_4 , Halons, TCE, TCA, etc).
- Traditional analysis: GC-MS $< 0.1 \text{ ug/L}$ ($= 100 \text{ ng/L} = 100,000 \text{ pg/L}$)
- GC-ECD: $< 1 \text{ pg/L}$ ($= 5$ orders of magnitude below normal reporting levels)

**Shenandoah National Park
Spring
Drinking Water
Modern Clean Water**




**Spring
Yorktown, VA
Drinking Water**




A Town well in VA
Drinking Water Well
Depth: 190 meters
Valley and Ridge

Relative Concentration




A Town well in VA
Drinking Water Well
Depth: 115 meters
Blue Ridge

Relative Concentration



Aquifer Susceptibility in Virginia, 1998-2000

Nelms, D.L. and others,
2003, USGS
WRIR 03-4278




<http://water.usgs.gov/pubs/of/2003/ofr03-246/ofr03-246.htm>


Shapiro, S.D., Busenberg, E., Focazio, M.J., and Plummer, L.N., 2004,
Historical trends in occurrence and atmospheric inputs of halogenated
volatile organic compounds in untreated ground water used as a source
of drinking water. *Science of the Total Environment*, v. 321, 201-217.

A photograph showing a red drilling rig mounted on a truck in a grassy field. To the left is a white box truck with two red diamond-shaped hazard symbols on its side. To the right is a white SUV. In the background, there is a dense line of bare trees under an overcast sky. A person in a yellow jacket is standing near the base of the drilling rig. A red caution tape is strung across the area.

**Halon 1211
Drilling Tracer**



Well MW-N3D. Large Halon peak. CFC-11 and CFC-12 indicate mid- to late 1970s. Since CFCs came with Halon, water is older than 1970s, but cannot be dated further with CFCs because of contamination with drilling air. Possible mixture of old water and water contaminated with drilling air. Without the Halon data, we would have interpreted a CFC age that is too young.



Well MW-N4D. Trace or no Halon present. CFC data probably unaffected by drilling air. CFC data suggest a modern age for the water. Without the Halon data, we would not know that the CFCs are valid in this sample.

CONCLUSIONS

- Use of multiple tracers and “tracer plots” can help to eliminate some mixing models and refine estimates of mean tracer age.
- To a first approximation, the ages and mixing fractions of many samples from karst or fractured rock can be interpreted using a simple binary mixing model. Is this the result of the sampling process?
- Young ages in Blue Ridge. Mixtures from wells.
- About half of the $^3\text{H}/^3\text{He}$ samples from the Valley and Ridge karst have initial tritium consistent with piston flow (0-15 yrs, unmixed). Rest are mixtures of 0-25 yrs (apparent age) mixed with old (pre-bomb) water.

CONCLUSIONS (cont.)

- **CFC-113/CFC-12 data demonstrate cases of piston flow and binary mixing too, but can be affected by contamination.**
- **Most ground water from fractured rock or karstic aquifers is vulnerable to contamination.**
- **Should include tracers in well drilling.**

CONCLUSIONS (cont.)

“... the concept of groundwater age has little significance” (Fontes, 1983).

Investigation of multiple environmental tracers in groundwater systems can often help to refine the interpretation of age, refine hydrologic concepts, and identify vulnerability to contamination.

Thanks

- **Chesapeake Bay Study.** Scott Phillips, Bruce Lindsey, Gary Spieran, Mike Focazio, J.K. Bohlke, Bill Burton, Colleen Donnelly, Ed Busenberg.
- **Virginia Aquifer Susceptibility Study.** Dave Nelms and George Harlow.
- **Shenandoah National Park Study.** Ed Busenberg, Dave Nelms, Jerry Casile, Julian Wayland, Wandee Kirkland, Stephanie Shapiro, Brian Norton.
- **Reston Chlorofluorocarbon Laboratory.**
- **Noble Gas Laboratory of Lamont-Doherty Earth Observatory, Columbia University.**