

Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida,

July 20-30, 1998



U.S. GEOLOGICAL SURVEY Open-File Report 99-226

Prepared in cooperation with the SOUTH FLORIDA WATER MANAGEMENT DISTRICT

Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998

By Mario Fernandez, Jr., Marci Marot, and Charles Holmes

U.S. Geological Survey Open-File Report 99–226

Prepared in cooperation with the SOUTH FLORIDA WATER MANAGEMENT DISTRICT



Tallahassee, Florida 1999

U.S. DEPARTMENT OF THE INTERIOR BRUCE BABBITT, Secretary

U.S. GEOLOGICAL SURVEY Charles G. Groat, Director

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey

For additional information write to:

District Chief U.S. Geological Survey Suite 3015 227 N. Bronough Street Tallahassee, FL 32301 Copies of this report can be purchased from:

U.S. Geological Survey Branch of Information Services Box 25286 Denver, CO 80225 888-ASK-USGS

CONTENTS

Abstract	1
Introduction	1
Purpose and Scope	3
Previous Studies	3
Description of Study Area	4
Sampling Strategy	5
Sample Collection and Analysis Methods	5
Collection, Preparation, and Preservation of Sediment Samples	5
Identification of Short-Term Sedimentation Patterns Using Beryllium-7 Radiodating	7
Analysis for Sediment Size	8
Scans for Trace Elements	8
Analysis for Total Organic Carbon	8
Immunoassay Analysis for Toxic Organic Compounds	
Laboratory Analysis for Toxic Organic Compounds and Trace Elements	10
Chemical and Physical Characteristics of Selected Bottom Sediments	10
Short-Term Sedimentation Patterns Using Beryllium-7 Radiodating	10
Sediment Size	10
Trace Elements	10
Total Organic Carbon	23
Immunoassay Results for Toxic Organic Compounds	23
Laboratory Results for 10 Selected Sites	27
Toxic Organic Compounds	28
Trace Elements	28
Summary	29
References	30
Appendix—Analytical report for organic compounds and trace elements in 10 selected bottom-sediment	
samples, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	31

Plates [in back pocket]

- 1. Location of previous bottom-sediment sampling sites (1982-1997), and present bottom-sediment sampling sites in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.
- 2. Distribution of beryllium-7 in bottom sediment in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.
- 3. Distribution of trace elements above Natural Range in bottom sediment in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.
- 4. Distribution of total organic carbon in bottom sediment in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.
- 5. Distribution of DDT enzyme-linked immunosorbent assay data in bottom sediment in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.
- 6. Distribution of chlordane enzyme-linked immunosorbent assay data in bottom sediment in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.
- 7. Distribution of carcinogenic polycyclic aromatic hydrocarbons enzyme-linked immunosorbent assay data in bottom sediment in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.

Figures

1. Map showing Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	2
2. Diagram showing sampling and analysis steps for the Caloosahatchee River and Estuary study area,	
Lee County, Florida, July 20-30, 1998	5

3-12. Graphs showing:

	T	
3.	Arsenic concentrations normalized to aluminum concentrations for 30 bottom-sediment samples	
	compared to Florida Department of Environmental Protection regression line and prediction limits,	
	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	16
4.	Cadmium concentrations normalized to aluminum concentrations for 30 bottom-sediment samples	
	compared to Florida Department of Environmental Protection regression line and prediction limits,	
	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	17
5.	Chromium concentrations normalized to aluminum concentrations for 30 bottom-sediment samples	
	compared to Florida Department of Environmental Protection regression line and prediction limits,	
	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	18
6.	Copper concentrations normalized to aluminum concentrations for 30 bottom-sediment samples	
	compared to Florida Department of Environmental Protection regression line and prediction limits,	
	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	19
7.	Nickel concentrations normalized to aluminum concentrations for 30 bottom-sediment samples	
	compared to Florida Department of Environmental Protection regression line and prediction limits,	
	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	20
8.	Lead concentrations normalized to aluminum concentrations for 30 bottom-sediment samples	
	compared to Florida Department of Environmental Protection regression line and prediction limits,	
	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	21
9.	Zinc concentrations normalized to aluminum concentrations for 30 bottom-sediment samples	
	compared to Florida Department of Environmental Protection regression line and prediction limits,	
	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	22
10.	DDT enzyme-linked immunosorbent assay data normalized to total organic carbon, Caloosahatchee	
	River and Estuary study area, Lee County, Florida, July 20-30, 1998	25
11.	Chlordane enzyme-linked immunosorbent assay data normalized to total organic carbon, Caloosahatchee	
	River and Estuary study area, Lee County, Florida, July 20-30, 1998	26
12.	Carcinogenic polycyclic aromatic hydrocarbons enzyme-linked immunosorbent assay data	
	normalized to total organic carbon, Caloosahatchee River and Estuary study area, Lee County, Florida,	
	July 20-30, 1998	26

Tables

1.	Summary of data collected during previous studies of sediment chemical analysis, Caloosahatchee River	-
	and Estuary, and contiguous bays, Lee County, Florida, 1978-88	3
2.	Bottom-sediment sampling sites in the Caloosahatchee River and Estuary study area, Lee County, Florida,	
	July 20-30, 1998	6
3.	Beryllium-7 radioisotope activity on bottom-sediment samples collected in the Caloosahatchee River and	
	Estuary study area, Lee County, Florida, July 20-30, 1998	11
4.	Sediment size analyses on bottom-sediment samples collected in the Caloosahatchee River and Estuary	
	study area, Lee County, Florida, July 20-30, 1998	12
5.	Scans for trace elements on selected bottom-sediment samples collected in the Caloosahatchee River and	
	Estuary, Lee County, Florida, July 20-30, 1998	14
6.	Scans for trace elements on selected bottom-sediment samples collected in the contiguous bays and	
	tributaries to the Caloosahatchee River Estuary, Lee County, Florida, July 20-30, 1998	
7	Laboratory analyses for total organic carbon on selected bottom-sediment samples collected in the	
,.	Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998	23
8	Immunoassay screening on selected bottom-sediment samples collected in the Caloosahatchee River and	
0.	Estuary study area, Lee County, Florida, July 20-30, 1998	24
0	DDT, cyclodienes as chlordane, and carcinogenic polycyclic aromatic hydrocarbons normalized to total	
9.	organic carbon on selected bottom-sediment samples collected in the contiguous bays and tributaries to the	
		25
10	Caloosahatchee River Estuary, Lee County, Florida, July 20-30, 1998	
10.	Laboratory analytical results for pesticides, polychlorinated biphenyls, and polycyclic aromatic	
	hydrocarbons on 10 bottom-sediment samples collected in the Caloosahatchee River Estuary, Lee County,	
	Florida, July 20-30, 1998	27
11.	Laboratory analytical results for trace elements on 10 bottom-sediment samples collected in the	
	Caloosahatchee River Estuary, Lee County, Florida, July 20-30, 1998	28

CONVERSION FACTORS AND ADDITIONAL ABBREVIATIONS

Multiply	Ву	To obtain	
inch (in.)	25.40	millimeter	
foot (ft)	0.3048	meter	
mile (mi)	1.609	kilometer	

- $\mu g/kg =$ micrograms per kilogram
 - < = less than
 - > = greater than
- BDL = below detection limit
- CaPAHs = carcinogenic polycyclic aromatic hydrocarbons
 - CBBI = Charlotte Harbor-Bird Island
- CBFM = Charlotte Harbor–Ft. Myers
 - cm = centimeter
- dpm/g = disintegrations per minute per gram
- EPA = Environmental Protection Agency
- FDEP = Florida Department of Environmental Protection
 - kev = kiloelectron volts
- mg/kg = milligrams per kilogram
- MWP = Mussel Watch Program
- NOAA = National Oceanic and Atmospheric Administration
- NS&T = National Status and Trends
- PAHs = polycyclic aromatic hydrocarbons
- PCBs = polychlorinated biphenyls
- PEL = Probable Effects Level
- ppb = parts per billion
- ppm = parts per million
- QC = quality control
- SFWMD = South Florida Water Management District
 - TEL = Threshold Effects Level
 - TOC = total organic carbon
 - USGS = United States Geological Survey

Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998

By Mario Fernandez, Jr., Marci Marot, and Charles Holmes

Abstract


This report summarizes a reconnaissance study, conducted July 20-30, 1998, of chemical and physical characteristics of recently deposited bottom sediments in the Caloosahatchee River and Estuary. Recently deposited sediments were identified using an isotopic chronometer, Beryllium-7 (⁷Be), a short-lived radioisotope. Fifty-nine sites were sampled in an area that encompasses the Caloosahatchee River (River) about three miles upstream from the Franklin Lock (S-79), the entire tidally affected length of the river (estuary), and the contiguous water bodies of Matlacha Pass, San Carlos Bay, Estero Bay, Tarpon Bay, and Pine Island Sound in Lee County, Florida.

Bottom sediments were sampled for ⁷Be at 59 sites. From the results of the ⁷Be analysis, 30 sites were selected for physical and chemical analysis. Sediments were analyzed for particle size, total organic carbon (TOC), trace elements, and toxic organic compounds, using semiquantitative methods for trace elements and organic compounds. The semiquantitative scans of trace elements indicated that cadmium, copper, lead, and zinc concentrations, when normalized to aluminum, were above the natural background range at 24 of 30 sites. Particle size and TOC were used to characterize sediment deposition patterns and organic content. Pesticides, polychlorinated biphenyls (PCBs), and carcinogenic polycyclic aromatic hydrocarbons (CaPAHs) were determined at 30 sites using immunoassay analysis. The semiquantitative immunoassay analyses of toxic organic compounds indicated that all of the samples contained DDT, cyclodienes as chlordane (pesticides), and CaPAHs. PCBs were not detected.

Based on analyses of the 30 sites, sediments at 10 of these sites were analyzed for selected trace elements and toxic organic compounds, including pesticides, PCBs, and PAHs, using quantitative laboratory procedures. No arsenic or cadmium was detected. Zinc was detected at two sites with concentrations greater than the lower limit of the range of sediment contaminant concentrations that are usually or always associated with adverse effects (Florida Department of Environmental Protection's Sediment Quality Assessment Guidelines). Organochlorine pesticides were detected at four sites at concentrations below the reporting limits; there were no organophosphorus pesticides or PCBs detected. PAHs were detected at eight sites; however, only four sites had concentrations above the reporting limit.

INTRODUCTION

The South Florida Water Management District (District) is developing a Caloosahatchee River Water Management Plan to address environmental conditions and water-supply needs of the Caloosahatchee watershed. As part of this plan, the District will evaluate nitrogen, phosphorus, and potential toxic trace elements in the sediments in the Caloosahatchee River and Estuary, and the contiguous water bodies of Matlacha Pass, San Carlos Bay, Estero Bay, Tarpon Bay, and Pine Island Sound, (fig. 1), in Lee County, Florida (study area). As part of this plan, the USGS in July 1998, in cooperation with the District, conducted a reconnaissance of chemical and physical characteristics of selected bottom-sediment samples in the study area for anthropogenic organic compounds and trace elements. The anthropogenic compounds are referenced in this study as toxic organic compounds and include organochlorine pesticides, organophosphorus pesticides, polychlorinated biphenyls (PCBs), and polycyclic aromatic hydrocarbons (PAHs) which include the carcinogenic PAHs (CaPAHs). The anthropogenic trace elements include arsenic, cadmium, chromium, copper, lead, mercury, nickel, and zinc.





2 Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998

Purpose and Scope

The purpose of this report is to summarize the findings of the USGS reconnaissance survey of bottom sediments in the Caloosahatchee River and Estuary, tributaries, and contiguous bays conducted on July 20-30, 1998.

The USGS reconnaissance survey consisted of the following activities:

- Determination of the chemical and physical characteristics of 60 selected bottom sediment-sampling sites including 2 sites above the Franklin Lock (S-79), and
- Selection of 10 sampling sites for analysis of toxic organic compounds and selected trace elements. The 10 sites were selected from the 60 bottom sediment-sampling sites.

A technical advisory group, consisting of USGS and South Florida Water Management District personnel, selected the initial 60 sampling sites. Fifty-eight sites were located in the estuary, tributaries, and the bays; and two sites were located upstream of the Franklin Lock.

Previous Studies

Several Federal and State agencies have previously conducted sediment chemistry investigations of the Caloosahatchee River and Estuary. The National Oceanic and Atmospheric Administration (NOAA) has monitored spatial and temporal trends of chemical contamination and biological responses to contamination in their National Status and Trends Program (NS&T). NOAA's Mussel Watch Program (MWP) has monitored bottom sediments at Charlotte Harbor-Bird Island (CBBI) (at the mouth of the Caloosahatchee River) in 1986 and 1987, and Charlotte Harbor-Ft. Myers (CBFM) in 1988 (table 1, plate 1). The constituents analyzed in the sediments included chlorinated pesticides, PCBs, PAHs, total organic carbon (TOC), and selected trace elements. Sediment-size distribution was also determined.

The Florida Department of Environmental Protection (FDEP) published the "Florida Coastal Sediment Contaminants Atlas" in 1994 which presented the quality of the coastal sediments for five sites in the Caloosahatchee Estuary, one site in San Carlos Bay, and one site in Estero Bay (table 1). The atlas presents summarized data and interpretations for the distribution of pesticides, PCBs, PAHs, TOC, total nitrogen, total phosphorus, and trace elements.

The FDEP further developed a statistical/graphical approach to differentiate concentrations of trace elements present in sediment samples from varied natural background concentrations (FDEP, 1988).

 Table 1.
 Summary of data collected during previous studies of sediment chemical analysis, Caloosahatchee River and Estuary, and contiguous bays, Lee County, Florida, 1978-88

[PAH, polycyclic aromatic hydrocarbons; PCBs, polychlorinated biphenyls; TOC, total organic carbon; TKN, total Kjedhal nitrogen; TP, total phosphorus; X, analyzed; -, not analyzed]

			Latituda				_		Constit	uents	analyze	əd			
Site	Sam- pling	Latitude			Longitude		Organic compounds					Trace ele-	Sedi- ment		
number	date	Degrees	Minutes	Seconds	Degrees	Degrees Minutes S		Pesti- cides	PCBs	PAH	тос	TKN	N TP	ments	size
CLR-00011	8/28/85	26	41	47	81	47	52	х	Х	Х	х	Х	Х	х	-
CLR-00021	8/28/85	26	36	39	81	53	50	Х	Х	Х	Х	Х	Х	Х	-
CLR-00031	8/28/85	26	36	7	81	54	5	Х	Х	Х	Х	Х	Х	Х	-
CLR-00041	8/28/85	26	32	8	81	56	48	Х	Х	Х	Х	Х	Х	Х	-
CLR-00051	8/28/85	26	31	43	81	58	26	Х	Х	Х	Х	Х	Х	Х	-
SCB-00011	8/28/85	26	30	32	82	1	4	Х	Х	Х	Х	Х	Х	Х	-
EST-00011	9/23/86	26	24	49	81	52	0	Х	Х	Х	Х	Х	Х	Х	-
CBBI ²	1986-87	26	30	52	82	2	4	Х	Х	Х	Х	-	-	Х	Х
CBFM ²	1988	26	33	30	81	55	22	Х	Х	Х	Х	-	-	Х	Х
S-79 ³	1978	26	43	25	81	41	55	-	Х	-	-	-	-	Х	-
CH-27 ⁴	1982	26	31	46	81	59	52	Х	Х	Х	-	Х	Х	Х	Х

¹Seal, Thomas, Florida Department of Environmental Protection, written commun.,1999. (CLR, Caloosahatchee River; SCB, San Carlos Bay; EST, Estero Bay).

²Cantillo, Adriana, National Oceanic and Atmospheric Administration, National Status and Trends, written commun., 1997; (CBBI, Charlotte Harbor-Bird Island; CBFM, Charlotte Island-Ft. Myers).

³La Rose, H.R., and McPherson, B.F.,1983, Chemical and hydrologic assessment of the Caloosahatchee River Basin, Lake Okeechobee to Franklin Lock, Florida: U.S. Geological Survey, Water-Resources Investigations Report 83-4126.

⁴Stoker, Y.E., 1986, Water quality of the Charlotte Harbor estuarine system, Florida, November 1982 through October 1984: U.S. Geological Survey Open-File Report 85-563.

The statistical approach was based on the normalization of trace elements concentrations to aluminum concentrations in sediment samples. The FDEP stated that regression lines and the 95 percent confidence limits resulting from normalizing the data provided a valid statistical estimate of the range of values to be expected from samples collected from clean sediments (Natural Range) in Florida (FDEP, 1988, p. 29). Data points plotting above the upper 95 percent confidence limit corresponded to "Above Natural Range (enrichment)"; data points plotting below the lower 95 percent confidence limit corresponded to "Below Natural Range (possible analytical error)". A detailed discussion about the development of this statistical method is described in "A guide to the interpretation of metal concentrations in estuarine sediments" (FDEP, 1988).

The FDEP developed effects-based sediment quality assessment guidelines (SQAGs) as a costeffective approach for screening chemical levels to estimate potential adverse biological effects (FDEP, 1994). The SQAGs may be used to:

- Assess the adverse biological effects that could, potentially, be associated with levels of sediment-associated contaminants.
- Evaluate existing sediment chemistry data, and rank areas of concern and chemicals of concern in terms of their potential to be associated with adverse biological effects.
- Evaluate existing data to determine if additional testing is needed to support regulatory decisions.

Three distinct ranges of contaminant concentrations were defined in the FDEP (1994) report: 1) a minimal effects range, 2) a possible effects range, and 3) a probable effects range. The upper limit of the minimal effects range was represented by the Threshold Effects Level (TEL). The TEL represents the upper limit of the range of sediment contaminant concentrations dominated by no adverse biological effects. The Probable Effects Level (PEL) represents the lower limit of the range of contaminant concentrations that are usually or always associated with adverse biological effects. The TEL and the PEL bracket the range of possible effects; data falling within this range suggest additional investigations.

Normalization of toxic organic compounds to TOC was used by the FDEP (1994) to account for the influence of organic carbon on the bioavailability of the organic compounds, and therefore the potential for toxicity. High TOC concentrations will tend to reduce the bioavailability of toxic organic compounds by effectively sequestering them in the sediments (FDEP, 1994). Two USGS reports contain information about chemical constituents in bottom sediments of the Caloosahatchee River and Estuary. LaRose and McPherson (1983) collected samples from four sites upstream of the Franklin Lock (S-79) and Stoker (1986) collected samples from five sites across a transect at the mouth of the estuary at Shell Point (table 1, plate 1).

Information about discharges that may affect the water quality of the Caloosahatchee watershed can be found at the Environmental Protection Agency's Index of Watershed Indicators web site (**www.epa.gov**/**surf2**). The web site contains environmental uses and discharges (provided by EPA Envirofacts) and facilities regulated by EPA including hazardous waste facilities (Resources Conservation Recovery Act – RCRA); superfund sites (Comprehensive Environmental Response, Compensation, and Liability Act – CERCLA); and toxic releases (Toxic Release inventory – TRI).

DESCRIPTION OF STUDY AREA

The study area encompasses a 1.3-mile length of the Caloosahatchee River upstream from the Franklin Lock (S-79) (River), the entire tidally affected length of the river (estuary) and the contiguous water bodies of Matlacha Pass, San Carlos Bay, Estero Bay, Tarpon Bay, and Pine Island Sound (plate 1). Boundaries of the contiguous water bodies are based on the delineation reported by Goodwin (1991). The estuary has 16 tributaries and two major cities, Cape Coral and Fort Myers. Cape Coral is on the west bank, and Fort Myers is on the east bank. The city of Cape Coral, with a population of approximately 84,000 (Pierce, 1995), has about 400 miles (mi) of canals that honeycomb the entire city. Of the 400 mi, about 130 mi are tidally affected (Goodwin, 1991). The city of Fort Myers has a population of approximately 46,000 (Pierce, 1995), and does not have a canal system like Cape Coral. Runoff from areas of Cape Coral, Fort Myers, and some unincorporated areas of Lee County discharge through stormwater conveyances into the Caloosahatchee River estuary (Tony Pellicer, Natural Resource Manager, Lee County, Florida, verbal commun., 1999).

SAMPLING STRATEGY

The sampling strategy for the study was developed with the cooperation of a technical advisory group assembled by the District. The strategy specified the collection of approximately the top 1-centimeter (cm) of soft bottom-sediment samples at 60 selected sites and the physical and chemical characterization of the sediment samples (excluding rocks, sand, gravel, and vegetation). A sampling and analysis step diagram is presented in figure 2.




Figure 2. Sampling and analysis steps for the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.

The characterization included short-term sediment accumulation patterns (determined from analyses of Beryllium-7 (⁷Be), sediment-size distributions, concentrations of trace elements, total organic carbon, and selected toxic organic compounds. Locations of the 60 sites and number of samples collected are listed below:

- *Caloosahatchee River*: 32 samples in the estuary including 2 samples upstream of the Franklin Lock (fig. 1),
- Matlacha Pass: 3 samples,
- San Carlos Bay: 13 samples,
- Estero Bay: 8 samples,
- Tarpon Bay: 2 samples, and
- Pine Island Sound: 2 samples.

Fifty-nine of the 60 selected sites were sampled; one site, located in San Carlos Bay, was not sampled because the bottom sediments consisted of sand within a 1-mile radius. Initially, samples from the 59 sites would be analyzed for ⁷Be to determine short-term sedimentation patterns. Samples from 30 sites with the most recent sedimentation based on the findings of the ⁷Be survey were selected for sediment-size distribution, scans for trace elements, total organic carbon, and semiquantitative analysis of toxic organic compounds.

Subsequently, based on the analytical findings for the 30 selected sites, samples from the 10 sites with the greatest percentage of fine sediments and concentrations of chemical constituents were analyzed by a contract laboratory. The 10 samples sent to the laboratory were analyzed for organochlorine and organophosphorus pesticides, PCBs, PAHs, and selected trace elements.

SAMPLE COLLECTION AND ANALYSIS METHODS

This section describes the sediment collection method and analytical methods used in this report.

Collection, Preparation, and Preservation of Sediment Samples

The descriptive information (site number, latitude and longitude, name of surface-water body, distance downstream from the Franklin Lock (S-79), date sampled, and depth to the top of the sediment layer) about the sediment-sampling sites in the study area is presented in table 2. Visual, mechanical, and depth-sounding methods were used to locate areas of recent sediment deposition (less than 6 months).
 Table 2.
 Bottom-sediment sampling sites in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998

[Site number and site location same as previous study except when noted by (n): (n), previous site is near the site sampled in this report; Previously reported, previously reported data collected by C (Crean and Chamberlain (1985), or H (Haunert, 1978); ---, not previously sampled; (miles), miles upstream from the confluence with the Caloosahatchee River Estuary; n, near the site sampled in this study; na, not applicable]

Site number		Latitude		Longitude			Distance		Depth to
This report	Previous report	Degrees	Minutes	Degrees	Minutes	- Surface- water body	down- stream ¹ (in miles)	Date sampled	sediments from surface (in feet)
1		26	42 279	01	41.060	Caloosahatchee River ²	1.1	07/29/09	24.0
1		26 26	43.278	81	41.060	Caloosahatchee River ²	-1.1	07/28/98	24.0 6.7
2	 C-156 ⁴	26 26	43.375	81	40.870		-1.3	07/28/98	
3		26 26	43.576	81	42.650	Caloosahatchee River Estuary	0.7	07/20/98	9.5
4	C-159 H-6 ⁵	26 26	43.061	81	45.045	Caloosahatchee River Estuary	3.1	07/20/98	8.3
5		26 26	42.154	81	46.504	Caloosahatchee River Estuary	5.4	07/20/98	11.2
6	C-163	26 26	41.860	81	47.389	Caloosahatchee River Estuary	6.4	07/21/98	9.5
7		26 26	41.365	81	45.993	Orange River (3.5 miles)	6.8	07/21/98	11.0
8	C-165	26 26	41.793	81	47.764	Caloosahatchee River Estuary	6.8	07/21/98	3.5
9		26 26	42.115	81	49.876	Daughtrey Creek (0.6 mile)	9.6	07/30/98	4.2
10	C-167	26	41.948	81	48.659	Caloosahatchee River Estuary	8.1	07/21/98	5.2
11	C-169(n)	26	41.417	81	49.722	Caloosahatchee River Estuary	8.8	07/20/98	3.1
12	C-170	26	40.853	81	49.689	Caloosahatchee River Estuary	9.3	07/22/98	6.3
13	C-171	26	40.986	81	49.395	Caloosahatchee River Estuary	8.9	07/21/98	7.5
14	C-173(n)	26	0.637	81	50.989	Caloosahatchee River Estuary	10.5	07/22/98	5.3
15	C-174	26	40.326	81	51.479	Caloosahatchee River Estuary	10.3	07/22/98	6.5
16		26	39.477	81	51.631	Caloosahatchee River Estuary	11.9	07/22/98	6.5
17		26	39.171	81	50.720	Billy Creek (0.9 mile)	11.3	07/22/98	2.9
18		26	39.176	81	51.564	Caloosahatchee River Estuary	12.2	07/22/98	7.1
19		26	38.739	81	52.402	Caloosahatchee River Estuary	13.0	07/22/98	9.1
20	C-228	26	39.766	81	52.415	Caloosahatchee River Estuary	12.1	07/22/98	5.1
21		26	39.453	81	53.722	Yellow Fever Creek (0.1 mile)	13.7	07/22/98	9.5
22		26	38.548	81	53.756	Caloosahatchee River Estuary	14.2	07/22/98	7.0
23	C-227	26	37.678	81	53.502	Caloosahatchee River Estuary	14.9	07/22/98	8.1
24		26	36.130	81	51.874	Caloosahatchee River Estuary	17.0	07/23/98	3.9
25		26	36.068	81	53.667	Caloosahatchee River Estuary	16.7	07/23/98	2.2
26		26	35.457	81	54.145	Caloosahatchee River Estuary	17.6	07/23/98	9.6
27		26	32.620	81	55.79000	Caloosahatchee River Estuary	21.2	07/23/98	7.2
28	C-218(n)	26	32.218	81	56.29200	Caloosahatchee River Estuary	21.8	07/30/98	2.0
29	C-214(n)	26	31.780	81	58.78600	Caloosahatchee River Estuary	24.3	07/29/98	14.0
30	C-211	26	31.381	82	0.3090	San Carlos Bay	na	07/23/98	5.0
31		26	32.194	81	59.91600	Caloosahatchee River Estuary	26.7	07/24/98	15.8
32		26	0.509	82	1.611	San Carlos Bay	na	07/23/98	5.3
33		26	30.238	82	1.042	San Carlos Bay	na	07/30/98	5.5
34		26	29.910	82	1.027	San Carlos Bay	na	07/23/98	2.7
35		26	29.630	82	0.137	San Carlos Bay	na	07/30/98	1.5
36			Not sat	mpled ⁶			na		
37	C-192(n)	26	28.081	82	2.656	San Carlos Bay	na	07/29/98	21.0
38	C-188	26	29.404	82	5.221	San Carlos Bay	na	07/29/98	4.5
39		26	29.964	82	2.543	San Carlos Bay	na	07/29/98	4.4
40	C-191(n)	26	29.563	82	2.259	San Carlos Bay	na	07/29/98	3.1
40A		26	29.056	82	2.259	San Carlos Bay	na	07/29/98	10.7
41		26	29.056	82	2.031	San Carlos Bay	na	07/23/98	4.5
42		26	30.358	82	2.669	San Carlos Bay	na	07/23/98	2.7
						-			

6 Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998 **Table 2.** Bottom-sediment sampling sites in the Caloosahatchee River and Estuary study area, Lee County, Florida,July 20-30, 1998 (Continued)

[Site number and site location same as previous study except when noted by (n): (n), previous site is near the site sampled in this report; Previously reported, previously reported data collected by C (Crean and Chamberlain (1985), or H (Haunert, 1978); ---, not previously sampled; (miles), miles upstream from the confluence with the Caloosahatchee River Estuary; n, near the site sampled in this study; na, not applicable]

Site number		number Latitude		Longitude			Distance		Depth to
This report	Previous report	Degrees	Minutes	Degrees	Minutes	- Surface- water body	down- stream ¹ (in miles)	Date sampled	sediments from surface (in feet)
43	C-180	26	30.860	82	3.648	San Carlos Bay	na	07/29/98	5.3
44	C-175	26	31.693	82	3.465	Matlacha Pass	na	07/29/98	8.5
45	C-177(n)	26	31.712	82	2.349	Matlacha Pass	na	07/23/98	1.3
46		26	31.955	82	4.472	Matlacha Pass	na	07/29/98	1.5
47		26	27.927	82	4.993	Tarpon Bay	na	07/23/98	2.9
48		26	27.485	82	4.436	Tarpon Bay	na	07/23/98	8.0
49		26	28.589	82	7.889	Pine Island Sound	na	07/29/98	2.0
50		26	29.986	82	7.284	Pine Island Sound	na	07/29/98	5.5
51		26	27.310	81	57.011	Estero Bay	na	07/29/98	10.0
52		26	27.912	81	51.965	Estero Bay	na	07/28/98	4.1
53		26	27.569	81	52.223	Estero Bay	na	07/28/98	3.5
54		26	25.566	81	52.635	Estero Bay	na	07/24/98	6.3
55		26	24.014	81	51.176	Estero Bay	na	07/24/98	6.5
56		26	22.115	81	51.003	Estero Bay	na	07/28/98	11.0
57		26	21.103	81	50.964	Estero Bay	na	07/24/98	4.7
58		26	39.061	81	51.246	Billy Creek (0.4 mile)	12.1	07/22/98	1.9
59		26	26.337	81	53.866	Estero Bay	na	07/24/98	3.7
60		26	32.635	81	58.486	Caloosahatchee River Estuary	23.8	07/30/98	10.4

¹Locations of bottom-sediment sampling sites in the estuary are referenced to midstream distances in miles downstream from the Franklin Lock (S-79).

²Above Franklin Lock (S-79).

³Oxbow above Franklin Lock (S-79).

⁴Site number for data collected by Crean and Chamberlain in 1985.

⁵Site number for data collected by Haunert in 1978.

⁶Not sampled, bottom sediments consisted of sand within a 1-mile radius.

A Rockwell geographical positioning system (GPS) was used from the sampling boat to record latitude and longitude for each sampling site. Sediment samples were collected from approximately the top 1-cm at each site using an Eckman dredge with a messenger for soft sediments (rocks, sand, gravel, and vegetation were avoided). The samples collected for each site were combined and sealed in plastic bags, labeled, iced in the field, and stored in a laboratory freezer for processing. In preparation for analysis, the 59 frozen combined sediment samples were subdivided in the USGS laboratory with a heated nickel-chromium wire. One set of samples was thawed at room temperature; the

other set of samples was kept frozen for future immunoassay and laboratory analysis. The thawed subset was divided into two parts: one part was dried in an oven at 60 °C for radioisotopes and trace elements, and the other part was analyzed for sediment-size analysis.

Identification of Short-Term Sedimentation Patterns Using Beryllium-7 Radiodating

The relative activity of ⁷Be in the surface layer of the bottom sediments was used to identify the location of recently deposited material. ⁷Be is a naturally produced radioisotope that is formed by cosmic ray bombardment of atmospheric nitrogen (N) and oxygen (O). The ⁷Be is transferred from the atmosphere to earth through precipitation (Geyh and Schleicher, 1990). This radioisotope is a highly reactive element that becomes rapidly and tightly associated with sedimentary substrata. ⁷Be has a half-life of 53 days, it can be used as a tracer of sedimentation that has occurred within approximately the past 6 months. Thus, detection of ⁷Be in the sediment indicates that the sediment and sedimentation patterns have been present for less than 6 months (Charles Holmes, USGS, verbal commun., 1998).

The 59 thawed samples were dried in an oven at 60°C and then ground finely using a mortar and pestle. Fifty grams were sealed in counting jars for gamma emission detection. Gamma emissions were measured using a Canberra high-purity germanium detector with a PC-based multichannel analyzer. Each sample was counted for a minimum of 24 hours, and the ⁷Be activity was calculated from the net area of the 477 kev (kiloelectron volts) photopeak. The ⁷Be activity was decay-corrected to the date of sampling using the following equation:

where,

A is the activity, decay-corrected to the date of sampling,

 $A = A_0 e^{-\lambda t}$

- A_0 is the measured activity,
- λ is the ⁷Be decay constant, and
- t is the time since collection.

The results were reported as ⁷Be disintegrations per minute per gram of sample (dpm/g) and statistical counting error.

Analysis for Sediment Size

The purpose of collecting bottom sediment for size analysis was to determine the distribution of fine sediments (silt and clays) because the amount of silt and clay-sized particles impacts the chemical properties of the sediments. Size analyses were performed on 30 sediment samples to determine the distribution of fine sediments (silts and clays). Potential sources of sediment particles for an estuary include rivers, net sediment flux from the marine environment, overland runoff, and anthropogenic point sources (Schoellhamer, 1991). Fine sediments can be resuspended into the water column, adsorb organic compounds and trace elements that may be present, and then become a vehicle for transport and redeposition of the sorbed organic compounds and trace elements.

A wet sieve analysis was performed on the samples as described by Matthes (1992). The analysis included 8 mesh sizes: >4, >2, >1, >0. 5, >0.25, >0.125, >0.0625, and <0.0625 millimeters. The sieved fractions were dried in an oven at 60° C, cooled, weighed, and the percentage of each fraction determined.

Scans for Trace Elements

Scans for trace elements were performed by a USGS contract laboratory on 30 sediment samples for 30 trace elements including seven trace elements often associated with anthropogenic activities. The selected trace elements included arsenic, cadmium, chromium, copper, lead, nickel, and zinc.

The samples were dried and then digested in a 4-acid digestion (HNO_3 - $HClO_4$ -HF-HCl) procedure. Digested samples were analyzed for 30 trace elements by an inductive coupled plasma/atomic emission spectroscopy analytical method. The 30 trace elements analyzed are listed below:

Aluminum	Iron	Silver
Antimony	Lead	Sodium
Arsenic	Magnesium	Strontium
Barium	Manganese	Tin
Bismuth	Molybdenum	Titanium
Cadmium	Nickel	Tungsten
Calcium	Phosphorus	Vanadium
Chromium	Potassium	Yttrium
Cobalt	Scandium	Zinc
Copper	Selenium	Zirconium

Analysis for Total Organic Carbon

Thirty sediment samples were sent to a contract laboratory for TOC analysis using EPA Method SW-9040/E415.1. TOC was used to identify the distribution of the total organic constituents in the sediments and to normalize the toxic organic compounds. The total organic constituents included those that are naturally occurring organic compounds (marine life residue and tannins from vegetation decay) and anthropogenic organic (toxic) compounds.

(1)

Immunoassay Analysis for Toxic Organic Compounds

Immunoassay technology was used at the USGS North Carolina District's laboratory to semiquantitatively identify anthropogenic organic compounds in 30 samples. This technology is based on biologically generated "immunoglobulin" proteins (antibodies) reacting with specific target compounds (antigens). Highly specific antibodies for target compounds, such as chlorinated pesticides, PCBs, and carcinogenic PAHs (CaPAHs) have been developed for use in environmental immunoassays. The antigen-antibody complexes of specific target compounds provided a high degree of analytical specificity.

Competitive enzyme-linked immunosorbent assay (ELISA) was used in this study (Strategic Diagnostics Inc., 1998). In competitive ELISAs, the unlabeled analyte (liquid extract of the sediment sample) is combined with a labeled enzyme analog of the analyte and the analyte specific antibody. Both the unlabeled and the enzyme-labeled (Horse Radish Peroxidase) analytes competed for a limited number of antibody sites, and bonded to antibodies in direct proportions to their relative concentrations in the reactive mixture. After an incubation period, the antigen-antibody complexes were separated from the unbound substances by decanting the liquid. Reagents were added to the complexes and a color developed which was read in a spectrophotometer. The color was proportional to the enzyme-labeled analyte and inversely proportional to the concentration of the sample analyte. The targeted toxic organic compounds, chlorinated pesticides, and PCBs were analyzed using the ELISA competitive method; CaPAHs were analyzed by ELISA/paramagnetic particle analysis.

In the paramagnetic particle analysis procedure, an enzyme-labeled CaPAH (enzyme conjugate) was added to the sample followed by paramagnetic particles with antibodies specific to CaPAHs. Both the sample and the enzyme conjugate competed for antibody sites on the magnetic particles. At the end of an incubation period, a magnetic field was applied to hold the paramagnetic particles that have the sample's CaPAH and the enzyme conjugate bound to the antibodies on the particle. The sample's CaPAH and the enzyme conjugate were bound in proportion to their original concentration in the tube. After the attached particles were washed, an enzyme substrate and chromogen was added and a color was developed during an incubation period. Because the enzyme conjugate was in competition with the unlabeled CaPAHs in the sample for antibody sites on the paramagnetic particles, the color was, therefore, inversely proportional to the concentrations of CaPAHs in the sample.

As with all ELISA tests, cross-reactivity needs to be considered. Because more than one compound will react with the ELISA test, a positive result only indicates the presence of one or more compounds that are listed with the individual kits.

The DDT immunoassay test does not differntiate between DDT and other organochlorine compounds. The following organochlorine compounds yield positive results with this assay:

p,p'-DDT	DDA
p,p'-DDD	Chloropropylate
p,p'-DDE	Chlorobenzilate
o,p'-DDT	Dicofol
o,p'-DDD	Tetradifon
o.p'-DDE	

The calibrating standard used for the DDT assay was p,p'-DDT. The standard concentrations were 0.2, 1.0, and 10 parts per million (ppm). Two sample extracts were analyzed in duplicate for quality control (QC).

The cyclodiene immunoassay test does not differentiate between chlordane and other cyclodienes. The following cyclodiene compounds yield positive results with this assay:

Chlordane	Dieldrin
Aldrin	Heptachlor
Endosulfan	Toxaphene
Endrin	

The calibrating standard used for the cyclodiene assay was chlordane. The calibrating standard concentrations were 20, 100, and 590 parts per billion (ppb) (0.02, 0.1 and 0.59 ppm). Two sample extracts were analyzed in duplicate for QC.

The PCBs (Aroclors) immunoassay test does not differentiate between various Aroclors. The following Aroclor compounds yield positive results with this assay:

Aroclor 1016	Aroclor 1254
Aroclor 1221	Aroclor 1260
Aroclor 1232	Aroclor 1268
Aroclor 1242	Aroclor 1282
Aroclor 1248	

The calibrating standard used for the PCBs was Aroclor 1248. The calibrating standard concentrations were 1, 5, 10, and 50 ppm. Two sample extracts were analyzed in duplicate for QC.

The CaPAHs immunoassay test does not differentiate between various CaPAHs of the pyrene series, selected PAHs, creosote, and fuels. The following CaPAH compounds and mixtures yield positive results with this assay:

CaPAH Compounds

Acenaphthalene Acenaphthylene Anthracene Benzo[a]anthracene Benzo[k]fluoranthene Benzo[b]fluoranthene

Benzo[a]pyreneFluoreneBenzo[g,h,i]peryleneIndeno[1,2,3-c,d]pyreneChryseneNaphthleneDibenzo[a,h]anthraceePhenanthreneFluoranthenePyrene

CaPAH Mixtures

Cresote Diesel Fuel Fuel Oil #4 Fuel Oil #5 Fuel Oil #6 Gasoline Heating Fuel Jet A Fuel Kerosene

The calibrating standard used for the CaPAHs was benzo(a)pyrene. The calibrating standard concentrations were 0.1, 1.0, and 5.0 ppb. Two sample extracts were analyzed in duplicate for QC.

Laboratory Analysis for Toxic Organic Compounds and Trace Elements

Based on the preliminary findings for the 30 selected sites, 10 sites were selected by the technical advisory group for further laboratory analysis of pesticides, PCBs, PAHs, and selected trace elements. The technical advisory group selected the 10 sites based on the percentage of fine sediments, ⁷Be activity, and concentrations of chemical constituents. The sites selected were site numbers 4, 12, 14, 18, 22, 26, 27, 30, 33, and 60. Samples from the 10 sites were analyzed according to EPA methods listed below:

Chemical Constituents	EPA analytical method
Pesticides	
Organochlorine	EPA Method SW-8080
Organophosphorus	EPA Method SW-8140
Polychlorinated Biphenyls	EPA Method SW-8080
Polycyclic Aromatic Hydrocar-	EPA Method SW-8310
bons	

Selected Trace Elements:

aluminum, arsenic, cadmium,	EPA Method SW-5910
chromium, copper, iron, lead,	EPA Method SW-5910
zinc, and	EPA Method SW-5910
mercury	EPA Method SW-7471A

CHEMICAL AND PHYSICAL CHARACTERISTICS OF SELECTED BOTTOM SEDIMENTS

This section of the report presents the results of physical and chemical analysis of bottom-sediment samples collected during the reconnaissance survey.

Short-Term Sedimentation Patterns Using Beryllium-7 Radiodating

The data for the ⁷Be activity of bottom sediments from the 59 sites in the study area are presented in table 3. The data ranged from not detected to 10.17 dpm/g. Thirty-one of the 59 sites sampled had detectable ⁷Be ranging between 0.95 and 10.17 dpm/g. The locations of the sites in table 3 are identified with reference to their respective surfacewater body. There were 19 recent sedimentation sites in the river and estuary, and 12 recent sedimentation sites in the bays and tributaries. The locations of these sites are presented in plate 2.

Sediment Size

Fifty-nine bottom-sediment samples were analyzed for eight particle sizes. The sizes represent the distribution at the top 1-cm of the bottom sediments. The eight particle sizes included mesh sizes >4, >2, >1, >0.5, >0.25, >0.125, >0.0625, and <0.0625. The total fines are represented by the sum of the percentages from mesh sizes >0.0625 and <0.0625. The corresponding sieve number and phi size are presented with the percentage of each size in table 4. The total percent of fines in the estuary ranged from 10.06 (site 28) to 85.55 (site 31). The total percent of fines in the bays ranged from 39.21 (site 33) to 89.95 (site 51).

Trace Elements

Scans for trace elements are presented in tables 5 and 6. The concentrations of the trace elements, arsenic through zinc, were normalized to aluminum by superimposing the data from this study (tables 5 and 6) on the regression curves published by FDEP (1984).

Table 3. Beryllium-7 radioisotope activity on bottom-sediment samples collected in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998

[cm, centimeter; ⁷Be, Beryllium-7; dpm/g, disintegrations per minute per gram; \pm , statistical counting error; (miles), miles upstream from the confluence with the Caloosahatche River Estuary; Tr, trace, too low to quantify; na, no data; nd, not detected]

Site No.	Depth (cm)	Surface-water body	⁷ Be (dpm/g)	± dpm/g	
1	0-1	Caloosahatchee River ²	3.19	1.24	
2	0-1	Caloosahatchee River ³	10.00	2.17	
3	0-1	Caloosahatchee River Estuary	3.99	1.15	
4	0-1	Caloosahatchee River Estuary	5.75	1.60	
5	0-1	Caloosahatchee River Estuary	1.02	0.02	
6	0-1	Caloosahatchee River Estuary	Tr	na	
7	0-1	Orange River (3.5 miles)	nd	na	
8	0-1	Caloosahatchee River Estuary	nd	na	
9	0-1	Daughtrey Creek (0.6 mile)	3.43	1.12	
10	0-1	Caloosahatchee River Estuary	0.95	0.48	
11	0-1	Caloosahatchee River Estuary	Tr	na	
12	0-1	Caloosahatchee River Estuary	2.26	0.75	
13	0-1	Caloosahatchee River Estuary	nd	na	
14	0-1	Caloosahatchee River Estuary	4.71	1.05	
15	0-1	Caloosahatchee River Estuary	3.88	1.01	
16	0-1	Caloosahatchee River Estuary	2.20	0.95	
17	0-1	Billy Creek (0.9 mile)	nd	na	
18	0-1	Caloosahatchee River Estuary	5.34	1.00	
19	0-1	Caloosahatchee River Estuary	5.25	0.98	
20	0-1	Caloosahatchee River Estuary	2.37	0.66	
21	0-1	Yellow Fever Creek (0.1 mile)	5.86	1.07	
22	0-1	Caloosahatchee River Estuary	3.18	0.80	
23	0-1	Caloosahatchee River Estuary	1.96	0.68	
24	0-1	Caloosahatchee River Estuary	Tr	na	
25	0-1	Caloosahatchee River Estuary	Tr	na	
26	0-1	Caloosahatchee River Estuary	3.42	1.02	
27	0-1	Caloosahatchee River Estuary	6.48	0.93	
28	0-1	Caloosahatchee River Estuary	nd	na	
29	0-1	Caloosahatchee River Estuary	1.31	0.60	
30	0-1	San Carlos Bay	3.63	1.06	
31	0-1	Caloosahatchee River Estuary	Tr	na	
32	0-1	San Carlos Bay	Tr	na	
33	0-1	San Carlos Bay	1.85	0.86	
34	0-1	San Carlos Bay	2.45	0.70	
35	0-1	San Carlos Bay	1.19	0.70	
36	0-1	Not Sampled	na	na 0.00	
30	0-1	San Carlos Bay	Tr	na	
38	0-1	San Carlos Bay	2.24	0.65	
39	0-1	San Carlos Bay			
39 40	0-1	San Carlos Bay	nd Tr	na	
		•		na	
41 42	0-1	San Carlos Bay	nd	na	
	0-1	San Carlos Bay	nd	na	
43	0-1	San Carlos Bay Matlacha Bass	nd	na 0.06	
44 45	0-1 0-1	Matlacha Pass Matlacha Pass	2.80 nd	0.96	
			na	na	

Table 3. Beryllium-7 radioisotope activity on bottom-sediment samples collected in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998 (Continued) [cm, centimeter; ⁷Be, Beryllium-7; dpm/g, disintegrations per minute per gram; ±, statistical counting error; (miles), miles upstream from the confluence with the Caloosahatche River Estuary; Tr, trace, too low to quantify; na, no data; nd, not detected]

Site No.	Depth (cm)	Surface-water body	⁷ Be (dpm/g)	± dpm/g
47	0-1	Tarpon Bay	1.66	0.55
48	0-1	Tarpon Bay	Tr	na
49	0-1	Pine Island Sound	nd	na
50	0-1	Pine Island Sound	Tr	na
51	0-1	Estero Bay	2.51	1.06
52	0-1	Estero Bay	nd	na
53	0-1	Estero Bay	nd	na
54	0-1	Estero Bay	Tr	na
55	0-1	Estero Bay	nd	na
56	0-1	Estero Bay	Tr	na
57	0-1	Estero Bay	Tr	na
58	0-1	Estero Bay	10.17	1.56
59	0-1	Estero Bay	nd	na
60	0-1	Caloosahatchee River Estuary	7.25	1.41

Table 4. Sediment size analyses on bottom-sediment samples collected in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998 [Sample depth is from 0 to 1 centimeter below the bottom sediments. Mesh size, in millimeters; >, greater than; <, less than]

	Sediment size								
Sieve No. Mesh size Phi size	5 >4 -2	10 >2 -1	18 >1 0	35 >0.5 1	60 >0.25 2	120 >0.125 3	230 >0.0625 4	<230 <0.0625 <4	
Site No.				Percent o	dry weight				Total fines ¹
12	7.52	0.70	4.38	9.81	13.38	8.07	16.95	39.18	56.13
2^{2}	0.11	0.21	2.57	21.76	10.11	11.55	11.51	42.17	53.68
3	0.08	0.08	1.17	5.30	3.36	14.01	23.04	52.96	76.00
4	0.54	0.24	4.31	9.74	8.50	5.20	14.86	56.62	71.47
5	0.33	0.64	1.42	3.35	22.36	35.95	26.23	9.72	35.95
6	0.15	0.26	0.50	1.84	19.13	35.99	33.50	8.63	42.13
7	2.75	3.75	2.09	4.15	37.95	13.80	17.08	18.44	35.52
8	0.14	0.34	0.70	0.86	8.35	64.57	22.31	2.74	25.04
9	0.78	2.55	2.68	3.53	12.64	26.06	30.66	21.11	51.76
10	26.96	0.69	0.82	0.63	8.67	27.32	26.80	8.12	34.91
11	0.56	0.89	0.60	0.67	3.03	51.39	38.89	3.98	42.86
12	0.50	1.17	2.46	3.66	8.48	39.77	28.80	15.16	43.96
13	2.40	1.91	4.16	4.30	17.23	34.68	20.62	14.71	35.32
14	1.12	0.90	1.62	2.15	3.48	13.05	19.44	58.22	77.67
15	0.24	1.62	1.96	2.96	5.17	26.71	34.91	26.43	61.34
16	0.35	3.52	3.59	2.10	5.49	38.18	39.69	7.08	46.77
17	0.63	0.43	0.38	0.97	24.21	52.47	14.12	6.79	20.91
18	20.87	2.32	1.79	1.26	0.87	10.40	22.30	40.19	62.49
19	0.39	0.37	0.67	1.06	2.69	21.28	46.26	27.29	73.55

¹² Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998

Table 4. Sediment size analyses on bottom-sediment samples collected in theCaloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998(Continued)

[Sample depth is from 0 to 1 centimeter b	low the bottom sediments. Mesh size, in millimeters;

Sediment size									
Sieve No. Mesh size Phi size	5 >4 -2	10 >2 -1	18 >1 0	35 >0.5 1	60 >0.25 2	120 >0.125 3	230 >0.0625 4	<230 <0.0625 <4	
Site No.				Percent of	dry weight	:			Total fines
20	0.72	0.96	1.41	1.25	3.11	27.36	34.49	30.70	65.19
21	1.39	1.07	1.58	2.12	4.06	11.08	50.99	27.71	78.70
22	1.32	2.29	2.74	2.58	4.20	23.99	44.34	18.54	62.88
23	0.56	0.22	0.34	0.35	1.00	42.86	47.33	7.33	54.67
24	0.14	0.25	0.39	0.47	4.35	53.13	40.30	0.97	41.27
25	0.08	0.09	0.14	0.17	8.73	63.29	27.03	0.48	27.50
26	0.00	0.08	0.47	0.49	0.99	29.36	45.68	22.92	68.61
27	0.22	0.31	0.67	0.70	1.29	21.98	45.70	29.13	74.83
28	0.01	0.01	0.02	0.34	30.52	59.03	9.42	0.64	10.06
29	0.02	0.22	0.54	0.76	7.82	48.11	34.12	8.41	42.52
30	1.61	5.07	12.05	7.77	4.95	10.46	11.35	46.74	58.09
31	0.00	0.04	0.22	4.03	3.00	7.16	16.98	68.56	85.55
32	0.03	0.02	0.23	0.30	1.86	21.94	64.29	11.34	75.63
33	2.66	7.57	4.14	3.00	8.54	34.88	26.43	12.78	39.21
34	0.06	0.16	0.11	0.07	0.74	38.65	43.52	16.70	60.22
35	0.72	0.10	0.65	0.57	3.54	22.04	37.50	34.87	72.37
36				Not sa	ampled				
37	1.92	4.70	10.12	5.21	3.83	17.17	29.75	27.29	57.05
38	0.13	0.07	0.30	0.15	0.15	13.61	55.52	30.07	85.59
39	0.08	0.20	0.44	0.38	1.48	38.32	50.75	8.35	59.10
40	0.18	0.20	0.27	0.25	4.19	44.54	36.29	14.08	50.37
41	0.73	0.22	0.39	0.54	3.06	32.16	49.84	13.06	62.90
42	0.20	0.10	0.18	0.40	2.04	32.94	51.68	12.46	64.14
43	0.10	0.15	0.43	0.40	2.36	30.45	61.19	4.92	66.11
44	0.03	0.20	0.29	0.27	3.84	45.78	38.58	11.01	49.60
45	0.00	0.02	0.03	0.05	2.12	39.33	56.93	1.52	58.45
46	0.45	0.08	0.28	0.78	1.44	43.73	35.80	17.45	53.25
47	0.85	0.63	0.52	0.34	0.50	19.84	58.10	19.22	77.32
48	0.00	0.10	0.11	0.32	1.38	28.08	62.07	7.94	70.01
49	0.30	0.25	0.32	1.74	4.94	48.31	35.24	8.88	44.13
50	0.21	0.58	0.52	0.57	2.47	28.00	61.01	6.65	67.66
51	0.00	0.03	0.25	0.46	0.71	8.60	24.79	65.16	89.95
52	0.29	0.25	0.28	0.36	8.46	49.44	32.50	8.42	40.92
53	0.00	0.17	0.25	0.38	2.12	36.97	49.75	10.36	60.11
54	0.19	0.26	0.29	0.58	4.21	47.87	31.57	15.02	46.59
55	0.00	0.02	0.05	0.05	1.20	32.36	41.59	24.72	66.31
56	0.00	0.11	0.09	0.05	0.92	14.40	56.35	27.92	84.27
57	0.00	0.02	0.14	0.24	0.52	14.50	28.56	56.02	84.58
58	3.43	2.26	2.44	1.45	3.18	48.08	19.75	19.40	39.15
59	0.22	0.28	0.27	0.45	2.21	53.63	38.80	4.14	42.94
60	0.22	0.28	2.84	3.44	2.21	3.60	4.04	82.35	86.39

¹Total fines, sum of 230 (coarse silt) and <230 (silt and clay).

²Sites 1 and 2 located above Franklin Lock (S-79).

 Table 5.
 Scans for trace elements on selected bottom-sediment samples collected in the Caloosahatchee River and Estuary,

 Lee County, Florida, July 20-30, 1998

[Sites 1 and 2 located upstream from Franklin Lock (S-79); distance downstream from Franklin Lock; Concentration in parts per million; D, duplicate sample; <, less than; NG, no guidelines]

Site	Distance				Concent	tration			
number	downstream (in miles)	Aluminum	Arsenic	Cadmium	Chromium	Copper	Lead	Nickel	Zinc
1	1.1	34,800	<5	2.0	74	36	17	16	77
2	1.3	33,600	<5	1.2	58	34	19	19	80
2D	1.3	36,600	<5	1.0	65	36	21	21	88
3	0.7	33,300	<5	1.2	58	32	20	18	78
4	3.1	31,300	<5	1.3	58	34	19	16	76
5	5.4	7,100	<5	0.4	18	7	5	4	17
12	9.3	12,400	<5	0.4	30	14	10	6	44
12D	9.3	11,000	<5	0.5	27	13	8	5	41
15	10.3	14,300	<5	0.7	34	16	13	8	50
14	10.5	23,700	<5	1.1	49	24	20	11	79
16	11.9	1,700	<10	<5	8	5	5	1	<1
20	12.1	7,100	<5	<4	12	9	9	3	26
18	12.2	24,000	<5	1.0	66	55	39	13	161
19	13.0	17,000	<5	0.7	54	34	38	9	120
21	13.7	26,000	8.00	1.1	73	56	32	32	150
22	14.2	12,600	<5	5.0	33	14	12	6	48
22D	14.2	12,600	<5	<4	33	14	12	6	49
23	14.9	4,700	<5	<4	15	5	5	2	19
26	17.6	8,500	<5	0.4	25	9	10	7	30
27	21.2	16,900	<5	0.6	40	30	18	9	74
29	21.8	2,600	<5	<4	10	2	<5	2	8
60	23.8	27,800	<5	0.9	81	34	25	18	94
			Se	diment quality	/ assessment g	juidelines for	trace elemen	its ¹	
	TEL ²	NG	7.24	0.676	52.3	18.7	30.2	15.6	124

¹Sediment Quality Assessment Guidelines, helps address concerns relative to contamination of coastal ecosystems with substances that tend to be associated with sediments (FDEP, 1994).

160

108

112

42.8

271

²TEL, Threshold Effects Level, defines the upper limit of the range of sediment contaminant concentrations dominated by no effects data entries (minimal effect range).

4.21

PEL³

NG

41.6

³PEL, Probable Effects Level, defines the lower limit of the range of sediment contaminant concentrations that are usually or always associated with adverse effects.

 Table 6.
 Scans for trace elements on selected bottom-sediment samples collected in the contiguous bays and tributaries to the Caloosahatchee River Estuary, Lee County, Florida, July 20-30, 1998

[Concentration in parts per million; D, duplicate sample; <, less than; NG, no guidelines]

Cita Na	Leastion				Concen	tration			
Site No.	Location	Aluminum	Arsenic	Cadmium	Chromium	Copper	Lead	Nickel	Zinc
9	Daughtrey Creek	6,000	<10	0.5	24	21	15	7	2
9D	Daughtrey Creek	4,200	<10	< 0.5	19	21	14	5	2
30	San Carlos Bay	4,300	<10	< 0.5	22	7	8	3	1
33	San Carlos Bay	1,400	<10	< 0.5	8	2	3	1	<1
34	San Carlos Bay	2,500	<10	< 0.5	14	4	5	3	<1
35	San Carlos Bay	10,600	<5	< 0.4	34	8	10	5	21
35D	San Carlos Bay	13,100	<5	0.4	38	11	11	6	26
38	San Carlos Bay	12,900	5	0.4	40	9	10	8	28
44	San Carlos Bay	6,600	<10	< 0.5	30	11	11	8	25
44D	San Carlos Bay	6,600	<10	< 0.5	27	7	6	7	16
46	Matlacha Pass	17,800	<5	< 0.4	40	12	18	13	26
47	Tarpon Bay	5,600	<5	< 0.4	20	4	5	5	8
51	Estero Bay	13,400	<10	0.6	87	29	15	12	47
58	Billy Creek	5,400	<10	1.1	62	64	100	14	318
			S	ediment Qualit	y Assessment (Guidelines for	trace elemen	ts ¹	
	TEL ²	NG	7.24	0.676	52.3	18.7	30.2	15.6	124
	PEL ³	NG	41.6	4.21	160	108	112	42.8	271

¹Sediment Quality Assessment Guidelines, helps address concerns relative to contamination of coastal ecosystems with substances that tend to be associated with sediments (FDEP, 1994).

²TEL, Threshold Effects Level, defines the upper limit of the range of sediment contaminant concentrations dominated by no effects data entries (minimal effect range).

³PEL, Probable Effects Level, defines the lower limit of the range of sediment contaminant concentrations that are usually or always associated with adverse effects.

Figures 3 to 9 provide a visual representation of where the bottom sediments in this study plot relative to FDEP's upper and lower 95 percent prediction limits for Natural Range. By superimposing the data from this study on FDEP's regression curves, several observations were noted:

- *Arsenic*: All samples plotted within the Natural Range,
- *Cadmium*: 17 samples plotted above the upper limits,

- *Chromium*: 6 samples plotted above the upper limits,
- Copper: 8 samples plotted above the upper limits,
- *Nickel*: All samples plotted within the Natural Range,
- *Lead*: 8 samples plotted above the upper limits, and
- Zinc: 18 samples plotted above the upper limits.

Plate 3 shows a map locating the sites from which samples were scanned and the number of sites with trace elements above the Natural Range.




Figure 3. Arsenic concentrations normalized to aluminum concentrations for 30 bottom-sediment samples compared to Florida Department of Environmental Protection regression line and prediction limits, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.

16 Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998




Figure 4. Cadmium concentrations normalized to aluminum concentrations for 30 bottom-sediment samples compared to Florida Department of Environmental Protection regression line and prediction limits, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.






Figure 5. Chromium concentrations normalized to aluminum concentrations for 30 bottom-sediment samples compared to Florida Department of Environmental Protection regression line and prediction limits, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.

18 Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998




Figure 6. Copper concentrations normalized to aluminum concentrations for 30 bottom-sediment samples compared to Florida Department of Environmental Protection regression line and prediction limits, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.




Figure 7. Nickel concentrations normalized to aluminum concentrations for 30 bottom-sediment samples compared to Florida Department of Environmental Protection regression line and prediction limits, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.




Figure 8. Lead concentrations normalized to aluminum concentrations for 30 bottom-sediment samples compared to Florida Department of Environmental Protection regression line and prediction limits, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.




Figure 9. Zinc concentrations normalized to aluminum concentrations for 30 bottom-sediment samples compared to Florida Department of Environmental Protection regression line and prediction limits, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.

Total Organic Carbon

TOC analyses of the 30 bottom-sediment samples collected in the study area are presented in table 7 and plate 4. The data for the Caloosahatchee River and Estuary are referenced to the respective site numbers and distance in miles above and below the Franklin Lock (S-79); data for the bays and tributaries are referenced only to the site numbers. TOC concentrations in the bottom sediments of the river and estuary ranged from 4,600 to 164,000 ppm; TOC concentrations in the bays and tributaries ranged from 4,290 to 142,000 ppm.

Immunoassay Results for Toxic Organic Compounds

ELISA analyses for the 30 bottom-sediment samples collected in the study area are presented in table 8 and plates 5, 6, and 7. The SQAGs that were established by FDEP (1994) are presented in table 8. At 11 sites, the concentrations of the DDT ELISA tests

were above the FDEP's PEL of 0.0517 mg/kg. All the sites had concentrations of cyclodienes as chlordane greater than the PEL. There were no samples that responded to the PCB ELISA test. All of the samples responded to the CaPAH ELISA test. The concentrations for CaPAH were below the TEL except for sites 19 and 58 which were above the TEL; there were no sites at or above the PEL. PEL defines the lower limit of the range of contaminant concentrations that are usually or always associated with adverse biological effects, whereas TEL defines the upper limit of the range of sediment contaminant concentrations dominated by the minimal effects range (FDEP, 1994). As with all ELISA tests, cross-reactivity needs to be considered (Mike Meyer, USGS, verbal commun., 1999). Therefore, the data presented in table 8 do not absolutely indicate that any individual compound exceeded the PEL. However, the data do suggest that quantitative analyses should be performed on samples to determine which compounds responded to the different ELISA tests and if any of these compounds exceeded the PEL.

Table 7. Laboratory analyses for total organic carbon on selected bottom-sediment samplescollected in the Caloosahatchee River and Estuary study area, Lee County, Florida,July 20-30, 1998

Sites in the Caloosahatchee River and Estuary								
Site No.	Distance downstream (in miles)	TOC (ppm)	Site No.	Distance downstream (in miles)	TOC (ppm)			
1	-1.1	74,800	18	12.2	36,600			
2	-1.3	164,000	19	13.0	16,400			
3	0.7	149,000	21	13.7	90,600			
4	3.1	124,000	22	14.2	26,000			
5	5.4	30,600	23	14.9	4,600			
12	9.3	29,100	26	17.6	20,600			
14	10.5	69,900	27	21.2	26,700			
15	10.3	34,500	29	24.3	5,570			
16	11.9	17,700	60	23.8	54,300			
20	12.1	71,400						

[TOC, total organic carbon; ppm, parts per million]

Sites in bays and tributaries to the Caloosahatchee River estuary

Site No.	Location	TOC (ppm)	Site No.	Location	TOC (ppm)
9	Daughtrey Creek	142,000	44	San Carlos Bay	4,290
30	San Carlos Bay	33,400	46	Matlacha Pass	27,500
33	San Carlos Bay	6,800	47	Matlacha Pass	10,000
34	San Carlos Bay	11,800	51	Estero Bay	27,500
35	San Carlos Bay	30,700	58	Billy Creek	34,300
38	San Carlos Bay	26,700			

Table 8. Immunoassay screening on selected bottom-sediment samples collected in the Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998

[Site number, site numbers 1 and 2 are sites located above the Franklin Lock (S-79); Distance downstream from Franklin Lock (S-79); PCBs, polychlorinated biphenyls; mg/kg, milligrams per kilogram; µg/kg, micrograms per kilogram; <, less than; >, greater than; concentrations based on dry weight; DDT, includes DDT metabolites; PCBs, polychlorobiphenyls; CaPAH, carcinogenic polycyclic hydrocarbons]

		DDT ¹	hee River and Estuary Chlordane ²	PCBs ³	CaPAH ⁴
Site No.	Distance down- stream (in miles)	mg/kg	Chlordane ² mg/kg	PCBs ³ mg/kg	CaPAH≁ μg/kg
1	-1.1	0.89	1.49		239
2	-1.3	1.00	0.89	<0.25	15
3	0.7	<0.05	1.52	(0.25	317
4	3.1	0.61	1.72	<0.25	333
5	5.4	<0.05	0.62	(0120	32
12	9.3	0.29	0.67		397
15	10.3	< 0.05	1.02		314
14	10.5	< 0.05	0.76	<0.25	183
16	11.9	< 0.05	0.47	<0.25	177
20	12.1	0.23	0.47		335
18	12.2	< 0.05	1.17		1378
19	13.0	< 0.05	0.88	< 0.25	1894
21	13.7	>0.05 but <0.1	1.62	< 0.25	620
22	14.2	< 0.05	1.01		401
23	14.9	< 0.05	0.29	< 0.25	257
26	17.6	< 0.05	0.52		419
27	21.2	< 0.05	0.68	< 0.25	274
29	21.8	< 0.05	0.37		81
60	23.8	>0.05 but <0.1	1.02	< 0.25	131
9	Daughtrey Creek	>0.05 but <0.1	1.34	< 0.25	310
30	San Carlos Bay	1.27	1.81	< 0.25	221
33	San Carlos Bay	0.21	0.41		33
34	San Carlos Bay	>0.05 but <0.1	0.89	<0.25	92
35	San Carlos Bay	< 0.05	0.46		96
38	San Carlos Bay	>0.05 but <0.1	0.87	< 0.25	100
44	San Carlos Bay	>0.05 but <0.1	0.33		23
46	Matlacha Pass	0.60	1.31	< 0.25	195
47	Tarpon Bay	< 0.05	0.77		23
51	Estero Bay	>0.05 but <0.1	0.82	< 0.25	280
58	Billy Creek	1.64	1.21		1,914
		Sediment Qual	ity Assessment Guid	elines ⁵	
	TEL ⁶	0.00389	0.00226	0.0216	1,684
	PEL ⁷	0.0517	0.00479	0.189	16,770

¹DDT, DDT test cross-reacts with DDTs, DDEs, DDDs, DDA, chloropropylate, chlorobenzilate, dicofol, and tetradifon; concentrations may represent one, or the sum of part or all the compounds.

²Chlordane, chlordane test cross-reacts with aldrin, endosulfan, endrin, dieldrin , heptachlor, and toxaphene. Concentrations may represent one, or the sum of part or all of the compounds.

³PCBs, PCBs test cross-reacts with Aroclors 1016, 1221, 1232, 1242, 1248, 1254, 1260, 1268, and 1282; concentrations may represent one, or the sum of more than one compound.

⁴CaPAHs; CaPAHs test cross-reacts with the compounds below; concentrations may represent one, or the sum of part or all the compounds.

Benzo[a]pyrene	Phenanthrene	Fuel Oil #4
Benzo[a]anthracene	Benzo[g,h,i]perylene	Fuel Oil #5
Benzo[k]fluoranthene	Pyrene	Fuel Oil #6
Chrysene	Fluorene	Heating Fuel
Benzo[b]fluoranthene	Naphthalene	Diesel Fuel
Indeno[1,2,3-c,d]pyrene	Acenaphthalene	Gasoline
Dibenzo[a,h]anthracene	Creosote	Kerosene
Anthracene	Fluoranthene	Jet A Fuel
Acenaphthylene		

⁵Sediment Quality Assessment Guidelines based on dry basis. FDEP suggests that SQAGs may be used to assess the adverse biological effects that could, potentially, be associated with levels of sediment-associated contaminants (FDEP, 1994).

⁶TEL, Threshold Effects Level, defines the upper limit of the range of sediment contaminant concentrations dominated by no effects data entries (FDEP, 1994).

⁷PEL, Probable Effects Level, defines the lower limit of the range of contaminant concentrations that are usually or always associated with adverse biological effects.

Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River 24 and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998

The concentrations of all samples that responded to the ELISA tests were normalized to the TOC concentration of each respective sample (FDEP, 1994). The normalized ELISA DDT test data for the sites in the river and estuary are presented in figure 10; the data for the bays and tributaries are presented in table 9. The normalized data for the ELISA Chlordane and CaPAH tests for the sites in the river and estuary are presented in figures 11 and 12 respectively; the data for the bays and tributaries are presented in table 9. The FDEP (1994) presents a discussion on the use of normalizing the sum of each group of organic compounds in a sample to the sample's TOC. The normalized toxic organic compound data can be used to compare the sampling sites on the relative potential impact on the benthic organisms at the respective sites (FDEP, 1994). The potential impact is relative to the size of the bar (figs. 10-12).




Figure 10. DDT enzyme-linked immunosorbent assay data normalized to total organic carbon, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.

Table 9. DDT, cyclodienes as chlordane, and carcinogenic polycyclic aromatic hydrocarbons normalized to total organic carbon on selected bottom-sediment samples collected in the contiguous bays and tributaries to the Caloosahatchee River Estuary, Lee County, Florida, July 20-30, 1998

[Factor, ratio multiplied by a number so that the ratio is reported as a whole number; NA, not applicable; reported findings are trace or not detected]

Site Location _	Ratio to total organic carbon x's factor					
	-	DDT ¹	Chlordane ²	CaPAH ³		
9	Daughtrey Creek	NA	9	22		
30	San Carlos Bay	38	54	66		
33	San Carlos Bay	31	60	49		
34	San Carlos Bay	NA	75	78		
35	San Carlos Bay	NA	15	31		
38	San Carlos Bay	NA	33	37		
44	San Carlos Bay	NA	77	54		
46	Matlacha Pass	22	48	71		
47	Tarpon Bay	NA	77	23		
51	Estero Bay	NA	30	102		
58	Billy Creek	48	35	558		

¹Factor equals 10⁶. DDT, DDT test cross-reacts with DDTs, DDEs, DDDs, DDA, chloropropylate, chlorobenzilate, dicofol, and tetradifon; concentrations may represent one, or the sum of part or all the compounds.

²Factor equals 10⁶. Chlordane, chlordane test cross-reacts with the cyclodiene compounds aldrin, endosulfan, endrin, dieldrin, heptachlor, and toxaphene. Concentrations may represent one, or the sum of part or all the compounds.

³Factor equals 10⁵. CaPAHs; CaPAHs test cross-reacts with the compounds below; concentrations may represent one, or the sum of part or all the compounds.

Benzo[a]pyrene	Acenaphthylene
Benzo[a]anthracene	Fluorene
Benzo[k]fluoranthene	Naphthalene
Chrysene	Creosote
Benzo[b]fluoranthene	Fuel Oil #4
Indeno[1,2,3-c,d]pyrene	Fuel Oil #5
Dibenzo[a,h]anthracene	Fuel Oil #6
Anthracene	Heating Fuel
Phenanthrene	Diesel Fuel
Benzo[g,h,i]perylene	Gasoline
Acenaphthalene	Kerosene
Pyrene	Jet A Fuel
Fluoranthene	





Figure 11. Chlordane enzyme-linked immunosorbent assay data normalized to total organic carbon, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.



RIVER MILES DOWNSTREAM FROM FRANKLIN LOCK (S-79)

Figure 12. Carcinogenic polycyclic aromatic hydrocarbons enzyme-linked immunosorbent assay data normalized to total organic carbon, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998.

26 Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998

Laboratory Results for 10 Selected Sites

The data for the bottom-sediment samples analyzed for toxic organic compounds and selected trace elements are presented in tables 10 and 11, and in the appendix. All the samples analyzed are from the Caloosahatchee River estuary. The reporting limits (detection limits) for the toxic organic compounds and trace elements varied with each sample because of the different moisture content of the samples. The reporting limits are directly related to the moisture content of the samples (Christina Mott, Quanterra Laboratories, verbal commun., April 1999).

Table 10. Laboratory analytical results for pesticides, polychlorinated biphenyls, and polycyclicaromatic hydrocarbons on 10 bottom-sediment samples collected in the Caloosahatchee RiverEstuary, Lee County, Florida, July 20-30, 1998

[μg/kg, micrograms per kilogram; PCBs, polychlorinated biphenyls; PAH, polycyclic aromatic hydrocarbon; nd, not detected; J, estimated result, result is less than reporting limit]

0:1-		Death		ration				
Site No.		Pestic			PCBs ¹	μ g/kg	PAH ³	μ g/kg
-	organo- chlorine ¹	μ g/kg	organo- phosphorus ²	μ g/kg	1003	μg/kg		μց/Νց
4	nd		nd		nd		Anthracene	130J
							Benzo(a)anthracene	150J
							Benzo(b)fluoranthene	840J
							Benzo(ghi)perylene	110J
							Benzo(k)fluoranthene	270J
							Fluoranthene	1,100J
12	4,4'-DDD	5.3J	nd		nd		Benzo(a)pyrene	120J
							Benzo(b)fluoranthene	240J
							Fluoranthene	230J
14	4,4'-DDD	3.3J	nd		nd		Benzo(a)anthracene	60J
							Benzo(a)pyrene	120J
							Benzo(b)fluoranthene	270
							Fluoranthene	210J
18	nd		nd		nd		Anthracene	95J
							Benzo(a)anthracene	940
							Fluoranthene	4,100
							Phenanthrene	730
							Pyrene	1,700
22	nd		nd		nd		Benzo(a)pyrene	58J
							Benzo(b)fluoranthene	91J
							Fluoranthene	78J
26	delta-BHC	3.8J	nd		nd		Benzo(a)pyrene	78
							Benzo(ghi)perylene	71J
							Benzo(k)fluoranthene	40J
							Fluoranthene	110J
27	gamma-BHC	1.5J	nd		nd		Benzo(k)fluoranthene	21J
	(Lindane)						Fluoranthene	58J
							Fluorene	130
							Pyrene	57J
30	nd		nd		nd		nd	
33	nd		nd		nd		nd	
60	nd		nd		nd		Benzo(a)pyrene Benzo(b)fluoranthene	120J 190J

¹Environmental Protection Agency Method SW8080A.

²Environmental Protection Agency Method SW8140.

³Environmental Protection Agency Method SW8310.

 Table 11.
 Laboratory analytical results for trace elements on 10 bottom-sediment samples collected in the Caloosahatchee

 River Estuary, Lee County, Florida, July 20-30, 1998

[Concentration, in milligrams per kilogram; nd, not detected; na, not applicable; B, estimated result. Result is less than reporting limits]

0:1- 1-	Concentration								
Site No.	Aluminum	Arsenic	Cadmium	Chromium	Copper	Iron	Lead	Mercury	Zinc
4	33,900	nd	nd	110.0	123.0	78,800	61.4B	nd	290
12	10,000	nd	nd	42.0	34.5	22,400	24.0B	nd	110
14	9,960	nd	nd	42.1	35.1	24,400	24.3B	0.21	119
18	15,200	nd	nd	78.2	107.0	35,700	72.2	nd	296
22	3,340	nd	nd	15.3	11.4	7,950	10.1B	nd	46.3
26	3,060	nd	nd	14.7	10.0	7,120	8.9B	nd	36.4
27	1,700	nd	nd	8.6	9.3	3,540	nd	nd	25.0
30	6,000	nd	nd	30.6	12.3	8,320	10.7B	nd	29.3
33	1,630	nd	nd	8.5	2.7B	2,220	nd	nd	7.4
60	26,300	nd	nd	127.0	80.4	41,400	55.5	nd	223
				Sediment Qu	ality Assessme	ent Guidelines ¹	I		
TEL ²	na	7.24	0.676	52.3	18.7	na	30.2	0.13	124
PEL ³	na	41.6	4.21	160	108	na	112	0.696	271

¹Sediment Quality Assessment Guidelines, helps address concerns relative to contamination of coastal ecosystems with substances that tend to be associated with sediments (FDEP, 1994).

²TEL, Threshold Effects Level, defines the upper limit of the range of sediment contaminant concentrations dominated by no effects data entries (minimal effect range).

³PEL, Probable Effects Level, defines the lower limit of the range of sediment contaminant concentrations that are usually or always associated with adverse effects.

Toxic Organic Compounds

The toxic organic compounds in the estuary are reported as pesticides (organochlorine and organophosphorus), PCBs, and PAHs. There were 3 organochlorine pesticide compounds, 4,4'-DDD, delta-BHC, and gamma-BHC (Lindane), detected in sediment samples from four sites (sites 12,14, 26, and 27) (table 10, appendix). The concentrations of the DDD's and BHC's were all below the reporting limit. There were no organophosphorus pesticides or PCBs reported for the 10 sediment samples.

There were eight sites with detectable PAHs; however, only four sites (14, 18, 26, and 27) had reportable concentrations (table 10). The remaining sites (4, 12, 22, and 60) had concentrations less than the reportable limits for the method of analysis. There were no PAHs detected in sites 30 and 33.

Trace Elements

All 10 sites in the estuary had sediment samples with trace elements (table 11). No arsenic or cadmium was detected in the samples; however, aluminum, chromium, copper, iron, lead, mercury, and zinc were detected. The ranges of concentrations of the trace elements detected are listed below:

- Aluminum: from 1,630 to 33,900 mg/kg,
- Chromium: from 8.5 to 127 mg/kg,
- Copper: from 2.7 to 123 mg/kg,
- Iron: from 2,220 to 78,800 mg/kg,
- Lead: from not detected to 72.2 mg/kg,
- *Mercury*: from not detected to 0.21 mg/kg (only at one site), and
- *Zinc*: from 7.4 to 296 mg/kg

There were five sites with trace elements greater than the TEL and PEL. The five sites and the concentrations in mg/kg at each site are shown below.

Site No.	Chromium	Copper	Lead	Mercury	Zinc
4	110	123	61.4		290
12		34.5			
14		35.1		0.21	
18	78.2	107	72.2		296
60	127	80.4	55.5		223
TEL	52.3	18.7	30.2	0.13	124
PEL	160	108	112	0.696	271

SUMMARY

The South Florida Water Management District (District) is developing a Caloosahatchee River Water Management Plan to address environmental and water-supply needs of the Caloosahatchee watershed. The study area of this report includes the Caloosahatchee River and Estuary, and the contiguous water bodies of Matlacha Pass, San Carlos Bay, Estero Bay, Tarpon Bay, and Pine Island Sound, in Lee County. As part of this plan, the District will evaluate potential toxic substances in the sediments of the study area. The toxic substances include anthropogenic organic compounds (polynuclear aromatic hydrocarbons (PAH) organochlorine pesticides, organophosphorus pesticides), and trace elements. The data in this report provide chemical and physical characterization of sediments at selected sites in the study area. The USGS reconnaissance survey consisted of the following activities:

- Determination of the chemistry and physical characteristics of 60 selected bottom sediment-sampling sites including 2 sites above the Franklin Lock (S-79), and
- Selection of 10 sampling sites for analysis of toxic organic compounds and selected trace elements. The 10 sites were selected from the 60 bottom sediment-sampling sites.

A technical advisory group, consisting of USGS and South Florida Water Management District personnel, selected the initial 60 sampling sites. Fifty-eight sites were located in the estuary, tributaries, and the bays; and two sites were located upstream of the Franklin Lock.

A sampling strategy was developed to provide chemical and physical characterization of bottom sediments at selected sites in the study area. The strategy was developed with the cooperation of the technical advisory group. The strategy specified collection of approximately the top 1-cm of bottom sediment samples at 60 selected sites and the physical and chemical characterization of the sediment samples. The characterization included short-term sediment accumulation patterns, sediment-size distribution, concentrations of trace elements, total organic carbon, and selected toxic organic compounds. Fifty-nine of the planned 60 sites were sampled; one site, located in San Carlos Bay, was not sampled because the bottom sediments consisted of sand within a 1-mile radius. Bottom samples from 59 sites were analyzed for ⁷Be activity; thirty-one of the 59 sites sampled had detectable ⁷Be; nineteen in the river/estuary, and twelve in the bays and tributaries. Sediment-size distribution analysis was used to determine the extent of fine bottom sediments. The total percent of fines in the estuary ranged from 10.06 (site 28) to 85.55 (site 31). The total percent of fines in the bays ranged from 39.21 (site 33) to 89.95 (site 51).

Scans for trace elements were used to identify the presence of trace elements above Natural Range. The trace elements were differentiated from varied natural background concentrations by comparing data with FDEP regression curves based on the normalization of trace elements concentrations to aluminum concentration in the sample. Cadmium, chromium, copper, lead, and zinc were above Natural Range at some sites in the estuary and bays.

Total organic carbon analyses were used to identify the locations of carbonaceous sediments. TOC concentration in 30 selected sites ranged between 4,290 and 164,000 parts per million. The TOC from sediments in 18 sites in the river and estuary ranged between 4,600 (at site 23) and 164,000 parts per million (at site 2). The TOC in the bays and tributaries ranged between 4,290 at site 44 (Matlacha Pass) and 142,000 parts per million at site 9 (Daughtrey Creek).

Immunoassay analysis of competitive enzymelinked immunosorbent assay (ELISA) was used to identify and estimate semiquantitatively the concentration of groups of toxic compounds. Three groups were identified; chlorinated pesticides (DDTs and cyclodienes as chlordane), polychlorinated biphenyls (Aroclors) and carcinogenic polycyclic aromatic hydrocarbons. All the concentrations from samples that responded to the DDT and Chlordane ELISA test were above the Florida Department of Environmental Protection's Probable Effect Level (PEL).

Laboratory analysis for toxic organic compounds and selected trace elements was performed on samples from 10 sites (4, 12, 14, 18, 22, 26, 27, 30, 33, and 60). The toxic organic compounds included the organochlorine and organophosphorus pesticides, PCBs, and PAHs. The selected trace elements included arsenic, cadmium, chromium, copper, iron, lead, mercury, and zinc. There were 3 organochlorine pesticide compounds, 4,4'-DDD, delta-BHC, and gamma-BHC (Lindane), detected in sediment samples from four sites (12,14, 26, and 27). The concentrations of the DDD and BHC's were all below the reporting limit. There were no organophosphorus pesticides or PCBs reported for the 10 sediment samples. There were eight sites with detectable PAHs; however, only four sites (14, 18, 26, and 27) had concentrations equal to or greater than the reportable limit. Sites 4, 12, 22, and 60 had concentrations less than the reportable limit for the method of analysis. There were no PAHs detected in sites 30 and 33.

Trace elements at 10 sites were analyzed in the USGS contract laboratory. No arsenic and cadmium were detected in the samples; however, there were five sites (4, 12, 14, 18, and 60) with trace elements greater than the TEL or the PEL.

REFERENCES

- Florida Department of Environmental Protection, 1988, A guide to the interpretation of metal concentrations in estuarine sediments: Tallahassee, Florida, 44 p.
- 1994, Florida coastal sediment contaminants atlas,
 A summary of coastal sediment quality surveys: Tallahassee, Florida, 111 p.
- Geyh, M.A., and Schleicher, H., 1990, Absolute age determination-Physical and chemical dating methods and their applications: Berlin, Springer-Verlag, 503 p.
- Goodwin, C.R., 1991, Simulation of the effects of proposed tide gates on circulation, flushing, and water quality in residential canals, Cape Coral: U.S. Geological Survey Open-File Report 91-237, 43 p.
- Holmes, C.W., 1998, Short-lived isotopic chronometers, A means of measuring decadal sedimentary dynamics: U.S. Geological Survey Fact Sheet FS-073-98.
- La Rose, H.R., and McPherson, B.F., 1983, Chemical and hydrologic assessment of the Caloosahatchee River basin, Lake Okeechobee to Franklin Lock: U.S. Geological Survey Water-Resources Investigations Report 83-4126, 63 p.

- Matthes, W.J., Jr., 1992, Quality-assurance plan for the analysis of fluvial sediment by laboratories of the U.S. Geological Survey: Open-File Report 91-467, 31 p.
- Pierce, A. C., ed., 1995, Florida Statistical Abstract (29th ed.): Gainesville, Fla., University Press of Florida, Gainesville, Florida, Bureau of Economic and Business Research, College of Business Administration.
- Schoellhamer, D.H., 1991, Size classification of bed sediment and selection of resuspension monitoring sites in Upper Tampa Bay, Florida: U.S. Geological Survey Water-Resources Investigations Report 91-4070, 23 p.
- Stoker, Y.E., 1986, Water quality of the Charlotte Harbor estuarine system, Florida, November 1982 through October 1984: U.S. Geological Survey Open-File Report 85-563, 213 p.
- Strategic Diagnostics Incorporated, 1998, Remediation, assessment, and industrial testing product information: Newark, New Jersey.

Verbal and written communications:

- Cantillo, Adriana, National Oceanic and Atmospheric Administration, 1997, written communication.
- Crean, D., and Chamberlain, R., South Florida Water Management District, 1985, written communication.
- Haunert, D., South Florida Water Management District, 1978, written communication.
- Holmes, Charles, U.S. Geological Survey, 1998, verbal communication.
- Meyer, Mike, U.S. Geological Survey, 1999, verbal communication.
- Mott, Christina, Quanterra Laboratory, Arvada, Colorado, 1999, verbal communication.
- Pellicer, Tony, Natural Resource Manager, Lee County, Florida, 1999, verbal communication.
- Seal, Thomas, Florida Department of Environmental Protection, 1999, written communication.

Appendix

Analytical report for organic compounds and trace elements in 10 selected bottom-sediment samples, Caloosahatchee River and Estuary study area, Lee County, Florida, July 20-30, 1998

32 Reconnaissance of Chemical and Physical Characteristics of Selected Bottom Sediments of the Caloosahatchee River and Estuary, Tributaries, and Contiguous Bays, Lee County, Florida, July 20-30, 1998



Quanterra Incorporated 4955 Yarrow Street Arvada, Colorado 80002

303 421-6611 Telephone 303 431-7171 Fax

ANALYTICAL REPORT

FL00108

Lot #: D9B240185

Richard Daddow

U.S. Geological Survey

QUANTERRA INCORPORATED

WW. hott

Christina M. Mott Project Manager

May 24, 1999

Case Narrative Lot D9B240185

This report has been revised to show the results as dry weight corrected.

Organophophorus Pesticides by GC, Method 8140

The spike compound, Ethyl Parathion, is reported above the acceptance limit for the Laboratory Control Samples. As the method blank and matrix spike samples are in control, and there are no samples that contain Ethyl Parathion above the reporting limit, the data are reported.

Polynuclear Aromatic Hydrocarbons by HPLC, Method 8310

The Relative Percent Difference value for Fluorene in the Laboratory Control Samples is reported above the acceptance limit. As the individual percent recoveries are in control, the data are reported.

Several percent recoveries and one RPD value are reported outside of the acceptance limits for several of the analytes for the matrix spike samples due to sample matrix.

With exceptions noted as flags or footnotes, standard analytical protocols were followed in the analysis of the samples and no problems were encountered or anomalies observed. All laboratory quality control samples analyzed in conjunction with the samples in this project were within established control limits.

EXECUTIVE SUMMARY - Detection Highlights

D9B240185

PARAMETER	RESULT	REPORTING	UNITS	ANALYTICAL METHOD
FL00108 SITE#4 02/22/99 001				
Anthracene	130 J	890	ug/kg	SW846 8310
Benzo(a)anthracene	150 J	890	ug/kg	SW846 8310
Benzo(b)fluoranthene	840 J	890	ug/kg	SW846 8310
Benzo(ghi)perylene	110 J	1800	ug/kg	SW846 8310
Benzo(k)fluoranthene	270 J	890	ug/kg	SW846 8310
Fluoranthene	1100 J	1800	ug/kg	SW846 8310
Aluminum	33900	223	mg/kg	SW846 6010B
Chromium	110	22.3	mg/kg	SW846 6010B
Copper	123	44.6	mg/kg	SW846 6010B
Iron	78800	223	mg/kg	SW846 6010B
Lead	61.4 B	112	mg/kg	SW846 6010B
Zinc	290	44.6	mg/kg	SW846 6010B
Percent Moisture	95.5	0.10	8	MCAWW 160.3 MOD
FL00108 SITE#12 02/22/99 002				
4,4'-DDD	5.3 J	24	ug/kg	SW846 8080A
Benzo(a)pyrene	120 J	290	ug/kg	SW846 8310
Benzo(b)fluoranthene	240 J	290	ug/kg	SW846 8310
Fluoranthene	230 J	570	ug/kg	SW846 8310
Aluminum	10000	71.8	mg/kg	SW846 6010B
Chromium	42.0	7.2	mg/kg	SW846 6010B
Copper	34.5	14.4	mg/kg	SW846 6010B
Iron	22400	71.8	mg/kg	SW846 6010B
Lead	24.0 B	35.9	mg/kg	SW846 6010B
Zinc	110	14.4	mg/kg	SW846 6010B
Percent Moisture	86.1	0.10	00	MCAWW 160.3 MOD
FL00108 SITE#14 02/22/99 003				
4,4'-DDD	3.3 J	19	ug/kg	SW846 8080A
Benzo(a)anthracene	60 J	230	ug/kg	SW846 8310
Benzo(a)pyrene	120 J	230	ug/kg	SW846 8310
Benzo(b)fluoranthene	270	230	ug/kg	SW846 8310
Fluoranthene	210 J	450	ug/kg	SW846 8310
Mercury	0.21	0.19	mg/kg	SW846 7471A
Aluminum	9960	56.6	mg/kg	SW846 6010B
Chromium	42.1	5.7	mg/kg	SW846 6010B
Copper	35.1	11.3	mg/kg	SW846 6010B
Iron	24400	56.6	mg/kg	SW846 6010B
Lead	24.3 B	28.3	mg/kg	SW846 6010B
Zinc	119	11.3	mg/kg	SW846 6010B
Percent Moisture	82.3	0.10	8	MCAWW 160.3 MOD

(Continued on next page)

EXECUTIVE SUMMARY - Detection Highlights

D9B240185

			REPORTING		ANALY	TICAL	
PARAMETH	SR	RESULT	LIMIT	UNITS	METHO	D	
FL00108 SITE#18	3 02/22/99 004						
Anthrace	ene	95 J	180	ug/kg	SW846	8310	
Benzo(a)	anthracene	940	180	ug/kg	SW846	8310	
Fluorant	hene	4100	360	ug/kg	SW846	8310	
Phenanth	irene	730	360	ug/kg	SW846	8310	
Pyrene		1700	360	ug/kg	SW846	8310	
Aluminum	n	15200	89.3	mg/kg	SW846	6010B	
Chromium	n	78.2	8.9	mg/kg	SW846	6010B	
Copper		107	17.9	mg/kg	SW846	6010B	
Iron		35700	89.3	mg/kg		6010B	
Lead		72.2	44.6	mg/kg		6010B	
Zinc		296	17.9	mg/kg		6010B	
Percent	Moisture	88.8	0.10	%		160.3	MOD
FL00108 SITE#22	2 02/22/99 005						
Benzo(a)	pyrene	58 J	140	ug/kg	SW846	8310	
Benzo(b)	fluoranthene	91 J	140	ug/kg	SW846	8310	
Fluorant	hene	78 J	280	ug/kg	SW846	8310	
Aluminum	1	3340	34.7	mg/kg	SW846	6010B	
Chromium	1	15.3	3.5	mg/kg	SW846	6010B	
Copper		11.4	6.9	mg/kg	SW846	6010B	
Iron		7950	34.7	mg/kg	SW846	6010B	
Lead		10.1 B	17.3	mg/kg	SW846	6010B	
Zinc		46.3	6.9	mg/kg	SW846	6010B	
Percent	Moisture	71.2	0.10	*	MCAWW	160.3	MOD
FL00108 SITE#26	02/22/99 006						
delta-BH	IC	3.8 J	5.5	ug/kg	SW846	8080A	
Benzo(a)	pyrene	78	65	ug/kg	SW846		
	i)perylene	71 J	130	ug/kg	SW846		
Benzo(k)	fluoranthene	40 J	65	ug/kg	SW846		
Fluorant	hene	110 J	130	ug/kg	SW846		
Aluminum	L	3060	32.5	mg/kg	SW846		
Chromium	L	14.7	3.3	mg/kg	SW846		
Copper		10	6.5	mg/kg	SW846		
Iron		7120	32.5	mg/kg	SW846		
Lead		8.9 B	16.3	mg/kg	SW846		
Zinc		36.4	6.5	mg/kg	SW846		
Percent	Moisture	69.3	0.10	8		160.3	MOD

(Continued on next page)

EXECUTIVE SUMMARY - Detection Highlights

D9B240185

PARAMETER	RESULT	REPORTING	UNITS	ANALYTICAL METHOD
FL00108 SITE#27 02/22/99 007				
gamma-BHC (Lindane)	1.5 J	4.9	uq/kq	SW846 8080A
Benzo(k)fluoranthene	21 J	57	ug/kg	SW846 8310
Fluoranthene	58 J	110	ug/kg	SW846 8310
Fluorene	130	110	ug/kg	SW846 8310
Pyrene	57 J	110	ug/kg	SW846 8310
Aluminum	1700	28.6	mg/kg	SW846 6010B
Chromium	8.6	2.9	mg/kg	SW846 6010B
Copper	9.3	5.7	mg/kg	SW846 6010B
Iron	3540	28.6	mg/kg	SW846 6010B
Zinc	25.0	5.7	mg/kg	SW846 6010B
Percent Moisture	65.1	0.10	8	MCAWW 160.3 MOD
FL00108 SITE#30 02/22/99 008				
Aluminum	6000	42.4	mg/kg	SW846 6010B
Chromium	30.6	4.2	mg/kg	SW846 6010B
Copper	12.3	8.5	mg/kg	SW846 6010B
Iron	8320	42.4	mg/kg	SW846 6010B
Lead	10.7 B	21.2	mg/kg	SW846 6010B
Zinc	29.3	8.5	mg/kg	SW846 6010B
Percent Moisture	76.4	0.10	3, J %	MCAWW 160.3 MOD
FL00108 SITE#60 02/22/99 009				
Benzo(a)pyrene	120 J	420	ug/kg	SW846 8310
Benzo(b)fluoranthene	190 J	420	ug/kg	SW846 8310
Aluminum	26300	105	mg/kg	SW846 6010B
Chromium	127	10.5	mg/kg	SW846 6010B
Copper	80.4	21.1	mg/kg	SW846 6010B
Iron	41400	105	mg/kg	SW846 6010B
Lead	55.5	52.7	mg/kg	SW846 6010B
Zinc	223	21.1	mg/kg	SW846 6010B
Percent Moisture	90.5	0.10	~	MCAWW 160.3 MOD
FL00108 SITE#33 02/22/99 010				
Aluminum	1630	26.1	mg/kg	SW846 6010B
Chromium	8.5	2.6	mg/kg	SW846 6010B
Copper	2.7 B	5.2	mg/kg	SW846 6010B
Iron	2220	26.1	mg/kg	SW846 6010B
Zinc	7.4	5.2	mg/kg	SW846 6010B
Percent Moisture	61.7	0.10	%	MCAWW 160.3 MOD

ANALYTICAL METHODS SUMMARY

D9B240185

PARAMETER	ANALY METHO		
Inductively Coupled Plasma (ICP) Metals Mercury in Solid Waste (Manual Cold-Vapor) Organochlorine Pesticides and PCBs Organophosphorous Pesticides by GC Percent Moisture	SW846 SW846 SW846 MCAWW	160.3	MOD
Polynuclear Aromatic Hydrocarbons by HPLC	SW846	8310	

References:

MCAWW	"Methods	for	Chem	ical	Analys	is of	1 Water	and	Wastes",	
	EPA-600/4	1-79-	-020,	Marc	h 1983	and	subsequ	lent	revision	s.

SW846 "Test Methods for Evaluating Solid Waste, Physical/Chemical Methods", Third Edition, November 1986 and its updates.

PREPARATION METHODS SUMMARY

D9B240185

PREPARATION DESCRIPTION	PREPARATION METHOD	ANALYTICAL METHOD
Acid Digestion of Sediments, Sludges, Soils	SW846 3050B	SW846 6010B
Low Concentration Utrasonic Extraction	SW846 3550	SW846 8140
Low Concentration Utrasonic Extraction	SW846 3550	SW846 8310
Mercury sample preparation	SW846 7471A	SW846 7471A
Sonication-Low Level	SW846 8080A	SW846 8080A
Total Residue as Percent Moisture	MCAWW 160.3 MOD	MCAWW 160.3 MOD

References:

- MCAWW "Methods for Chemical Analysis of Water and Wastes", EPA-600/4-79-020, March 1983 and subsequent revisions.
- SW846 "Test Methods for Evaluating Solid Waste, Physical/Chemical Methods", Third Edition, November 1986 and its updates.

METHOD / ANALYST SUMMARY

D9B240185

ANALYTICAL METHOD	ANALYST	ANALYST ID
MCAWW 160.3 MOD	Andrea Sporleder	001971
SW846 6010B	Tracy Anderson	009690
SW846 7471A	William G. Logan	002179
SW846 8080A	Mike Schmitt	005925
SW846 8140	Matthew Graves	001801
SW846 8310	Dane Rodgers	007407

References:

- MCAWW "Methods for Chemical Analysis of Water and Wastes", EPA-600/4-79-020, March 1983 and subsequent revisions.
- SW846 "Test Methods for Evaluating Solid Waste, Physical/Chemical Methods", Third Edition, November 1986 and its updates.

SAMPLE SUMMARY

D9B240185

<u>WO #</u>	SAMPLE#	# CLIENT SAMPLE ID	DATE TIME
CR3CF	001	FL00108 SITE#4	02/22/99
CR3DN	002	FL00108 SITE#12	02/22/99
CR3DV	003	FL00108 SITE#14	02/22/99
CR3E1	004	FL00108 SITE#18	02/22/99
CR3E3	005	FL00108 SITE#22	02/22/99
CR3E6	006	FL00108 SITE#26	02/22/99
CR3E9	007	FL00108 SITE#27	02/22/99
CR3EP	008	FL00108 SITE#30	02/22/99
CR3EW	009	FL00108 SITE#60	02/22/99
CR3F5	010	FL00108 SITE#33	02/22/99

NOTE(S):

- The analytical results of the samples listed above are presented on the following pages.

- All calculations are performed before rounding to avoid round-off errors in calculated results.

- Results noted as "ND" were not detected at or above the stated limit.

- This report must not be reproduced, except in full, without the written approval of the laboratory.

- Results for the following parameters are never reported on a dry weight basis: color, corrosivity, density, flashpoint, ignitability, layers, odor,

paint filter test, pH, porosity pressure, reactivity, redox potential, specific gravity, spot tests, solids, solubility, temperature, viscosity, and weight.

Client Sample ID: FL00108 SITE#4

GC Semivolatiles

Lot-Sample #:	D9B240185-001	Work Order #: CR3CF103	Matrix SOLID
Date Sampled:	02/22/99	Date Received: 02/24/99	
Prep Date:	03/08/99	Analysis Date: 03/28/99	
Prep Batch #:	9067354	Analysis Time: 07:16	
Dilution Factor:	1		

Method.....: SW846 8080A

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	38	ug/kg
alpha-BHC	ND	38	ug/kg
beta-BHC	ND	38	ug/kg
delta-BHC	ND	38	ug/kg
gamma-BHC (Lindane)	ND	38	ug/kg
Chlordane (technical)	ND	380	ug/kg
alpha-Chlordane	ND	38	ug/kg
gamma-Chlordane	ND	38	ug/kg
Chlorobenzilate	ND	740	ug/kg
4,4'-DDD	ND	74	ug/kg
4,4'-DDE	ND	74	ug/kg
4,4'-DDT	ND	74	ug/kg
Diallate	ND	740	ug/kg
Dieldrin	ND	74	ug/kg
Endosulfan I	ND	38	ug/kg
Endosulfan II	ND	74	ug/kg
Endosulfan sulfate	ND	74	ug/kg
Endrin	ND	74	ug/kg
Endrin aldehyde	ND	74	ug/kg
Endrin ketone	ND	74	ug/kg
Heptachlor	ND	38	ug/kg
Heptachlor epoxide	ND	38	ug/kg
Isodrin	ND	74	ug/kg
Kepone	ND	1900	ug/kg
Methoxychlor	ND	380	ug/kg
Aroclor 1016	ND	740	ug/kg
Aroclor 1221	ND	740	ug/kg
Aroclor 1232	ND	740	ug/kg
Aroclor 1242	ND	740	ug/kg
Aroclor 1248	ND	740	ug/kg
Aroclor 1254	ND	740	ug/kg
Aroclor 1260	ND	740	ug/kg
Toxaphene	ND	3800	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Decachlorobiphenyl	87	(62 - 139)	-
Tetrachloro-m-xylene	60	(46 - 117)	
-			

NOTE(S):

Client Sample ID: FL00108 SITE#12

GC Semivolatiles

Lot-Sample #:	D9B240185-002	Work Order #: CR3DN	Matrix: SOLID
Date Sampled:	02/22/99	Date Received: 02/24	1/99
Prep Date:	03/08/99	Analysis Date: 03/28	3/99
Prep Batch #:	9067354	Analysis Time: 07:51	L
Dilution Factor:	1		

Method.....: SW846 8080A

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	12	ug/kg
alpha-BHC	ND	12	ug/kg
beta-BHC	ND	12	ug/kg
delta-BHC	ND	12	ug/kg
gamma-BHC (Lindane)	ND	12	ug/kg
Chlordane (technical)	ND	120	ug/kg
alpha-Chlordane	ND	12	ug/kg
gamma-Chlordane	ND	12	ug/kg
Chlorobenzilate	ND	240	ug/kg
4,4'-DDD	5.3 J	24	ug/kg
4,4'-DDE	ND	24	ug/kg
4,4'-DDT	ND	24	ug/kg
Diallate	ND	240	ug/kg
Dieldrin	ND	24	ug/kg
Endosulfan I	ND	12	ug/kg
Endosulfan II	ND	24	ug/kg
Endosulfan sulfate	ND	24	ug/kg
Endrin	ND	24	ug/kg
Endrin aldehyde	ND	24	ug/kg
Endrin ketone	ND	24	ug/kg
Heptachlor	ND	12	ug/kg
Heptachlor epoxide	ND	12	ug/kg
Isodrin	ND	24	ug/kg
Kepone	ND	600	ug/kg
Methoxychlor	ND	120	ug/kg
Aroclor 1016	ND	240	ug/kg
Aroclor 1221	ND	240	ug/kg
Aroclor 1232	ND	240	ug/kg
Aroclor 1242	ND	240	ug/kg
Aroclor 1248	ND	240	ug/kg
Aroclor 1254	ND	240	ug/kg
Aroclor 1260	ND	240	ug/kg
Toxaphene	ND	1200	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	_
Decachlorobiphenyl	90	(62 - 139)	_
Tetrachloro-m-xylene	75	(46 - 117)	

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#14

GC Semivolatiles

Lot-Sample #:	D9B240185-003	Work Order #: CR3DV103	Matrix SOLID
Date Sampled:	02/22/99	Date Received: 02/24/99	
Prep Date:	03/08/99	Analysis Date: 03/28/99	
Prep Batch #:	9067354	Analysis Time: 08:25	
Dilution Factor:	1		

Method..... SW846 8080A

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	9.6	ug/kg
alpha-BHC	ND	9.6	ug/kg
beta-BHC	ND	9.6	ug/kg
delta-BHC	ND	9.6	ug/kg
gamma-BHC (Lindane)	ND	9.6	ug/kg
Chlordane (technical)	ND	96	ug/kg
alpha-Chlordane	ND	9.6	ug/kg
gamma-Chlordane	ND	9.6	ug/kg
Chlorobenzilate	ND	190	ug/kg
4,4'-DDD	3.3 J	19	ug/kg
4,4'-DDE	ND	19	ug/kg
4,4'-DDT	ND	19	ug/kg
Diallate	ND	190	ug/kg
Dieldrin	ND	19	ug/kg
Endosulfan I	ND	9.6	ug/kg
Endosulfan II	ND	19	ug/kg
Endosulfan sulfate	ND	19	ug/kg
Endrin	ND	19	ug/kg
Endrin aldehyde	ND	19	ug/kg
Endrin ketone	ND	19	ug/kg
Heptachlor	ND	9.6	ug/kg
Heptachlor epoxide	ND	9.6	ug/kg
Isodrin	ND	19	ug/kg
Kepone	ND	470	ug/kg
Methoxychlor	ND	96	ug/kg
Aroclor 1016	ND	190	ug/kg
Aroclor 1221	ND	190	ug/kg
Aroclor 1232	ND	190	ug/kg
Aroclor 1242	ND	190	ug/kg
Aroclor 1248	ND	190	ug/kg
Aroclor 1254	ND	190	ug/kg
Aroclor 1260	ND	190	ug/kg
Toxaphene	ND	960	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Decachlorobiphenyl	85	(62 - 139)	
Tetrachloro-m-xylene	71	(46 - 117)	

NOTE (S) :

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#18

GC Semivolatiles

Lot-Sample #: D9B240185-00	04 Work Order #: CR3E1103	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/28/99	
Prep Batch #: 9067354	Analysis Time: 08:59	
Dilution Factor: 1		

Method.....: SW846 8080A

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	15	ug/kg
alpha-BHC	ND	15	ug/kg
beta-BHC	ND	15	ug/kg
delta-BHC	ND	15	ug/kg
gamma-BHC (Lindane)	ND	15	ug/kg
Chlordane (technical)	ND	150	ug/kg
alpha-Chlordane	ND	15	ug/kg
gamma-Chlordane	ND	15	ug/kg
Chlorobenzilate	ND	290	ug/kg
4,4'-DDD	ND	29	ug/kg
4,4'-DDE	ND	29	ug/kg
4,4'-DDT	ND	29	ug/kg
Diallate	ND	290	ug/kg
Dieldrin	ND	29	ug/kg
Endosulfan I	ND	15	ug/kg
Endosulfan II	ND	29	ug/kg
Endosulfan sulfate	ND	29	ug/kg
Endrin	ND	29	ug/kg
Endrin aldehyde	ND	29	ug/kg
Endrin ketone	ND	29	ug/kg
Heptachlor	ND	15	ug/kg
Heptachlor epoxide	ND	15	ug/kg
Isodrin	ND	29	ug/kg
Kepone	ND	740	ug/kg
Methoxychlor	ND	150	ug/kg
Aroclor 1016	ND	290	ug/kg
Aroclor 1221	ND	290	ug/kg
Aroclor 1232	ND	290	ug/kg
Aroclor 1242	ND	290	ug/kg
Aroclor 1248	ND	290	ug/kg
Aroclor 1254	ND	290	ug/kg
Aroclor 1260	ND	290	ug/kg
Toxaphene	ND	1500	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	_
Decachlorobiphenyl	73	(62 - 139))
Tetrachloro-m-xylene	71	(46 - 117))

NOTE (S) :

Client Sample ID: FL00108 SITE#22

GC Semivolatiles

Lot-Sample #:	D9B240185-005	Work Order #: CR3E3103	Matrix: SOLID
Date Sampled:	02/22/99	Date Received: 02/24/99	
Prep Date:	03/08/99	Analysis Date: 03/28/99	
Prep Batch #:	9067354	Analysis Time: 09:33	
Dilution Factor:	1		

Method.....: SW846 8080A

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	5.9	ug/kg
alpha-BHC	ND	5.9	ug/kg
beta-BHC	ND	5.9	ug/kg
delta-BHC	ND	5.9	ug/kg
gamma-BHC (Lindane)	ND	5.9	ug/kg
Chlordane (technical)	ND	59	ug/kg
alpha-Chlordane	ND	5.9	ug/kg
gamma-Chlordane	ND	5.9	ug/kg
Chlorobenzilate	ND	110	ug/kg
4,4'-DDD	ND	11	ug/kg
4,4'-DDE	ND	11	ug/kg
4,4'-DDT	ND	11	ug/kg
Diallate	ND	110	ug/kg
Dieldrin	ND	11	ug/kg
Endosulfan I	ND	5.9	ug/kg
Endosulfan II	ND	11	ug/kg
Endosulfan sulfate	ND	11	ug/kg
Endrin	ND	11	ug/kg
Endrin aldehyde	ND	11	ug/kg
Endrin ketone	ND	11	ug/kg
Heptachlor	ND	5.9	ug/kg
Heptachlor epoxide	ND	5.9	ug/kg
Isodrin	ND	11	ug/kg
Kepone	ND	290	ug/kg
Methoxychlor	ND	59	ug/kg
Aroclor 1016	ND	110	ug/kg
Aroclor 1221	ND	110	ug/kg
Aroclor 1232	ND	110	ug/kg
Aroclor 1242	ND	110	ug/kg
Aroclor 1248	ND	110	ug/kg
Aroclor 1254	ND	110	ug/kg
Aroclor 1260	ND	110	ug/kg
Toxaphene	ND	590	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Decachlorobiphenyl	82	(62 - 139)	
Tetrachloro-m-xylene	74	(46 - 117)	

NOTE(S):

Client Sample ID: FL00108 SITE#26

GC Semivolatiles

Lot-Sample #: D9B24	0185-006 Work Order #.	: CR3E6103	Matrix SOLID
Date Sampled: 02/22	2/99 Date Received	: 02/24/99	
Prep Date: 03/08	Analysis Date	: 03/28/99	
Prep Batch #: 90673	Analysis Time	: 10:08	
Dilution Factor: 1			

Method.....: SW846 8080A

		REPORTIN	IG
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	5.5	ug/kg
alpha-BHC	ND	5.5	ug/kg
beta-BHC	ND	5.5	ug/kg
delta-BHC	3.8 J	5.5	ug/kg
gamma-BHC (Lindane)	ND	5.5	ug/kg
Chlordane (technical)	ND	55	ug/kg
alpha-Chlordane	ND	5.5	ug/kg
gamma-Chlordane	ND	5.5	ug/kg
Chlorobenzilate	ND	110	ug/kg
4,4'-DDD	ND	11	ug/kg
4,4'-DDE	ND	11	ug/kg
4,4'-DDT	ND	11	ug/kg
Diallate	ND	110	ug/kg
Dieldrin	ND	11	ug/kg
Endosulfan I	ND	5.5	ug/kg
Endosulfan II	ND	11	ug/kg
Endosulfan sulfate	ND	11	ug/kg
Endrin	ND	11	ug/kg
Endrin aldehyde	ND	11	ug/kg
Endrin ketone	ND	11	ug/kg
Heptachlor	ND	5.5	ug/kg
Heptachlor epoxide	ND	5.5	ug/kg
Isodrin	ND	11	ug/kg
Kepone	ND	270	ug/kg
Methoxychlor	ND	55	ug/kg
Aroclor 1016	ND	110	ug/kg
Aroclor 1221	ND	110	ug/kg
Aroclor 1232	ND	110	ug/kg
Aroclor 1242	ND	110	ug/kg
Aroclor 1248	ND	110	ug/kg
Aroclor 1254	ND	110	ug/kg
Aroclor 1260	ND	110	ug/kg
Toxaphene	ND	550	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Decachlorobiphenyl	87	(62 - 13	9)
Tetrachloro-m-xylene	81	(46 - 11	7)

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#27

GC Semivolatiles

Lot-Sample #: D9B240185	5-007 Work Order #: CR3E9103	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/28/99	
Prep Batch #: 9067354	Analysis Time: 10:42	
Dilution Factor: 1	_	

Method.....: SW846 8080A

		REPORTIN	IG
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	4.9	ug/kg
alpha-BHC	ND	4.9	ug/kg
beta-BHC	ND	4.9	ug/kg
delta-BHC	ND	4.9	ug/kg
gamma-BHC (Lindane)	1.5 J	4.9	ug/kg
Chlordane (technical)	ND	49	ug/kg
alpha-Chlordane	ND	4.9	ug/kg
gamma-Chlordane	ND	4.9	ug/kg
Chlorobenzilate	ND	94	ug/kg
4,4'-DDD	ND	9.4	ug/kg
4,4'-DDE	ND	9.4	ug/kg
4,4'-DDT	ND	9.4	ug/kg
Diallate	ND	94	ug/kg
Dieldrin	ND	9.4	ug/kg
Endosulfan I	ND	4.9	ug/kg
Endosulfan II	ND	9.4	ug/kg
Endosulfan sulfate	ND	9.4	ug/kg
Endrin	ND	9.4	ug/kg
Endrin aldehyde	ND	9.4	ug/kg
Endrin ketone	ND	9.4	ug/kg
Heptachlor	ND	4.9	ug/kg
Heptachlor epoxide	ND	4.9	ug/kg
Isodrin	ND	9.4	ug/kg
Kepone	ND	240	ug/kg
Methoxychlor	ND	49	ug/kg
Aroclor 1016	ND	94	ug/kg
Aroclor 1221	ND	94	ug/kg
Aroclor 1232	ND	94	ug/kg
Aroclor 1242	ND	94	ug/kg
Aroclor 1248	ND	94	ug/kg
Aroclor 1254	ND	94	ug/kg
Aroclor 1260	ND	94	ug/kg
Toxaphene	ND	490	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Decachlorobiphenyl	88	(62 - 13	9)
Tetrachloro-m-xylene	80	(46 - 11	7)

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#30

GC Semivolatiles

Lot-Sample #: D9B240185-00	8 Work Order #: CR3EP103	Matrix: SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/28/99	
Prep Batch #: 9067354	Analysis Time: 12:24	
Dilution Factor: 1		

Method.....: SW846 8080A

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	7.2	ug/kg
alpha-BHC	ND	7.2	ug/kg
beta-BHC	ND	7.2	ug/kg
delta-BHC	ND	7.2	ug/kg
gamma-BHC (Lindane)	ND	7.2	ug/kg
Chlordane (technical)	ND	72	ug/kg
alpha-Chlordane	ND	7.2	ug/kg
gamma-Chlordane	ND	7.2	ug/kg
Chlorobenzilate	ND	140	ug/kg
4,4'-DDD	ND	14	ug/kg
4,4'-DDE	ND	14	ug/kg
4,4'-DDT	ND	14	ug/kg
Diallate	ND	140	ug/kg
Dieldrin	ND	14	ug/kg
Endosulfan I	ND	7.2	ug/kg
Endosulfan II	ND	14	ug/kg
Endosulfan sulfate	ND	14	ug/kg
Endrin	ND	14	ug/kg
Endrin aldehyde	ND	14	ug/kg
Endrin ketone	ND	14	ug/kg
Heptachlor	ND	7.2	ug/kg
Heptachlor epoxide	ND	7.2	ug/kg
Isodrin	ND	14	ug/kg
Kepone	ND	350	ug/kg
Methoxychlor	ND	72	ug/kg
Aroclor 1016	ND	140	ug/kg
Aroclor 1221	ND	140	ug/kg
Aroclor 1232	ND	140	ug/kg
Aroclor 1242	ND	140	ug/kg
Aroclor 1248	ND	140	ug/kg
Aroclor 1254	ND	140	ug/kg
Aroclor 1260	ND	140	ug/kg
Toxaphene	ND	720	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	_
Decachlorobiphenyl	89	(62 - 139))
Tetrachloro-m-xylene	77	(46 - 117))

NOTE (S) :

Client Sample ID: FL00108 SITE#60

GC Semivolatiles

Lot-Sample #: D9B240185-009	Work Order #: CR3EW103	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/28/99	
Prep Batch #: 9067354	Analysis Time: 12:59	
Dilution Factor: 1	_	

Method.....: SW846 8080A

		REPORTIN	IG
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	18	ug/kg
alpha-BHC	ND	18	ug/kg
beta-BHC	ND	18	ug/kg
delta-BHC	ND	18	ug/kg
gamma-BHC (Lindane)	ND	18	ug/kg
Chlordane (technical)	ND	180	ug/kg
alpha-Chlordane	ND	18	ug/kg
gamma-Chlordane	ND	18	ug/kg
Chlorobenzilate	ND	350	ug/kg
4,4'-DDD	ND	35	ug/kg
4,4'-DDE	ND	35	ug/kg
4,4'-DDT	ND	35	ug/kg
Diallate	ND	350	ug/kg
Dieldrin	ND	35	ug/kg
Endosulfan I	ND	18	ug/kg
Endosulfan II	ND	35	ug/kg
Endosulfan sulfate	ND	35	ug/kg
Endrin	ND	35	ug/kg
Endrin aldehyde	ND	35	ug/kg
Endrin ketone	ND	35	ug/kg
Heptachlor	ND	18	ug/kg
Heptachlor epoxide	ND	18	ug/kg
Isodrin	ND	35	ug/kg
Kepone	ND	870	ug/kg
Methoxychlor	ND	180	ug/kg
Aroclor 1016	ND	350	ug/kg
Aroclor 1221	ND	350	ug/kg
Aroclor 1232	ND	350	ug/kg
Aroclor 1242	ND	350	ug/kg
Aroclor 1248	ND	350	ug/kg
Aroclor 1254	ND	350	ug/kg
Aroclor 1260	ND	350	ug/kg
Toxaphene	ND	1800	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Decachlorobiphenyl	69	(62 - 13	9)
Tetrachloro-m-xylene	69	(46 - 11	7)

NOTE(S):

Client Sample ID: FL00108 SITE#33

GC Semivolatiles

Lot-Sample #: D9B240185-010	Work Order #: CR3F5103	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/28/99	
Prep Batch #: 9067354	Analysis Time: 13:33	
Dilution Factor: 1		

Method.....: SW846 8080A

		REPORTIN	IG
PARAMETER	RESULT	LIMIT	UNITS
Aldrin	ND	4.4	ug/kg
alpha-BHC	ND	4.4	ug/kg
beta-BHC	ND	4.4	ug/kg
delta-BHC	ND	4.4	ug/kg
gamma-BHC (Lindane)	ND	4.4	ug/kg
Chlordane (technical)	ND	44	ug/kg
alpha-Chlordane	ND	4.4	ug/kg
gamma-Chlordane	ND	4.4	ug/kg
Chlorobenzilate	ND	86	ug/kg
4,4'-DDD	ND	8.6	ug/kg
4,4'-DDE	ND	8.6	ug/kg
4,4'-DDT	ND	8.6	ug/kg
Diallate	ND	86	ug/kg
Dieldrin	ND	8.6	ug/kg
Endosulfan I	ND	4.4	ug/kg
Endosulfan II	ND	8.6	ug/kg
Endosulfan sulfate	ND	8.6	ug/kg
Endrin	ND	8.6	ug/kg
Endrin aldehyde	ND	8.6	ug/kg
Endrin ketone	ND	8.6	ug/kg
Heptachlor	ND	4.4	ug/kg
Heptachlor epoxide	ND	4.4	ug/kg
Isodrin	ND	8.6	ug/kg
Kepone	ND	220	ug/kg
Methoxychlor	ND	44	ug/kg
Aroclor 1016	ND	86	ug/kg
Aroclor 1221	ND	86	ug/kg
Aroclor 1232	ND	86	ug/kg
Aroclor 1242	ND	86	ug/kg
Aroclor 1248	ND	86	ug/kg
Aroclor 1254	ND	86	ug/kg
Aroclor 1260	ND	86	ug/kg
Toxaphene	ND	440	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Decachlorobiphenyl	85	(62 - 13	9)
Tetrachloro-m-xylene	78	(46 - 11	7)

NOTE(S):

Client Sample ID: FL00108 SITE#4

GC Semivolatiles

Lot-Sample #: D9B240185-001	Work Order #: CR3CF10H	Matrix: SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 06:47	
Dilution Factor: 1		

Method.....: SW846 8140

PARAMETER Azinphos-methyl Bolstar Chlorpyrifos Coumaphos Demeton (total) Diazinon Dichlorvos Dimethoate Disulfoton Ethoprop Ethyl parathion Fensulfothion Fenthion Malathion	RESULT ND ND ND ND ND ND ND ND ND ND ND ND ND	REPORTING LIMIT 1900 380 380 380 380 380 380 380 380 380 3	UNITS ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg
Malathion Merphos Methyl parathion Mevinphos Naled Phorate Ronnel Sulfotepp Tokuthion Trichloronate Tetrachlorvinphos (Stirophos)	ND ND ND ND ND ND ND ND ND	940 380 380 4700 7400 380 380 380 380 380 380	ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg
SURROGATE Ethyl Pirimifos Chlormefos	PERCENT <u>RECOVERY</u> 80 74	RECOVERY LIMITS (55 - 105) (50 - 150)	

NOTE (S) :

Client Sample ID: FL00108 SITE#12

GC Semivolatiles

Lot-Sample #: D9B2	240185-002 Work Order #:	CR3DN10F Matrix	: SOLID
Date Sampled: 02/2	Date Received:	02/24/99	
Prep Date: 03/0	08/99 Analysis Date:	03/10/99	
Prep Batch #: 9067	Analysis Time	07:19	
Dilution Factor: 1			

Method.....: SW846 8140

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Azinphos-methyl	ND	600	ug/kg
Bolstar	ND	120	ug/kg
Chlorpyrifos	ND	120	ug/kg
Coumaphos	ND	120	ug/kg
Demeton (total)	ND	120	ug/kg
Diazinon	ND	120	ug/kg
Dichlorvos	ND	120	ug/kg
Dimethoate	ND	120	ug/kg
Disulfoton	ND	120	ug/kg
Ethoprop	ND	120	ug/kg
Ethyl parathion	ND	120	ug/kg
Fensulfothion	ND	600	ug/kg
Fenthion	ND	120	ug/kg
Malathion	ND	300	ug/kg
Merphos	ND	120	ug/kg
Methyl parathion	ND	120	ug/kg
Mevinphos	ND	1500	ug/kg
Naled	ND	2400	ug/kg
Phorate	ND	120	ug/kg
Ronnel	ND	120	ug/kg
Sulfotepp	ND	120	ug/kg
Tokuthion	ND	120	ug/kg
Trichloronate	ND	120	ug/kg
Tetrachlorvinphos (Stirophos)	ND	600	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Ethyl Pirimifos	85	(55 - 105)	-
Chlormefos	77	(50 - 150)	

NOTE (S) :

Client Sample ID: FL00108 SITE#14

GC Semivolatiles

Lot-Sample #: D9B240185-00	3 Work Order #: CR3DV10F	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 07:52	
Dilution Factor: 1	-	

Method.....: SW846 8140

PARAMETER	RESULT	REPORTING LIMIT	UNITS
Azinphos-methyl	ND	470	ug/kg
Bolstar	ND	96	ug/kg
Chlorpyrifos	ND	96	ug/kg
Coumaphos	ND	96	ug/kg
Demeton (total)	ND	96	ug/kg
Diazinon	ND	96	ug/kg
Dichlorvos	ND	96	ug/kg
Dimethoate	ND	96	ug/kg
Disulfoton	ND	96	ug/kg
Ethoprop	ND	96	ug/kg
Ethyl parathion	ND	96	ug/kg
Fensulfothion	ND	470	ug/kg
Fenthion	ND	96	ug/kg
Malathion	ND	240	ug/kg
Merphos	ND	96	ug/kg
Methyl parathion	ND	96	ug/kg
Mevinphos	ND	1200	ug/kg
Naled	ND	1900	ug/kg
Phorate	ND	96	ug/kg
Ronnel	ND	96	ug/kg
Sulfotepp	ND	96	ug/kg
Tokuthion	ND	96	ug/kg
Trichloronate	ND	96	ug/kg
Tetrachlorvinphos (Stirophos)	ND	470	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Ethyl Pirimifos Chlormefos	86	(55 - 105)	
CHIOTHEIOS	76	(50 - 150)	

NOTE(S):

Client Sample ID: FL00108 SITE#18

GC Semivolatiles

Lot-Sample #: D9B2403	185-004 Work Order #:	CR3E110F Ma	atrix SOLID
Date Sampled: 02/22/9	99 Date Received:	02/24/99	
Prep Date: 03/08/9	99 Analysis Date	03/10/99	
Prep Batch #: 9067270	Analysis Time	08:25	
Dilution Factor: 1	_		

Method.....: SW846 8140

PARAMETER	RESULT	REPORTING LIMIT	UNITS
Azinphos-methyl	ND	740	ug/kg
Bolstar	ND	150	ug/kg
Chlorpyrifos	ND	150	ug/kg
Coumaphos	ND	150	ug/kg
Demeton (total)	ND	150	ug/kg
Diazinon	ND	150	ug/kg
Dichlorvos	ND	150	ug/kg
Dimethoate	ND	150	ug/kg
Disulfoton	ND	150	ug/kg
Ethoprop	ND	150	ug/kg
Ethyl parathion	ND	150	ug/kg
Fensulfothion	ND	740	ug/kg
Fenthion	ND	150	ug/kg
Malathion	ND	380	ug/kg
Merphos	ND	150	ug/kg
Methyl parathion	ND	150	ug/kg
Mevinphos	ND	1900	ug/kg
Naled	ND	2900	ug/kg
Phorate	ND	150	ug/kg
Ronnel	ND	150	ug/kg
Sulfotepp	ND	150	ug/kg
Tokuthion	ND	150	ug/kg
Trichloronate	ND	150	ug/kg
Tetrachlorvinphos (Stirophos)	ND	740	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Ethyl Pirimifos	88	(55 - 105)	•
Chlormefos	78	(50 - 150)	

NOTE(S):

Client Sample ID: FL00108 SITE#22

GC Semivolatiles

Lot-Sample #: D9B240185-00	5 Work Order #: CR3E310F	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 08:58	
Dilution Factor: 1	_	

Method.....: SW846 8140

PARAMETER Azinphos-methyl	RESULT	REPORTING <u>LIMIT</u> 290	UNITS
Bolstar	ND	290 59	ug/kg ug/kg
Chlorpyrifos	ND	59	ug/kg
Coumaphos	ND	59	ug/kg
Demeton (total)	ND	59	ug/kg
Diazinon	ND	59	ug/kg
Dichlorvos	ND	59	ug/kg
Dimethoate	ND	59	ug/kg
Disulfoton	ND	59	ug/kg
Ethoprop	ND	59	ug/kg
Ethyl parathion	ND	59	ug/kg
Fensulfothion	ND	290	ug/kg
Fenthion	ND	59	ug/kg
Malathion	ND	150	ug/kg
Merphos	ND	59	ug/kg
Methyl parathion	ND	59	ug/kg
Mevinphos	ND	730	ug/kg
Naled	ND	1100	ug/kg
Phorate	ND	59	ug/kg
Ronnel	ND	59	ug/kg
Sulfotepp	ND	59	ug/kg
Tokuthion	ND	59	ug/kg
Trichloronate	ND	59	ug/kg
Tetrachlorvinphos (Stirophos)	ND	290	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Ethyl Pirimifos	59	(55 - 105)	
Chlormefos	79	(50 - 100)	

NOTE(S):

Client Sample ID: FL00108 SITE#26

GC Semivolatiles

Lot-Sample #: D9B240185-006	Work Order #: CR3E610F	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 11:42	
Dilution Factor: 1		

Method..... SW846 8140

PARAMETER	RESULT	REPORTING LIMIT	UNITS
Azinphos-methyl	ND	270	ug/kg
Bolstar	ND	55	ug/kg
Chlorpyrifos	ND	55	ug/kg
Coumaphos	ND	55	ug/kg
Demeton (total)	ND	55	ug/kg
Diazinon	ND	55	ug/kg
Dichlorvos	ND	55	ug/kg
Dimethoate	ND	55	ug/kg
Disulfoton	ND	55	ug/kg
Ethoprop	ND	55	ug/kg
Ethyl parathion	ND	55	ug/kg
Fensulfothion	ND	270	ug/kg
Fenthion	ND	55	ug/kg
Malathion	ND	140	ug/kg
Merphos	ND	55	ug/kg
Methyl parathion	ND	55	ug/kg
Mevinphos	ND	680	ug/kg
Naled	ND	1100	ug/kg
Phorate	ND	55	ug/kg
Ronnel	ND	55	ug/kg
Sulfotepp	ND	55	ug/kg
Tokuthion	ND	55	ug/kg
Trichloronate	ND	55	ug/kg
Tetrachlorvinphos (Stirophos)	ND	270	ug/kg
SURROGATE	PERCENT	RECOVERY	
Ethyl Pirimifos	RECOVERY	LIMITS	
Chlormefos	85	(55 - 105)	
CHIOTHETOS	77	(50 - 150)	

NOTE(S):

Client Sample ID: FL00108 SITE#27

GC Semivolatiles

Lot-Sample #: D9B240185-007	Work Order #: CR3E910F	Matrix: SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 12:15	
Dilution Factor: 1	_	

Method.....: SW846 8140

PARAMETER	RESULT	REPORTING LIMIT	UNITS
Azinphos-methyl	ND	240	ug/kg
Bolstar	ND	49	ug/kg
Chlorpyrifos	ND	49	ug/kg
Coumaphos	ND	49	ug/kg
Demeton (total)	ND	49	ug/kg
Diazinon	ND	49	ug/kg
Dichlorvos	ND	49	ug/kg
Dimethoate	ND	49	ug/kg
Disulfoton	ND	49	ug/kg
Ethoprop	ND	49	ug/kg
Ethyl parathion	ND	49	ug/kg
Fensulfothion	ND	240	ug/kg
Fenthion	ND	49	ug/kg
Malathion	ND	120	ug/kg
Merphos	ND	49	ug/kg
Methyl parathion	ND	49	ug/kg
Mevinphos	ND	600	ug/kg
Naled	ND	940	ug/kg
Phorate	ND	49	ug/kg
Ronnel	ND	49	ug/kg
Sulfotepp	ND	49	ug/kg
Tokuthion	ND	49	ug/kg
Trichloronate	ND	49	ug/kg
Tetrachlorvinphos (Stirophos)	ND	240	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Ethyl Pirimifos Chlormefos	85	(55 - 105)	
CHIOIMEIOS	75	(50 - 150)	

NOTE (S) :

Client Sample ID: FL00108 SITE#30

GC Semivolatiles

Lot-Sample #: D9B240185-008	Work Order #: CR3EP10F	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 12:48	
Dilution Factor: 1	_	

Method.....: SW846 8140

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Azinphos-methyl	ND	350	ug/kg
Bolstar	ND	72	ug/kg
Chlorpyrifos	ND	72	ug/kg
Coumaphos	ND	72	ug/kg
Demeton (total)	ND	72	ug/kg
Diazinon	ND	72	ug/kg
Dichlorvos	ND	72	ug/kg
Dimethoate	ND	72	ug/kg
Disulfoton	ND	72	ug/kg
Ethoprop	ND	72	ug/kg
Ethyl parathion	ND	72	ug/kg
Fensulfothion	ND	350	ug/kg
Fenthion	ND	72	ug/kg
Malathion	ND	180	ug/kg
Merphos	ND	72	ug/kg
Methyl parathion	ND	72	ug/kg
Mevinphos	ND	890	ug/kg
Naled	ND	1400	ug/kg
Phorate	ND	72	ug/kg
Ronnel	ND	72	ug/kg
Sulfotepp	ND	72	ug/kg
Tokuthion	ND	72	ug/kg
Trichloronate	ND	72	ug/kg
Tetrachlorvinphos (Stirophos)	ND	350	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	-
Ethyl Pirimifos	85	(55 - 105)	
Chlormefos	77	(50 - 150)	

NOTE(S):

Client Sample ID: FL00108 SITE#60

GC Semivolatiles

Lot-Sample #: D9B240185-009	Work Order #: CR3EW10F	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 13:20	
Dilution Factor: 1	-	

Method..... SW846 8140

PARAMETER Azinphos-methyl Bolstar Chlorpyrifos Coumaphos Demeton (total) Diazinon Dichlorvos Dimethoate Disulfoton Ethoprop Ethyl parathion Fensulfothion Fenthion Malathion Merphos Methyl parathion	RESULT ND ND ND ND ND ND ND ND ND ND ND ND ND	REPORTING LIMIT 870 180 180 180 180 180 180 180 180 180 18	UNITS ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg
Methyl parathion	ND	180	ug/kg ug/kg
Naled Phorate Ronnel	ND ND	3500 180	ug/kg ug/kg
Sulfotepp Tokuthion Trichloronate	ND ND ND ND	180 180 180 180	ug/kg ug/kg ug/kg ug/kg
Tetrachlorvinphos (Stirophos)	ND PERCENT	870 RECOVERY	ug/kg
SURROGATE Ethyl Pirimifos Chlormefos	RECOVERY 83 75	<u>LIMITS</u> (55 - 105) (50 - 150)	

NOTE(S):

Client Sample ID: FL00108 SITE#33

GC Semivolatiles

Lot-Sample #: D9B240185-01	Work Order #: CR3F510F	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/10/99	
Prep Batch #: 9067270	Analysis Time: 13:53	
Dilution Factor: 1	· · · · · · · · · · · · · · · · · · ·	

Method.....: SW846 8140

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Azinphos-methyl	ND	220	ug/kg
Bolstar	ND	44	ug/kg
Chlorpyrifos	ND	44	ug/kg
Coumaphos	ND	44	ug/kg
Demeton (total)	ND	44	ug/kg
Diazinon	ND	44	ug/kg
Dichlorvos	ND	44	ug/kg
Dimethoate	ND	44	ug/kg
Disulfoton	ND	44	ug/kg
Ethoprop	ND	44	ug/kg
Ethyl parathion	ND	44	ug/kg
Fensulfothion	ND	220	ug/kg
Fenthion	ND	44	ug/kg
Malathion	ND	110	ug/kg
Merphos	ND	44	ug/kg
Methyl parathion	ND	44	ug/kg
Mevinphos	ND	550	ug/kg
Naled	ND	860	ug/kg
Phorate	ND	44	ug/kg
Ronnel	ND	44	ug/kg
Sulfotepp	ND	44	ug/kg
Tokuthion	ND	44	ug/kg
Trichloronate	ND	44	ug/kg
Tetrachlorvinphos (Stirophos)	ND	220	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Ethyl Pirimifos	83	(55 - 105)	
Chlormefos	78	(50 - 150)	

NOTE(S):

Client Sample ID: FL00108 SITE#4

HPLC

Lot-Sample #:	D9B240185-001	Work Order #:	CR3CF102	Matrix SOLID
Date Sampled:	02/22/99	Date Received:	02/24/99	
Prep Date:	02/25/99	Analysis Date:	03/12/99	
Prep Batch #:	9056304	Analysis Time:	14:51	
Dilution Factor:	2			
		Nothod	0500000000000	

Method.....: SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	8900	ug/kg
Acenaphthylene	ND	8900	ug/kg
Anthracene	130 J	890	ug/kg
Benzo(a)anthracene	150 J	890	ug/kg
Benzo(a)pyrene	ND	890	ug/kg
Benzo(b)fluoranthene	840 J	890	ug/kg
Benzo(ghi)perylene	110 J	1800	ug/kg
Benzo(k)fluoranthene	270 J	890	ug/kg
Chrysene	ND	1800	ug/kg
Dibenzo(a,h)anthracene	ND	1800	ug/kg
Fluoranthene	1100 J	1800	ug/kg
Fluorene	ND	1800	ug/kg
Indeno(1,2,3-cd)pyrene	ND	1800	ug/kg
1-Methylnaphthalene	ND	8900	ug/kg
2-Methylnaphthalene	ND	8900	ug/kg
Naphthalene	ND	8900	ug/kg
Phenanthrene	ND	1800	ug/kg
Pyrene	ND	1800	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	67	(57 - 140))

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#12

HPLC

Lot-Sample #:	D9B240185-002	Work Order #:	CR3DN102	Matrix SOLID
Date Sampled:	02/22/99	Date Received:	02/24/99	
Prep Date:	02/25/99	Analysis Date	03/12/99	
Prep Batch #:	9056304	Analysis Time:	16:21	
Dilution Factor:	2	_		
		Method	SW846 8310	

PARAMETER		REPORTING	INTERO
	RESULT	LIMIT	UNITS
Acenaphthene	ND	2900	ug/kg
Acenaphthylene	ND	2900	ug/kg
Anthracene	ND	290	ug/kg
Benzo(a)anthracene	ND	290	ug/kg
Benzo(a)pyrene	120 J	290	ug/kg
Benzo(b)fluoranthene	240 J	290	ug/kg
Benzo(ghi)perylene	ND	570	ug/kg
Benzo(k)fluoranthene	ND	290	ug/kg
Chrysene	ND	570	ug/kg
Dibenzo(a,h)anthracene	ND	570	ug/kg
Fluoranthene	230 J	570	ug/kg
Fluorene	ND	570	ug/kg
Indeno(1,2,3-cd)pyrene	ND	570	ug/kg
1-Methylnaphthalene	ND	2900	ug/kg
2-Methylnaphthalene	ND	2900	ug/kg
Naphthalene	ND	2900	ug/kg
Phenanthrene	ND	570	ug/kg
Pyrene	ND	570	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	61	(57 - 140)	

NOTE (S) :

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#14

HPLC

Lot-Sample #: D9B240185-	003 Work Order #: CR3DV102	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 02/25/99	Analysis Date: 03/12/99	
Prep Batch #: 9056304	Analysis Time: 16:50	
Dilution Factor: 2		
	Method SW846 8310	

REPORTING PARAMETER RESULT LIMIT UNITS Acenaphthene ND 2300 ug/kg Acenaphthylene ND 2300 ug/kg Anthracene ND 230 ug/kg Benzo(a) anthracene 60 J 230 ug/kg Benzo(a) pyrene 120 J 230 ug/kg Benzo(b) fluoranthene 270 230 ug/kg Benzo(ghi)perylene ND 450 ug/kg Benzo(k)fluoranthene ND 230 ug/kg Chrysene ND 450 ug/kg Dibenzo(a,h)anthracene ND 450 ug/kg Fluoranthene 210 J 450 ug/kg Fluorene ND 450 ug/kg Indeno(1,2,3-cd)pyrene ND 450 ug/kg 1-Methylnaphthalene ND 2300 ug/kg 2-Methylnaphthalene ND 2300 ug/kg Naphthalene ND 2300 ug/kg Phenanthrene ND 450 ug/kg Pyrene ND 450 ug/kg PERCENT RECOVERY SURROGATE RECOVERY LIMITS Terphenyl-d14 72 (57 - 140)

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#18

HPLC

Lot-Sample #: D9B240185-004	Work Order #: C	R3E1102 M	atrix SOLID
Date Sampled: 02/22/99	Date Received: 0.	2/24/99	
Prep Date: 02/25/99	Analysis Date: 0.	3/05/99	
Prep Batch #: 9056304	Analysis Time: 0		
Dilution Factor: 1	-		

Method.....: SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	1800	ug/kg
Acenaphthylene	ND	1800	ug/kg
Anthracene	95 J	180	ug/kg
Benzo(a)anthracene	940	180	ug/kg
Benzo(a)pyrene	ND	180	ug/kg
Benzo(b)fluoranthene	ND	180	ug/kg
Benzo(ghi)perylene	ND	360	ug/kg
Benzo(k)fluoranthene	ND	180	ug/kg
Chrysene	ND	360	ug/kg
Dibenzo(a,h)anthracene	ND	360	ug/kg
Fluoranthene	4100	360	ug/kg
Fluorene	ND	360	ug/kg
Indeno(1,2,3-cd)pyrene	ND	360	ug/kg
1-Methylnaphthalene	ND	1800	ug/kg
2-Methylnaphthalene	ND	1800	ug/kg
Naphthalene	ND	1800	ug/kg
Phenanthrene	730	360	ug/kg
Pyrene	1700	360	ug/kg
	PERCENT	RECOVERY	
SURROGATE	<u>RECOVERY</u>	LIMITS	-
Terphenyl-d14	73	(57 - 140)	

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#22

HPLC

Lot-Sample #: D9B240185-0	05 Work Order #: CR3E3102	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 02/25/99	Analysis Date. :: 03/12/99	
Prep Batch #: 9056304	Analysis Time: 17:20	
Dilution Factor: 2		

Method.....: SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	1400	ug/kg
Acenaphthylene	ND	1400	ug/kg
Anthracene	ND	140	ug/kg
Benzo(a)anthracene	ND	140	ug/kg
Benzo(a)pyrene	58 J	140	ug/kg
Benzo(b)fluoranthene	91 J	140	ug/kg
Benzo(ghi)perylene	ND	280	ug/kg
Benzo(k)fluoranthene	ND	140	ug/kg
Chrysene	ND	280	ug/kg
Dibenzo(a,h)anthracene	ND	280	ug/kg
Fluoranthene	78 J	280	ug/kg
Fluorene	ND	280	ug/kg
Indeno(1,2,3-cd)pyrene	ND	280	ug/kg
1-Methylnaphthalene	ND	1400	ug/kg
2-Methylnaphthalene	ND	1400	ug/kg
Naphthalene	ND	1400	ug/kg
Phenanthrene	ND	280	ug/kg
Pyrene	ND	280	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	66	(57 - 140)	,

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#26

HPLC

Lot-Sample #: D9B2401	185-006 Work Order #: CR3	E6102 Matrix SOLID
Date Sampled: 02/22/9	9 Date Received: 02/	24/99
Prep Date: 02/25/9	Analysis Date: 03/	05/99
Prep Batch #: 9056304	Analysis Time: 10:	03
Dilution Factor: 1		

Method.....: SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	650	ug/kg
Acenaphthylene	ND	650	ug/kg
Anthracene	ND	65	ug/kg
Benzo(a) anthracene	ND	65	ug/kg
Benzo(a)pyrene	78	65	ug/kg
Benzo(b)fluoranthene	ND	65	ug/kg
Benzo(ghi)perylene	71 J	130	ug/kg
Benzo(k)fluoranthene	40 J	65	ug/kg
Chrysene	ND	130	ug/kg
Dibenzo(a,h)anthracene	ND	130	ug/kg
Fluoranthene	110 J	130	ug/kg
Fluorene	ND	130	ug/kg
Indeno(1,2,3-cd)pyrene	ND	130	ug/kg
1-Methylnaphthalene	ND	650	ug/kg
2-Methylnaphthalene	ND	650	ug/kg
Naphthalene	ND	650	ug/kg
Phenanthrene	ND	130	ug/kg
Pyrene	ND	130	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	58	(57 - 140)	-

NOTE (S) :

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#27

HPLC

Lot-Sample #: D9B240185-0	07 Work Order #: CR3E9102	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 02/25/99	Analysis Date: 03/05/99	
Prep Batch #: 9056304	Analysis Time: 10:33	
Dilution Factor: 1		

Method.....: SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	570	ug/kg
Acenaphthylene	ND	570	ug/kg
Anthracene	ND	57	ug/kg
Benzo(a)anthracene	ND	57	ug/kg
Benzo(a)pyrene	ND	57	ug/kg
Benzo(b)fluoranthene	ND	57	ug/kg
Benzo(ghi)perylene	ND	110	ug/kg
Benzo(k)fluoranthene	21 J	57	ug/kg
Chrysene	ND	110	ug/kg
Dibenzo(a,h)anthracene	ND	110	ug/kg
Fluoranthene	58 J	110	ug/kg
Fluorene	130	110	ug/kg
Indeno(1,2,3-cd)pyrene	ND	110	ug/kg
1-Methylnaphthalene	ND	570	ug/kg
2-Methylnaphthalene	ND	570	ug/kg
Naphthalene	ND	570	ug/kg
Phenanthrene	ND	110	ug/kg
Pyrene	57 J	110	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	57	(57 - 140)	_

NOTE (S) :

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#30

HPLC

Lot-Sample #:	D9B240185-008	Work Order #: CR3EP102	Matrix SOLID
Date Sampled:	02/22/99	Date Received: 02/24/99	
Prep Date:	02/25/99	Analysis Date: 03/12/99	
Prep Batch #:	9056304	Analysis Time: 17:50	
Dilution Factor:	2		

Method.....: SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	1700	ug/kg
Acenaphthylene	ND	1700	ug/kg
Anthracene	ND	170	ug/kg
Benzo(a)anthracene	ND	170	ug/kg
Benzo(a)pyrene	ND	170	ug/kg
Benzo(b)fluoranthene	ND	170	ug/kg
Benzo(ghi)perylene	ND	340	ug/kg
Benzo(k)fluoranthene	ND	170	ug/kg
Chrysene	ND	340	ug/kg
Dibenzo(a,h)anthracene	ND	340	ug/kg
Fluoranthene	ND	340	ug/kg
Fluorene	ND	340	ug/kg
Indeno(1,2,3-cd)pyrene	ND	340	ug/kg
1-Methylnaphthalene	ND	1700	ug/kg
2-Methylnaphthalene	ND	1700	ug/kg
Naphthalene	ND	1700	ug/kg
Phenanthrene	ND	340	ug/kg
Pyrene	ND	340	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	63	(57 - 140)	-

NOTE(S):

Client Sample ID: FL00108 SITE#60

HPLC

Lot-Sample #: D9B240185-009	Work Order #: CR3EW102	Matrix SOLID
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 02/25/99	Analysis Date: 03/12/99	
Prep Batch #: 9056304	Analysis Time: 18:20	
Dilution Factor: 2	_	

Method..... SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	4200	ug/kg
Acenaphthylene	ND	4200	ug/kg
Anthracene	ND	420	ug/kg
Benzo(a)anthracene	ND	420	ug/kg
Benzo(a)pyrene	120 J	420	ug/kg
Benzo(b)fluoranthene	190 J	420	ug/kg
Benzo(ghi)perylene	ND	840	ug/kg
Benzo(k)fluoranthene	ND	420	ug/kg
Chrysene	ND	840	ug/kg
Dibenzo(a,h)anthracene	ND	840	ug/kg
Fluoranthene	ND	840	ug/kg
Fluorene	ND	840	ug/kg
Indeno(1,2,3-cd)pyrene	ND	840	ug/kg
1-Methylnaphthalene	ND	4200	ug/kg
2-Methylnaphthalene	ND	4200	ug/kg
Naphthalene	ND	4200	ug/kg
Phenanthrene	ND	840	ug/kg
Pyrene	ND	840	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	62	(57 - 140)	-

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#33

HPLC

Lot-Sample #:	D9B240185-010	Work Order #: CR3F5102	Matrix SOLID
Date Sampled:	02/22/99	Date Received: 02/24/99	
Prep Date	02/25/99	Analysis Date: 03/12/99	
Prep Batch #:	9056304	Analysis Time: 18:50	
Dilution Factor: 2	2		

Method.....: SW846 8310

		REPORTING	
PARAMETER	RESULT	LIMIT	UNITS
Acenaphthene	ND	1000	ug/kg
Acenaphthylene	ND	1000	ug/kg
Anthracene	ND	100	ug/kg
Benzo(a)anthracene	ND	100	ug/kg
Benzo(a)pyrene	ND	100	ug/kg
Benzo(b)fluoranthene	ND	100	ug/kg
Benzo(ghi)perylene	ND	210	ug/kg
Benzo(k)fluoranthene	ND	100	ug/kg
Chrysene	ND	210	ug/kg
Dibenzo(a,h)anthracene	ND	210	ug/kg
Fluoranthene	ND	210	ug/kg
Fluorene	ND	210	ug/kg
Indeno(1,2,3-cd)pyrene	ND	210	ug/kg
1-Methylnaphthalene	ND	1000	ug/kg
2-Methylnaphthalene	ND	1000	ug/kg
Naphthalene	ND	1000	ug/kg
Phenanthrene	ND	210	ug/kg
Pyrene	ND	210	ug/kg
	PERCENT	RECOVERY	
SURROGATE	RECOVERY	LIMITS	
Terphenyl-d14	68	(57 - 140)	

NOTE(S):

Client Sample ID: FL00108 SITE#4

TOTAL Metals

_	Lot-Sample #: D9B240185-001 Date Sampled: 02/22/99 Date Received: 02/24/99						
PARAMETER	RESULT	REPOR	TING UNITS	METHO	D	PREPARATION- ANALYSIS DATE	WORK ORDER #
Prep Batch # Mercury	.: 9056141 ND	0.74 Dilution	mg/kg Factor: 1		7471A Time: 11:21	02/25-02/26/99	CR3CF10E
Prep Batch # Aluminum	.: 9056181 33900	223 Dilution	mg/kg Factor: 1		6010B Time: 19:06	02/26/99	CR3CF105
Arsenic	ND	223 Dilution	mg/kg Factor: 1		6010B Time: 19:06	02/26/99	CR3CF106
Cadmium	ND	11.2 Dilution	mg/kg Factor: 1		6010B Time: 19:06	02/26/99	CR3CF107
Chromium	110	22.3 Dilution	mg/kg Factor: 1		6010B Time: 19:06	02/26/99	CR3CF108
Copper	123	44.6 Dilution 1	mg/kg Factor: 1		6010B Time: 19:06	02/26/99	CR3CF109
Iron	78800	223 Dilution 3	mg/kg Factor: 1	SW846 Analysis	6010B Time: 20:55	02/26/99	CR3CF10A
Lead	61.4 B	112 Dilution D	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:06	02/26/99	CR3CF10C
Zinc	290	44.6 Dilution 1	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:06	02/26/99	CR3CF10D

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#12

TOTAL Metals

Lot-Sample #: D9B240185-002 Date Sampled: 02/22/99 Date Received: 02/24/99						Matrix:	SOLID
PARAMETER	RESULT	REPOR		METHO	D	PREPARATION- ANALYSIS DATE	WORK ORDER #
Prep Batch # Mercury	.: 9056141 ND	0.24 Dilution	mg/kg Factor: 1		7471A Time: 11:23	02/25-02/26/99	CR3DN10E
Prep Batch # Aluminum	.: 9056181 10000	71.8 Dilution	mg/kg Factor: 1		6010B Time: 19:11	02/26/99	CR3DN105
Arsenic	ND	71.8 Dilution	mg/kg Factor: 1		6010B Time: 19:11	02/26/99	CR3DN106
Cadmium	ND	3.6 Dilution	mg/kg Factor: 1		6010B Time: 19:11	02/26/99	CR3DN107
Chromium	42.0	7.2 Dilution	mg/kg Factor: 1		6010B Time: 19:11	02/26/99	CR3DN108
Copper	34.5	14.4 Dilution	mg/kg Factor: 1		6010B Time: 19:11	02/26/99	CR3DN109
Iron	22400	71.8 Dilution	mg/kg Factor: 1		6010B Time: 21:00	02/26/99	CR3DN10A
Lead	24.0 B	35.9 Dilution	mg/kg Factor: 1		6010B Time: 19:11	02/26/99	CR3DN10C
Zinc	110	14.4 Dilution :	mg/kg Factor: 1		6010B Time: 19:11	02/26/99	CR3DN10D

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#14

TOTAL Metals

	Lot-Sample #: D9B240185-003 Date Sampled: 02/22/99 Date Received: 02/24/99						SOLID
PARAMETER	RESULT	REPOR	TING <u>UNITS</u>	METHOI	<u>)</u>	PREPARATION- ANALYSIS DATE	WORK ORDER #
Prep Batch # Mercury	.: 9056141 0.21	0.19 Dilution	mg/kg Factor: 1		7471A Time: 11:25	02/25-02/26/99	CR3DV10E
Prep Batch # Aluminum	.: 9056181 9960	56.6 Dilution	mg/kg Factor: 1		6010B Time: 19:15	02/26/99	CR3DV105
Arsenic	ND	56.6 Dilution	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:15	02/26/99	CR3DV106
Cadmium	ND	2.8 Dilution	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:15	02/26/99	CR3DV107
Chromium	42.1	5.7 Dilution	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:15	02/26/99	CR3DV108
Copper	35.1	11.3 Dilution :	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:15	02/26/99	CR3DV109
Iron	24400	56.6 Dilution :	mg/kg Factor: 1	SW846 Analysis	6010B Time: 21:04	02/26/99	CR3DV10A
Lead	24.3 B	28.3 Dilution	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:15	02/26/99	CR3DV10C
Zinc	119	11.3 Dilution 1	mg/kg Factor: 1	SW846 Analysis	6010B Time: 19:15	02/26/99	CR3DV10D

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#18

TOTAL Metals

Lot-Sample # Date Sampled			.: 02/24/99	Matrix:	SOLID	
PARAMETER	RESULT	REPORTING	G <u>UNITS</u>	METHOD	PREPARATION- ANALYSIS DATE	WORK ORDER #
Prep Batch # Mercury	.: 9056141 ND	0.29 Dilution Fact	mg/kg or: 1	SW846 7471A Analysis Time: 11:2	02/25-02/26/99 8	CR3E110E
Prep Batch # Aluminum	.: 9056181 15200	89.3 Dilution Fact	mg/kg .or: 1	SW846 6010B Analysis Time: 19:3	02/26/99	CR3E1105
Arsenic	ND	89.3 Dilution Fact	mg/kg or: 1	SW846 6010B Analysis Time: 19:3	02/26/99	CR3E1106
Cadmium	ND	4.5 Dilution Fact	mg/kg or: 1	SW846 6010B Analysis Time: 19:3	02/26/99	CR3E1107
Chromium	78.2	8.9 Dilution Fact	mg/kg or: 1	SW846 6010B Analysis Time: 19:3	02/26/99	CR3E1108
Copper	107	17.9 Dilution Fact	mg/kg or: 1	SW846 6010B Analysis Time: 19:3	02/26/99	CR3E1109
Iron	35700	89.3 Dilution Fact	mg/kg or: 1	SW846 6010B Analysis Time: 21:10	02/26/99	CR3E110A
Lead	72.2	44.6 Dilution Fact	mg/kg or: 1	SW846 6010B Analysis Time: 19:30	02/26/99	CR3E110C
Zinc	296	17.9 Dilution Facto	mg/kg pr: 1	SW846 6010B Analysis Time: 19:30	02/26/99	CR3E110D

NOTE(S):

Client Sample ID: FL00108 SITE#22

TOTAL Metals

	Lot-Sample #: D9B240185-005 Date Sampled: 02/22/99 Date Received: 02/24/99					
PARAMETER	RESULT	REPOR		METHOD	PREPARATION- ANALYSIS DATE	WORK <u>ORDER #</u>
Prep Batch # Mercury	.: 9056141 ND	0.11 Dilution	mg/kg Factor: 1	SW846 7471A Analysis Time: 11:30	02/25-02/26/99	CR3E310E
Prep Batch # Aluminum	.: 9056181 3340	34.7 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:34	02/26/99	CR3E3105
Arsenic	ND	34.7 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:34	02/26/99	CR3E3106
Cadmium	ND	1.7 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:34	02/26/99	CR3E3107
Chromium	15.3	3.5 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:34	02/26/99	CR3E3108
Copper	11.4	6.9 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:34	02/26/99	CR3E3109
Iron	7950	34.7 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 21:20	02/26/99	CR3E310A
Lead	10.1 B	17.3 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:34	02/26/99	CR3E310C
Zinc	46.3	6.9 Dilution :	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:34	02/26/99	CR3E310D

NOTE (S) :

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#26

TOTAL Metals

Lot-Sample # Date Sampled	Matrix:	SOLID					
PARAMETER	RESULT	REPORTIN	NG UNITS	METHO	D	PREPARATION- ANALYSIS DATE	WORK ORDER #
Prep Batch # . Mercury	.: 9056141 ND	0.11 Dilution Fac	mg/kg tor: 1		7471A Time: 11:37	02/25-02/26/99	CR3E610E
Prep Batch # Aluminum	.: 9056181 3060	32.5 Dilution Fac	mg/kg stor: 1		6010B Time: 19:39	02/26/99	CR3E6105
Arsenic	ND	32.5 Dilution Fac	mg/kg tor: 1		6010B Time: 19:39	02/26/99	CR3E6106
Cadmium	ND	1.6 Dilution Fac	mg/kg tor: 1		6010B Time: 19:39	02/26/99	CR3E6107
Chromium	14.7	3.3 Dilution Fac	mg/kg tor: 1		6010B Time: 19:39	02/26/99	CR3E6108
Copper	10	6.5 Dilution Fac	mg/kg tor: 1		6010B Time: 19:39	02/26/99	CR3E6109
Iron	7120	32.5 Dilution Fac	mg/kg tor: 1		6010B Time: 21:24	02/26/99	CR3E610A
Lead	8.9 B	16.3 Dilution Fac	mg/kg tor: 1		6010B Time: 19:39	02/26/99	CR3E610C
Zinc	36.4	6.5 Dilution Fac	mg/kg tor: 1		6010B Time: 19:39	02/26/99	CR3E610D

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#27

TOTAL Metals

Lot-Sample # Date Sampled	Received.	.: 02/24/	99	Matrix:	SOLID		
PARAMETER	RESULT	REPORTING LIMIT UNITS METHOD				PREPARATION- ANALYSIS DATE	WORK ORDER #
Prep Batch # Mercury	.: 9056141 ND	0.094 Dilution Fac	mg/kg stor: 1		7471A Time: 11:39	02/25-02/26/99	CR3E910E
Prep Batch # Aluminum	.: 9056181 1700	28.6 Dilution Fac	mg/kg stor: 1		6010B Time: 19:44	02/26/99	CR3E9105
Arsenic	ND	28.6 Dilution Fac	mg/kg stor: 1		6010B Time: 19:44	02/26/99	CR3E9106
Cadmium	ND	1.4 Dilution Fac	mg/kg stor: 1		6010B Time: 19:44	02/26/99	CR3E9107
Chromium	8.6	2.9 Dilution Fac	mg/kg stor: 1		6010B Time: 19:44	02/26/99	CR3E9108
Copper	9.3	5.7 Dilution Fac	mg/kg stor: 1		6010B Time: 19:44	02/26/99	CR3E9109
Iron	3540	28.6 Dilution Fac	mg/kg tor: 1		6010B Time: 21:28	02/26/99	CR3E910A
Lead	ND	14.3 Dilution Fac	mg/kg tor: 1		6010B Time: 19:44	02/26/99	CR3E910C
Zinc	25.0	5.7 Dilution Fac	mg/kg tor: 1		6010B Time: 19:44	02/26/99	CR3E910D

NOTE(S):

Client Sample ID: FL00108 SITE#30

TOTAL Metals

Lot-Sample #: D9B240185-008 Date Sampled: 02/22/99 Date Received: 02/24					99	Matrix:	SOLID
PARAMETER	RESULT	REPORT	ING UNITS	METHO	D	PREPARATION- ANALYSIS DATE	WORK <u>ORDER #</u>
Prep Batch # . Mercury	.: 9056141 ND	0.14 Dilution F	mg/kg actor: 1		7471A Time: 11:42	02/25-02/26/99	CR3EP10E
Prep Batch # Aluminum	.: 9056181 6000	42.4 Dilution F	mg/kg actor: 1		6010B Time: 19:48	02/26/99	CR3BP105
Arsenic	ND	42.4 Dilution F	mg/kg actor: 1		6010B Time: 19:48	02/26/99	CR3EP106
Cadmium	ND	2.1 Dilution Fa	mg/kg actor: 1		6010B Time: 19:48	02/26/99	CR3EP107
Chromium	30.6	4.2 Dilution Fa	mg/kg actor: 1		6010B Time: 19:48	02/26/99	CR3EP108
Copper	12.3	8.5 Dilution Fa	mg/kg actor: 1		6010B Time: 19:48	02/26/99	CR3EP109
Iron	8320	42.4 Dilution Fa	mg/kg actor: 1		6010B Time: 21:32	02/26/99	CR3EP10A
Lead	10.7 B	21.2 Dilution Fa	mg/kg actor: 1		6010B Time: 19:48	02/26/99	CR3EP10C
Zinc	29.3	8.5 Dilution Fa	mg/kg actor: 1		6010B Time: 19:48	02/26/99	CR3EP10D

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#60

TOTAL Metals

—	Dot-Sample #: D9B240185-009 Date Sampled: 02/22/99 Date Received: 02/24/99					SOLID
PARAMETER	RESULT	REPORT	TING <u>UNITS</u>	METHOD	PREPARATION- ANALYSIS DATE	WORK ORDER #
Prep Batch # Mercury	.: 9056141 ND	0.35 Dilution	mg/kg Factor: 1 `	SW846 7471A Analysis Time: 11:44	02/25-02/26/99	CR3EW10E
Prep Batch # Aluminum	.: 9056181 26300	105 Dilution	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:53	02/26/99	CR3EW105
Arsenic	ND	105 Dilution 3	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:53	02/26/99	CR3EW106
Cadmium	ND	5.3 Dilution 1	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:53	02/26/99	CR3EW107
Chromium	127	10.5 Dilution 1	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:53	02/26/99	CR3EW108
Copper	80.4	21.1 Dilution 1	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:53	02/26/99	CR3EW109
Iron	41400	105 Dilution H	mg/kg Factor: 1	SW846 6010B Analysis Time: 21:36	02/26/99	CR3EW10A
Lead	55.5	52.7 Dilution H	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:53	02/26/99	CR3EW10C
Zinc	223	21.1 Dilution H	mg/kg Factor: 1	SW846 6010B Analysis Time: 19:53	02/26/99	CR3EW10D

NOTE(S):

Client Sample ID: FL00108 SITE#33

TOTAL Metals

	ot-Sample #: D9B240185-010 Date Sampled: 02/22/99 Date Received: 02/24/99							
PARAMETER	RESULT	REPORTI	ING UNITS	METHOD	PREPARATION- ANALYSIS DATE	WORK <u>ORDER #</u>		
Prep Batch # Mercury	.: 9056141 ND	0.086 Dilution Fa	mg/kg actor: 1	SW846 7471A Analysis Time: 11	02/25-02/26/99 :46	CR3F510E		
Prep Batch # Aluminum	.: 9056181 1630	26.1 Dilution Fa	mg/kg actor: 1	SW846 6010B Analysis Time: 19	02/26/99 :58	CR3F5105		
Arsenic	ND	26.1 Dilution Fa	mg/kg .ctor: 1	SW846 6010B Analysis Time: 19	02/26/99 :58	CR3F5106		
Cadmium	ND	1.3 Dilution Fa	mg/kg ctor: 1	SW846 6010B Analysis Time: 19	02/26/99	CR3F5107		
Chromium	8.5	2.6 Dilution Fa	mg/kg ctor: 1	SW846 6010B Analysis Time: 19	02/26/99	CR3F5108		
Copper	2.7 B	5.2 Dilution Fa	mg/kg ctor: 1	SW846 6010B Analysis Time: 19	02/26/99	CR3F5109		
Iron	2220	26.1 Dilution Fa	mg/kg ctor: 1	SW846 6010B Analysis Time: 21	02/26/99	CR3F510A		
Lead	ND	13.1 Dilution Fa	mg/kg ctor: 1	SW846 6010B Analysis Time: 19:	02/26/99	CR3F510C		
Zinc	7.4	5.2 Dilution Fa	mg/kg ctor: 1	SW846 6010B Analysis Time: 19:	02/26/99 58	CR3F510D		

NOTE(S):

Results and reporting limits have been adjusted for dry weight.

Client Sample ID: FL00108 SITE#4

General Chemistry

Lot-Sample #: Date Sampled:			Order #: Received:		t rix: S	OLID
PARAMETER Percent Moisture	2010	RL 0.10 ution Fact	UNITS	METHOD MCAWW 160.3 MOD Analysis Time: 00:00	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE (S) :

RL Reporting Limit

Client Sample ID: FL00108 SITE#12

General Chemistry

Lot-Sample #: Date Sampled:			Order #: Received:		S	OLID
PARAMETER Percent Moisture	0012	RL 0.10 ution Fact	UNITS	METHOD MCAWW 160.3 MOD Analysis Time: 00:00	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE(S):

RL Reporting Limit

Client Sample ID: FL00108 SITE#14

General Chemistry

Lot-Sample #: Date Sampled:			Order #: Received:		trix: S	OLID
PARAMETER Percent Moisture	0-10	RL 0.10 ution Fac	UNITS %	METHOD MCAWW 160.3 MOD Analysis Time: 00:00	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE(S):

RL Reporting Limit

Client Sample ID: FL00108 SITE#18

General Chemistry

Lot-Sample #: Date Sampled:		Order #: Received:		trix S	OLID
PARAMETER Percent Moisture	 RL 0.10 ution Fact	UNITS	METHOD MCAWW 160.3 MOD Analysis Time: 00:00	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE (S) :

RL Reporting Limit

Client Sample ID: FL00108 SITE#22

General Chemistry

Lot-Sample #: Date Sampled:		Order #: Received:		trix: S	SOLID
PARAMETER Percent Moisture	 RL 0.10 ution Fact	_ <u>UNITS</u> % for: 1	METHOD MCAWW 160.3 MOD Analysis Time: 00:0	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE(S):

RL Reporting Limit

Client Sample ID: FL00108 SITE#26

General Chemistry

Lot-Sample #: Date Sampled:		Order #: Received:		trix S	OLID
PARAMETER Percent Moisture	 RL 0.10 ation Fact	UNITS	METHOD MCAWW 160.3 MOD Analysis Time: 00:00	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE (S) :

RL Reporting Limit

Client Sample ID: FL00108 SITE#27

General Chemistry

Lot-Sample #: D Date Sampled: 02			rder #: eceived:		trix 8	SOLID
PARAMETER Percent Moisture	RESULT 65.1 Dilu	RL 0.10 ution Facto	UNITS * pr: 1	METHOD MCAWW 160.3 MOD Analysis Time: 00:00	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE (S) :

RL Reporting Limit

Client Sample ID: FL00108 SITE#30

General Chemistry

Lot-Sample #: Date Sampled:		Order #: Received:		atrix S	OLID
PARAMETER Percent Moisture	 RL 0.10 ution Fact	UNITS	METHOD MCAWW 160.3 MOD Analysis Time: 00:0	PREPARATION- ANALYSIS DATE 03/03/99	PREP <u>BATCH #</u> 9062255

NOTE(S):

RL Reporting Limit

Client Sample ID: FL00108 SITE#60

General Chemistry

Lot-Sample #: Date Sampled:			Order #: Received:		atrix: S	SOLID
PARAMETER Percent Moisture	2010	RL 0.10 ution Fact	UNITS	METHOD MCAWW 160.3 MOD Analysis Time: 00:0	• •	PREP <u>BATCH #</u> 9062255

NOTE(S):

RL Reporting Limit

Client Sample ID: FL00108 SITE#33

General Chemistry

Lot-Sample #: Date Sampled: 0			Order #: Received:	-	trix S	OLID
PARAMETER	RESULT	RL	UNITS	METHOD	PREPARATION- ANALYSIS DATE	PREP BATCH #
Percent Moisture	61.7	0.10	\$	MCAWW 160.3 MOD	03/03/99	9062255
	Dilu	ution Fact	or: 1	Analysis Time: 00:0)	

NOTE(S):

RL Reporting Limit

QC DATA ASSOCIATION SUMMARY

D9B240185

Sample Preparation and Analysis Control Numbers

		ANALYTICAL	LEACH	PREP	
SAMPLE#	MATRIX	METHOD	BATCH #	BATCH #	MS RUN#
001	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	2020710
	SOLID	SW846 8140		9067270	9067126
002	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126
003	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126
004	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126
005	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126
006	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126

(Continued on next page)

QC DATA ASSOCIATION SUMMARY

D9B240185

Sample Preparation and Analysis Control Numbers

		ANALYTICAL	LEACH	PREP	
SAMPLE#	MATRIX	METHOD	BATCH #	BATCH #	MS RUN#
0.07					
007	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126
008	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	
	SOLID	SW846 8310			9056053
	SOLID	MCAWW 160.3 MOD		9056304	9056118
	SOLID	SW846 8140		9062255	0000000
	SOLID	5W040 0140		9067270	9067126
009	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126
010					
010	SOLID	SW846 8080A		9067354	9067175
	SOLID	SW846 7471A		9056141	9056036
	SOLID	SW846 6010B		9056181	9056053
	SOLID	SW846 8310		9056304	9056118
	SOLID	MCAWW 160.3 MOD		9062255	
	SOLID	SW846 8140		9067270	9067126

LABORATORY CONTROL SAMPLE EVALUATION REPORT

GC Semivolatiles

Client Lot #:	D9B240185	Work Order #:	CRF2K102-LCS	Matrix: SOLID
LCS Lot-Sample#:	D9C080000-354		CRF2K103-LCSD	
Prep Date:	03/08/99	Analysis Date:	03/28/99	
Prep Batch #:		Analysis Time:	06:08	
Dilution Factor:	1			

	PERCENT	RECOVERY		RPD	
PARAMETER	RECOVERY	LIMITS	RPD	LIMITS	METHOD
4,4'-DDT	72	(60 - 133)			SW846 8080A
	74	(60 - 133)	2.2	(0-16)	SW846 8080A
Aldrin	95	(73 - 129)			SW846 8080A
	95	(73 - 129)	0.27	(0-16)	SW846 8080A
Dieldrin	90	(58 - 118)			SW846 8080A
	91	(58 - 118)	0.90	(0-16)	SW846 8080A
Endrin	92	(64 - 125)			SW846 8080A
	93	(64 - 125)	1.0	(0-17)	SW846 8080A
gamma-BHC (Lindane)	36	(34 - 158)			SW846 8080A
	36	(34 - 158)	0.18	(0-15)	SW846 8080A
Heptachlor	98	(74 - 134)		• •	SW846 8080A
	98	(74 - 134)	0.54	(0-15)	SW846 8080A
		PERCENT	RECOVI	ERY	
SURROGATE		RECOVERY	LIMITS		
Decachlorobiphenyl		95	(62 -		
		97	(62 -		
Tetrachloro-m-xylene		82	(46 -	•	
-		82	(46 -		

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results. Bold print denotes control parameters

LABORATORY CONTROL SAMPLE DATA REPORT

GC Semivolatiles

Client Lot #:	D9B240185	Work Order #:	CRF2K102-LCS	Matrix SOLID
LCS Lot-Sample#:	D9C080000-354		CRF2K103-LCSD	
Prep Date	03/08/99	Analysis Date:	03/28/99	
Prep Batch #:	9067354	Analysis Time:	06:08	
Dilution Factor:	1	-		

PARAMETER	SPIKE AMOUNT	MEASURE AMOUNT	D UNITS	PERCENT RECOVERY	RPD	MERIOD
4,4'-DDT	16.7	12.1	ug/kg	72	RPD	METHOD SW846 8080A
-	16.7	12.4	ug/kg	74	2.2	SW846 8080A SW846 8080A
Aldrin	6.67	6.31	ug/kg	95	4.4	SW846 8080A
	6.67	6.33	ug/kg	95	0.27	SW846 8080A
Dieldrin	16.7	15.1	ug/kg	90	0.27	SW846 8080A
	16.7	15.2	ug/kg	91	0.90	SW846 8080A
Endrin	16.7	15.4	ug/kg	92	0.90	SW846 8080A
	16.7	15.6	ug/kg	93	1.0	SW846 8080A
gamma-BHC (Lindane)	6.67	2.42	ug/kg	36	1.0	SW846 8080A
	6.67	2.41	ug/kg	36	0.18	SW846 8080A
Heptachlor	6.67	6.50	ug/kg	98	0.10	
	6.67	6.54	ug/kg	98	0 54	SW846 8080A
	0.07	0.54	ug/ng	30	0.54	SW846 8080A
			PERCENT	RECOVERY		
SURROGATE			RECOVERY	LIMITS		
Decachlorobiphenyl			95	(62 - 139	1	
- 1			97	(62 - 139) (62 - 139)		
Tetrachloro-m-xylene			82	(62 - 13) (46 - 117)	•	
			82	(46 - 117		
				(40 - 11)	/	

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results. Bold print denotes control parameters

METHOD BLANK REPORT

GC Semivolatiles

Client Lot #: D9B240185 MB Lot-Sample #: D9C080000-354	Work Order #: CRF2K101	Matrix SOLID
Analysis Date: 03/28/99 Dilution Factor: 1	Prep Date: 03/08/99 Prep Batch #: 9067354	Analysis Time: 05:34

		REPORTING	1	
PARAMETER	RESULT	LIMIT	UNITS	METHOD
Aldrin	ND	1.7	ug/kg	SW846 8080A
alpha-BHC	ND	1.7	ug/kg	SW846 8080A
beta-BHC	ND	1.7	ug/kg	SW846 8080A
delta-BHC	ND	1.7	ug/kg	SW846 8080A
gamma-BHC (Lindane)	ND	1.7	ug/kg	SW846 8080A
Chlordane (technical)	ND	17	ug/kg	SW846 8080A
alpha-Chlordane	ND	1.7	ug/kg	SW846 8080A
gamma-Chlordane	ND	1.7	ug/kg	SW846 8080A
Chlorobenzilate	ND	33	ug/kg	SW846 8080A
4,4'-DDD	ND	3.3	ug/kg	SW846 8080A
4,4'-DDE	ND	3.3	ug/kg	SW846 8080A
4,4'-DDT	ND	3.3	ug/kg	SW846 8080A
Diallate	ND	33	ug/kg	SW846 8080A
Dieldrin	ND	3.3	ug/kg	SW846 8080A
Endosulfan I	ND	1.7	ug/kg	SW846 8080A
Endosulfan II	ND	3.3	ug/kg	SW846 8080A
Endosulfan sulfate	ND	3.3	ug/kg	SW846 8080A
Endrin	ND	3.3	ug/kg	SW846 8080A
Endrin aldehyde	ND	3.3	ug/kg	SW846 8080A
Endrin ketone	ND	3.3	ug/kg	SW846 8080A
Heptachlor	ND	1.7	ug/kg	SW846 8080A
Heptachlor epoxide	ND	1.7	ug/kg	SW846 8080A
Isodrin	ND	3.3	ug/kg	SW846 8080A
Kepone	ND	83	ug/kg	SW846 8080A
Methoxychlor	ND	17	ug/kg	SW846 8080A
Aroclor 1016	ND	33	ug/kg	SW846 8080A
Aroclor 1221	ND	33	ug/kg	SW846 8080A
Aroclor 1232	ND	33	ug/kg	SW846 8080A
Aroclor 1242	ND	33	ug/kg	SW846 8080A
Aroclor 1248	ND	33	ug/kg	SW846 8080A
Aroclor 1254	ND	33	ug/kg	SW846 8080A
Aroclor 1260	ND	33	ug/kg	SW846 8080A
Toxaphene	ND	170	ug/kg	SW846 8080A
			-979	Shorto booom
	PERCENT	RECOVERY		
SURROGATE	RECOVERY	LIMITS		
Decachlorobiphenyl	89	(62 - 139)	-	
Tetrachloro-m-xylene	82	(46 - 117)		
		,		

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

MATRIX SPIKE SAMPLE EVALUATION REPORT

GC Semivolatiles

Client Lot #:	D9B240185	Work Order #:	CR3F510G-MS	Matrix: SOLID
MS Lot-Sample #:	D9B240185-010		CR3F510H-MSD	
Date Sampled:	02/22/99	Date Received:	02/24/99	
Prep Date:		Analysis Date:	. ,	
Prep Batch #:		Analysis Time:	• •	
Dilution Factor:				

	PERCENT	RECOVERY		RPD	
PARAMETER	RECOVERY	LIMITS R	RPD	LIMITS	METHOD
Aldrin	106	(73 - 129)			SW846 8080A
	106	(73 - 129) 0	.70	(0-16)	SW846 8080A
gamma-BHC (Lindane)	39	(34 - 158)			SW846 8080A
	39	(34 - 158) 0	.31	(0-15)	SW846 8080A
4,4'-DDT	74	(60 - 133)		•	SW846 8080A
	76	(60 - 133) 2	.3	(0-16)	SW846 8080A
Dieldrin	98	(58 - 118)		••	SW846 8080A
	95	(58 - 118) 2	.5	(0-16)	SW846 8080A
Endrin	100	(64 - 125)			SW846 8080A
	98	(64 - 125) 2	.4	(0-17)	SW846 8080A
Heptachlor	96	(74 - 134)			SW846 8080A
	98	(74 - 134) 1	.4	(0-15)	SW846 8080A
		PERCENT		RECOVERY	
SURROGATE		RECOVERY		LIMITS	
Decachlorobiphenyl	_	87		(62 - 139)	<u> </u>
		87		(62 - 139)	
Tetrachloro-m-xylene		78		(46 - 117)	
-		80		(46 - 117)	
				(10 11/)	

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

MATRIX SPIKE SAMPLE DATA REPORT

GC Semivolatiles

Client Lot #: D9B240185	Work Order #: CR3F510G-MS	Matrix SOLID
MS Lot-Sample #: D9B240185-010	CR3F510H-MSD	
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 03/08/99	Analysis Date: 03/28/99	
Prep Batch #: 9067354	Analysis Time: 14:07	
Dilution Factor: 1	• • • • • • •	

	SAMPLE	SPIKE	MEASRD		PERCENT			
PARAMETER	AMOUNT	AMT	AMOUNT	UNITS	RECOVERY	RPD	METHO	D
Aldrin	ND	17.4	18.5	ug/kg	106			8080A
	ND	17.4	18.4	ug/kg	106	0.70		8080A
gamma-BHC (Lindane)	ND	17.4	6.79	ug/kg	39		SW846	8080A
	ND	17.4	6.81	ug/kg	39	0.31	SW846	8080A
4,4'-DDT	ND	43.6	32.4	ug/kg	74		SW846	8080A
	ND	43.6	33.1	ug/kg	76	2.3	SW846	8080A
Dieldrin	ND	43.6	42.7	ug/kg	98		SW846	8080A
	ND	43.6	41.6	ug/kg	95	2.5	SW846	8080A
Bndrin	ND	43.6	43.8	ug/kg	100		SW846	8080A
	ND	43.6	42.8	ug/kg	98	2.4	SW846	8080A
Heptachlor	ND	17.4	16.8	ug/kg	96		SW846	8080A
	ND	17.4	17.0	ug/kg	98	1.4	SW846	8080A
			PERCENT		RECOVERY			
SURROGATE			RECOVERY	Y	LIMITS			
Decachlorobiphenyl			87	-	(62 - 139	$\overline{)}$		
			87		(62 - 139			
Tetrachloro-m-xylene			78		(46 - 117			
			80		(46 - 117	•		

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

LABORATORY CONTROL SAMPLE EVALUATION REPORT

GC Semivolatiles

Client Lot #:		Work Order #	#: CREM31	02-LCS	Matri	x		SOLID
LCS Lot-Sample#:	D9C080000-270		CREM31	03-LCSI				
Prep Date:	03/08/99	Analysis Dat	: 03/10/	99				
Prep Batch #:	9067270	Analysis Tim						
Dilution Factor:	1	-						
	1	PERCENT	RECOVERY		RPD			
PARAMETER	1	RECOVERY	LIMITS	RPD	LIMITS	METHOI	2	
Diazinon	:	104	(52 - 139)	<u> </u>		SW846	8140	
	:	100	(52 - 139)	4.0	(0-22)	SW846	8140	
Bthyl parathion	2	228 a	(61 - 150)			SW846	8140	
	:	198 a	(61 - 150)	14	(0-23)	SW846	8140	
Malathion	1	112	(58 - 121)			SW846	8140	
	1	106	(58 - 121)	5.3	(0-16)	SW846	8140	
Methyl parathion	1	107	(67 - 115)			SW846	8140	
	1	103	(67 - 115)	3.8	(0-15)	SW846	8140	
Phorate	<u>c</u>	99	(1.0- 116)			SW846	8140	
	5	97	(1.0- 116)	1.9	(0-50)	SW846	8140	
(III) DOG MER			PERCENT	RECOVE	RY			
SURROGATE			RECOVERY	LIMITS				
Ethyl Pirimifos			101	(55 -	105)			
Ch I arrest Cart			99	(55 -	105)			
Chlormefos			90	(50 -	150)			
			^ ^	(

89

(50 - 150)

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

a Spiked analyte recovery is outside stated control limits.

LABORATORY CONTROL SAMPLE DATA REPORT

GC Semivolatiles

Client Lot #:	D9B240185	Work Order #:	CREM3102-LCS	Matrix SOLID
LCS Lot-Sample#:	D9C080000-270		CREM3103-LCSD	
Prep Date	03/08/99	Analysis Date:	03/10/99	
Prep Batch #:		Analysis Time:		
Dilution Factor:	1	-		

PARAMETER Diazinon Ethyl parathion Malathion Methyl parathion Phorate	SPIKE AMOUNT 167 167 167 167 167 167 167 167 167 167	MEASURE AMOUNT 173 166 381 a 330 a 186 177 179 172 165 162	UNITS ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg	PERCENT RECOVERY 104 100 228 198 112 106 107 103 99 97	RPD 4.0 14 5.3 3.8 1.9	<u>METHOI</u> SW846 SW846 SW846 SW846 SW846 SW846 SW846 SW846 SW846 SW846	8140 8140 8140 8140 8140 8140 8140 8140	
<u>SURROGATE</u> Ethyl Pirimifos Chlormefos			PERCENT <u>RECOVERY</u> 101 99 90 89	RECOVERY LIMITS (55 - 105 (55 - 105 (50 - 150 (50 - 150))			

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

a Spiked analyte recovery is outside stated control limits.

METHOD BLANK REPORT

GC Semivolatiles

Client Lot #: MB Lot-Sample #:		Work Order #: CREM3101	Matrix: SOLID
Analysis Date: Dilution Factor:	03/10/99	Prep Date: 03/08/99 Prep Batch #: 9067270	Analysis Time: 05:08

PARAMETER	RESULT	REPORTING	-	
Azinphos-methyl	ND	LIMIT 83	UNITS	METHOD
Bolstar	ND		ug/kg	SW846 8140
Chlorpyrifos	ND	17	ug/kg	SW846 8140
Coumaphos	ND	17	ug/kg	SW846 8140
Demeton (total)	ND	17	ug/kg	SW846 8140
Diazinon	ND	17	ug/kg	SW846 8140
Dichlorvos		17	ug/kg	SW846 8140
Dimethoate	ND	17	ug/kg	SW846 8140
Disulfoton	ND	17	ug/kg	SW846 8140
Ethoprop	ND	17	ug/kg	SW846 8140
Ethyl parathion	ND	17	ug/kg	SW846 8140
Fensulfothion	ND	17	ug/kg	SW846 8140
Fenthion	ND	83	ug/kg	SW846 8140
Malathion	ND	17	ug/kg	SW846 8140
	ND	42	ug/kg	SW846 8140
Merphos	ND	17	ug/kg	SW846 8140
Methyl parathion	ND	17	ug/kg	SW846 8140
Mevinphos	ND	210	ug/kg	SW846 8140
Naled	ND	330	ug/kg	SW846 8140
Phorate	ND	17	ug/kg	SW846 8140
Ronnel	ND	17	ug/kg	SW846 8140
Sulfotepp	ND	17	ug/kg	SW846 8140
Tokuthion	ND	17	ug/kg	SW846 8140
Trichloronate	ND	17	ug/kg	SW846 8140
Tetrachlorvinphos (Stirop	ND	83	ug/kg	SW846 8140
			5. 5	
	PERCENT	RECOVERY		
SURROGATE	RECOVERY	LIMITS		
Ethyl Pirimifos	78	(55 - 105)	
Chlormefos	83	(50 - 150		

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

MATRIX SPIKE SAMPLE EVALUATION REPORT

GC Semivolatiles

Client Lot #: D9B2401	85 Work Order #:	CR3E310G-MS	Matrix SOLID
MS Lot-Sample #: D9B2401	85-005	CR3E310H-MSD	
Date Sampled: 02/22/9	9 Date Received:	02/24/99	
Prep Date: 03/08/99		· ·	
Prep Batch #: 9067270			
Dilution Factor: 1	-		

	PERCENT	RECOVERY		RPD		
PARAMETER	RECOVERY	LIMITS	RPD	LIMITS	METHO	D
Diazinon	110	(52 - 139)			SW846	8140
	137	(52 - 139)	22	(0-22)	SW846	8140
Bthyl parathion	112	(61 - 150)			SW846	8140
	115	(61 - 150)	2.9	(0-23)	SW846	8140
Malathion	98	(58 - 121)			SW846	8140
	101	(58 - 121)	2.8	(0-16)	SW846	8140
Methyl parathion	102	(67 - 115)			SW846	8140
	104	(67 - 115)	1.9	(0-15)	SW846	8140
Phorate	82	(1.0- 116)			SW846	8140
	83	(1.0- 116)	0.79	(0-50)	SW846	8140
		PERCENT		RECOVERY		
SURROGATE		RECOVERY		LIMITS		
Ethyl Pirimifos		94		(55 - 105)	-	
		96		(55 - 105)		
Chlormefos		81		(50 - 150)		
		80		(50 - 150)		

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

Results and reporting limits have been adjusted for dry weight.

MATRIX SPIKE SAMPLE DATA REPORT

GC Semivolatiles

Client Lot #: D9B2401	85 Work Order #:	CR3E310G-MS	Matrix SOLID
MS Lot-Sample #: D9B2401		CR3E310H-MSD	
Date Sampled: 02/22/9	9 Date Received:	02/24/99	
Prep Date: 03/08/9			
Prep Batch #: 9067270	Analysis Time		
Dilution Factor: 1	_		

	SAMPLE	SPIKE	MEASRD		PERCENT			
PARAMETER	AMOUNT	AMT	AMOUNT	UNITS	RECOVERY	RPD	METHO	D
Diazinon	ND	578	636	ug/kg	110		SW846	8140
	ND	578	792	ug/kg	137	22	SW846	8140
Ethyl parathion	ND	578	645	ug/kg	112		SW846	8140
	ND	578	664	ug/kg	115	2.9	SW846	8140
Malathion	ND	578	566	ug/kg	98		SW846	8140
	ND	578	582	ug/kg	101	2.8	SW846	8140
Methyl parathion	ND	578	592	ug/kg	102		SW846	8140
	ND	578	603	ug/kg	104	1.9	SW846	8140
Phorate	ND	578	477	ug/kg	82		SW846	8140
	ND	578	480	ug/kg	83	0.79	SW846	8140
			PERCENT		RECOVERY			
SURROGATE			RECOVERY	Ľ	LIMITS			
Ethyl Pirimifos			94	-	(55 - 105	5)		
			96		(55 - 105	i)		
Chlormefos			81		(50 - 150			
			80		(50 - 150)		

NOTE (S) :

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

Results and reporting limits have been adjusted for dry weight.

LABORATORY CONTROL SAMPLE EVALUATION REPORT

HPLC

Client Lot #: D9B240185 LCS Lot-Sample#: D9B250000-30 Prep Date: 02/25/99 Prep Batch #: 9056304 Dilution Factor: 1	Analysis Da	<pre>#: CR51K1</pre>	103-LCS		x: SOLID
PARAMETER Benzo(a)pyrene	PERCENT RECOVERY 94	RECOVERY LIMITS (61 - 128)	RPD	RPD LIMITS	METHOD SW846 8310
Fluorene	101 68 91 p	(61 - 128) (63 - 138) (63 - 138)	7.1 29	(0-34) (0-22)	SW846 8310 SW846 8310 SW846 8310
Indeno(1,2,3-cd)pyrene Naphthalene	106 105 127	(69 - 127) (69 - 127) (60 - 138)	0.70	(0-15)	SW846 8310 SW846 8310 SW846 8310
Pyrene	104 93 99	(60 - 138) (59 - 136) (59 - 136)	20 6.3	(0-26) (0-20)	SW846 8310 SW846 8310 SW846 8310
SURROGATE Terphenyl-d14		PERCENT <u>RECOVERY</u> 62 64	RECOV <u>LIMIT</u> (57 - (57 -	ERY <u>S</u> 140)	

NOTE (S) :

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

p Relative percent difference (RPD) is outside stated control limits.

LABORATORY CONTROL SAMPLE DATA REPORT

HPLC

Client Lot #: D9B240185 Work Order #: CR51K102-LCS Matrix: SOLID LCS Lot-Sample#: D9B250000-304 CR51K103-LCSD Prep Date: 02/25/99 Analysis Date: 03/05/99 Prep Batch #: 9056304 Analysis Time: 05:02 Dilution Factor: 1 Analysis Time: 05:02								
	SPIKE	MEASURE	D	PERCENT				
PARAMETER	AMOUNT	AMOUNT	UNITS	RECOVERY	RPD	METHOD		
Benzo(a)pyrene	533	501	ug/kg	94		SW846 8310		
	533	538	ug/kg	101	7.1	SW846 8310		
Fluorene	533	362	ug/kg	68		SW846 8310		
	533	487 p	ug/kg	91	29	SW846 8310		
Indeno(1,2,3-cd)pyrene	533	566	ug/kg	106		SW846 8310		
Naphthalene	533	562	ug/kg	105	0.70	SW846 8310		
Napricilatelle	533	678	ug/kg	127		SW846 8310		
Pyrene	533	555	ug/kg	104	20	SW846 8310		
. yrene	533	495	ug/kg	93		SW846 8310		
	533	527	ug/kg	99	6.3	SW846 8310		
			PERCENT	DECOVEDY				
SURROGATE			RECOVERY	RECOVERY LIMITS				
Terphenyl-d14			62	(57 - 140)	<u>_</u>			
				(3) - 140	/			

64

(57 - 140)

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

p Relative percent difference (RPD) is outside stated control limits.

METHOD BLANK REPORT

HPLC

Client Lot #: D9B240185 MB Lot-Sample #: D9B250000-304	Work Order #: CR51K101	Matrix SOLID
Analysis Date: 03/05/99 Dilution Factor: 1	Prep Date: 02/25/99 Prep Batch #: 9056304	Analysis Time: 04:33

		REPORTI	ING	
PARAMETER	RESULT	LIMIT	UNITS	METHOD
Acenaphthene	ND	200	ug/kg	SW846 8310
Acenaphthylene	ND	200	ug/kg	SW846 8310
Anthracene	ND	20	ug/kg	SW846 8310
Benzo(a)anthracene	ND	20	ug/kg	SW846 8310
Benzo(a)pyrene	ND	20	ug/kg	SW846 8310
Benzo(b)fluoranthene	ND	20	ug/kg	SW846 8310
Benzo(ghi)perylene	ND	40	ug/kg	SW846 8310
Benzo(k)fluoranthene	ND	20	ug/kg	SW846 8310
Chrysene	ND	40	ug/kg	SW846 8310
Dibenzo(a,h)anthracene	ND	40	ug/kg	SW846 8310
Fluoranthene	ND	40	ug/kg	SW846 8310
Fluorene	ND	40	ug/kg	SW846 8310
Indeno(1,2,3-cd)pyrene	ND	40	ug/kg	SW846 8310
1-Methylnaphthalene	ND	200	ug/kg	SW846 8310
2-Methylnaphthalene	ND	200	ug/kg	SW846 8310
Naphthalene	ND	200	ug/kg	SW846 8310
Phenanthrene	ND	40	ug/kg	SW846 8310
Pyrene	ND	40	ug/kg	SW846 8310
				50010 0510
	PERCENT	RECOVER	Y	
SURROGATE	RECOVERY	LIMITS		
Terphenyl-d14	63	(57 - 14	40)	
<i>i</i> i				

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

MATRIX SPIKE SAMPLE EVALUATION REPORT

HPLC

Client Lot #:	D9B240185	Work Order #:	CR3CF10F-MS	Matrix SOLID
MS Lot-Sample #:			CR3CF10G-MSD	
Date Sampled:	02/22/99	Date Received:	02/24/99	
Prep Date		Analysis Date:		
Prep Batch #:		Analysis Time		
Dilution Factor:	2			

	PERCENT	RECOVERY		RPD		
PARAMETER	RECOVERY	LIMITS	RPD	LIMITS	METHO	D
Benzo(a)pyrene	73	(61 - 128)			SW846	8310
	47 a,p	(61 - 128)	42	(0-34)	SW846	8310
Fluorene	113	(63 - 138)			SW846	8310
	101	(63 - 138)	11	(0-22)	SW846	8310
Indeno (1,2,3-cd) pyrene	109	(69 - 127)			SW846	8310
	95	(69 - 127)	14	(0-15)	SW846	8310
Naphthalene	86	(60 - 138)			SW846	8310
_	64 p	(60 - 138)	29	(0-26)	SW846	8310
Pyrene	109	(59 - 136)			SW846	8310
	85 p	(59 - 136)	25	(0-20)	SW846	8310
		PERCENT		RECOVERY		
SURROGATE	_	RECOVERY		LIMITS		
Terphenyl-d14		74		(57 - 140)	
		71		(57 - 140)	

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

a Spiked analyte recovery is outside stated control limits.

p Relative percent difference (RPD) is outside stated control limits.

Results and reporting limits have been adjusted for dry weight.

MATRIX SPIKE SAMPLE DATA REPORT

HPLC

Client Lot #: D9B240185	Work Order #: CR3CF10F-MS	Matrix SOLID
MS Lot-Sample #: D9B240185-001	CR3CF10G-MSD	Solid
Date Sampled: 02/22/99	Date Received: 02/24/99	
Prep Date: 02/25/99	Analysis Date: 03/12/99	
Prep Batch #: 9056304	Analysis Time: 15:21	
Dilution Factor: 2	· · · · · · · · · · · · · · · · · · ·	

PARAMETER Benzo (a) pyrene Fluorene	SAMPLE AMOUNT ND ND ND	<u>AMT</u> 11900 11900	MEASRD AMOUNT 8660 5620 a, 13500	UNITS ug/kg ug/kg ug/kg	PERCENT RECOVERY 73 47 113	<u>RPD</u> 42	<u>METHOI</u> SW846 SW846 SW846	8310 8310
Indeno (1,2,3-cd) pyrene Naphthalene Pyrene	ND ND ND ND ND ND ND	11900 11900 11900 11900 11900 11900	12100 13000 11300 10200 7630 p 12900 10100 p	ug/kg ug/kg ug/kg ug/kg ug/kg ug/kg	101 109 95 86 64 109	11 14 29	SW846 SW846 SW846 SW846 SW846 SW846	8310 8310 8310 8310 8310 8310 8310
SURROGATE Terphenyl-d14			PERCENT RECOVERY 74 71		85 RECOVERY LIMITS (57 - 140 (57 - 140		SW846	8310

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Bold print denotes control parameters

a Spiked analyte recovery is outside stated control limits.

p Relative percent difference (RPD) is outside stated control limits.

Results and reporting limits have been adjusted for dry weight.

LABORATORY CONTROL SAMPLE EVALUATION REPORT

TOTAL Metals

Lot-Sample	#: D9B24	10185		Matrix SOLID
PARAMETER	PERCENT <u>RECOVERY</u>	RECOVERY RPD LIMITS <u>RPD</u> LIMITS	METHOD	PREPARATION - PREP - <u>ANALYSIS DATE</u> BATCH #
Mercury	98	(82 - 114)	SW846 7471A	02/25-02/26/99 9056141
	101	(82 - 114) 3.5 (0-10)	SW846 7471A	02/25-02/26/99 9056141
		Dilution Factor: 1		, =,,
Aluminum	100	(88 - 120)	SW846 6010B	02/26/99 9056181
	100	(88 - 120) 0.12 (0-10)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		. ,
Arsenic	96	(80 - 120)	SW846 6010B	02/26/99 9056181
	96	(80 - 120) 0.01 (0-10)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		
Cadmium	102	(80 - 120)	SW846 6010B	02/26/99 9056181
	101	(80 - 120) 0.26 (0-16)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		
Chromium	102	(83 - 112)	SW846 6010B	02/26/99 9056181
	102	(83 - 112) 0.25 (0-10)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		
Copper	103	(84 - 115)	SW846 6010B	02/26/99 9056181
	103	(84 - 115) 0.53 (0-10)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		
Iron	103	(87 - 117)	SW846 6010B	02/26/99 9056181
	102	(87 - 117) 0.34 (0-10)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		
Lead	101	(82 - 114)	SW846 6010B	02/26/99 9056181
	100	(82 - 114) 0.18 (0-11)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		
Zinc	99	(80 - 120)	SW846 6010B	02/26/99 9056181
	99	(80 - 120) 0.28 (0-10)	SW846 6010B	02/26/99 9056181
		Dilution Factor: 1		

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Lot-Sample #...: D9B240185

LABORATORY CONTROL SAMPLE DATA REPORT

TOTAL Metals

Lot-Sample #...: D9B240185

Matrix..... SOLID

PARAMETER Mercury	SPIKE <u>AMOUNT</u> 0.417 0.417	MEASUR <u>AMOUNT</u> 0.407 0.421		PERCNT <u>RECVRY</u> 98 101 Metor: 1	<u>RPD</u> 3.5		D 7471A 7471A	PREPARATION- <u>ANALYSIS DATE</u> 02/25-02/26/99 02/25-02/26/99	
Aluminum	200 200	200 200	mg/kg mg/kg Dilution Fa	100 100 .ctor: 1	0.12		6010B 6010B	02/26/99 02/26/99	9056181 9056181
Arsenic	200 200	192 192	mg/kg mg/kg Dilution Fa	96 96 ctor: 1	0.01		6010B 6010B	02/26/99 02/26/99	9056181 9056181
Cadmium	5.00 5.00	5.08 5.07	mg/kg mg/kg Dilution Fa	102 101 ctor: 1	0.26		6010B 6010B	02/26/99 02/26/99	9056181 9056181
Chromium	20.0 20.0	20.5 20.4	mg/kg mg/kg Dilution Fa	102 102 ctor: 1	0.25	SW846 SW846	6010B 6010B	02/26/99 02/26/99	9056181 9056181
Copper	25.0 25.0	25.8 25.7	mg/kg mg/kg Dilution Fac	103 103 ctor: 1	0.53	SW846 SW846	6010B 6010B	02/26/99 02/26/99	9056181 9056181
Iron	100 100	103 102	mg/kg mg/kg Dilution Fac		0.34	SW846 SW846		02/26/99 02/26/99	9056181 9056181
Lead	50.0 50.0	50.3 50.2	mg/kg mg/kg Dilution Fac		0.18	SW846 SW846		02/26/99 02/26/99	9056181 9056181
Zinc	50.0 50.0	49.5 49.4	mg/kg mg/kg Dilution Fac			SW846 SW846			9056181 9056181

NOTE (S) :

Calculations are performed before rounding to avoid round-off errors in calculated results.

METHOD BLANK REPORT

TOTAL Metals

Client	Lot	#:	D9B240185
--------	-----	----	-----------

Matrix..... SOLID

PARAMETER	RESULT	REPORTING LIMIT UNIT	S METHOD	PREPARATION- ANALYSIS DATE	WORK <u>ORDER #</u>
MB Lot-Sample Mercury	e #: D9B2500(ND	00-141 Prep Batch # 0.033 mg/kg Dilution Factor: 1 Analysis Time: 10:	g SW846 7471A	02/25-02/26/99	CR407101
MB Lot-Sample	e #: D9B25000	0-181 Prep Batch #			
Aluminum	ND	10.0 mg/kg Dilution Factor: 1 Analysis Time: 18::	g SW846 6010B	02/26/99	CR46912A
Arsenic	ND	10.0 mg/kg Dilution Factor: 1 Analysis Time: 18:3		02/26/99	CR46912C
Cadmium	ND	0.50 mg/kg Dilution Factor: 1 Analysis Time: 18:3		02/26/99	CR46912D
Chromium	ND	1.0 mg/kg Dilution Factor: 1 Analysis Time: 18:3		02/26/99	CR46912E
Copper	ND	2.0 mg/kg Dilution Factor: 1 Analysis Time: 18:3		02/26/99	CR46912F
Iron	1.4 B	10.0 mg/kg Dilution Factor: 1 Analysis Time: 20:2		02/26/99	CR469127
Lead	ND	5.0 mg/kg Dilution Factor: 1 Analysis Time: 18:3		02/26/99	CR469128
Zinc	ND	2.0 mg/kg Dilution Factor: 1 Analysis Time: 18:33		02/26/99	CR469129

NOTE (S) :

Calculations are performed before rounding to avoid round-off errors in calculated results.

B Estimated result. Result is less than RL.

MATRIX SPIKE SAMPLE EVALUATION REPORT

TOTAL Metals

	#: D9B24	3/99 15:15 Date Received	1: 02/24/99	Matrix	: SOLID
PARAMETER	PERCENT RECOVERY	RECOVERY RPD LIMITS RPD LIMITS	5 METHOD	PREPARATION- ANALYSIS DATE	WORK ORDER #
MS Lot-Samp)le #: D9B24	0161-002 Prep Batch #.	9056141		
Mercury	86	(82 - 114)	SW846 7471A	02/25-02/26/99	CR37R13
	86	(82 - 114) 0.74 (0-10) Dilution Factor: 1 Analysis Time: 11:1	SW846 7471A	02/25-02/26/99	
MS Lot-Samp	le #: D9B24	0161-002 Prep Batch #.	: 9056181		
Aluminum	NC,MSB	(88 - 120)	SW846 6010B	02/26/99	CR37R12F
	NC,MSB	(88 - 120) (0-10) Dilution Factor: 1 Analysis Time: 18:5		02/26/99	CR37R121
Arsenic	90	(80 - 120)	SW846 6010B	02/26/99	CR37R12V
	90	(80 - 120) 0.28 (0-10) Dilution Factor: 1 Analysis Time: 18:5	SW846 6010B	02/26/99	CR37R12V
Cadmium	94	(80 - 120)	SW846 6010B	02/26/99	CR37R130
	95	(80 - 120) 0.48 (0-16) Dilution Factor: 1 Analysis Time: 18:5	SW846 6010B	02/26/99	CR37R131
Chromium	110	(83 - 112)	SW846 6010B	02/26/99	00000100
	107	(83 - 112) 1.9 (0-10) Dilution Factor: 1 Analysis Time: 18:5	SW846 6010B	02/26/99	CR37R133 CR37R134
Copper	98	(84 - 115)	SW846 6010B	02/26/99	CR37R136
	95	(84 - 115) 1.7 (0-10) Dilution Factor: 1 Analysis Time: 18:5	SW846 6010B	02/26/99	CR37R137
Iron	NC,MSB	(87 - 117)	SW846 6010B	02/26/99	CR37R12G
	NC,MSB	(87 - 117) (0-10) Dilution Factor: 1 Analysis Time: 20:47	SW846 6010B	/ _ /	CR37R12H
ead	93	(82 - 114)	SW846 6010B	02/26/99	CR37R12K
	93	<pre>(82 - 114) 0.07 (0-11) Dilution Factor: 1 Analysis Time: 18:57</pre>	SW846 6010B		CR37R12L

(Continued on next page)

MATRIX SPIKE SAMPLE EVALUATION REPORT

TOTAL Metals

Client Lot Date Sample		40185 3/99 15:15 Date F	Received.	: 02/24/99	Matrix	: SOLID
PARAMETER Zinc	PERCENT <u>RECOVERY</u> 96 93	RECOVERY <u>LIMITS</u> <u>RPD</u> (80 - 120) (80 - 120) 1.7 Dilution Fac Analysis Tim	tor: 1	METHOD SW846 6010B SW846 6010B	PREPARATION- ANALYSIS DATE 02/26/99 02/26/99	WORK ORDER # CR37R12N CR37R12P
NOTE (S) :						

Calculations are performed before rounding to avoid round-off errors in calculated results.

Results and reporting limits have been adjusted for dry weight.

NC The recovery and/or RPD were not calculated.

MSB The recovery and RPD were not calculated because the sample amount was greater than four times the spike amount.

MATRIX SPIKE SAMPLE DATA REPORT

TOTAL Metals

Client Lot #...: D9B240185 Date Sampled...: 02/23/99 15:15 Date Received..: 02/24/99

Matrix..... SOLID

<u>PARAMETER</u> Lead			MEASURED AMOUNT	UNITS	PERCNT RECVRY	RPD	METHO	D	PREPARATION- ANALYSIS DATE	WORK ORDER #
	6.5 6.5	58.7 58.7		mg/kg mg/kg ion Factor: 1 sis Time: 18	93 93 :57	0.07	SW846 SW846		02/26/99 02/26/99	CR37R12K CR37R12L
Zinc	34.4 34.4	58.7 58.7		Mg/kg Mg/kg ion Factor: 1 sis Time: 18:	96 93 :57	1.7	SW846 SW846		02/26/99 02/26/99	CR37R12N CR37R12P

NOTE(S):

Calculations are performed before rounding to avoid round-off errors in calculated results.

Results and reporting limits have been adjusted for dry weight.

NC The recovery and/or RPD were not calculated.

MSB The recovery and RPD were not calculated because the sample amount was greater than four times the spike amount.

MATRIX SPIKE SAMPLE DATA REPORT

TOTAL Metals

Client Lo Date Samp				.5 : 15	Date Rece	ived:	02/24/	' 99	Matr	ix: SOI	JD
PARAMETER	SAMPLE AMOUNT		MEAS AMOU		UNITS	PERCI	NT RY RPD	METHO		PREPARATION - <u>ANALYSIS</u> DATE	WORK ORDER \$
MS Lot-Sa Mercury	mple #:	D9B24	0161-	002	Prep Batc	h #:	905614	1			
-	ND ND	0.489 0.489		0 Dilu	mg/kg mg/kg tion Factor: ysis Time:		0.74		7471A 7471A	02/25-02/26/99 02/25-02/26/99	
MS Lot-Sa Aluminum	mple #:	D9B24	0161-	002	Prep Batch	n #:	905618	1			
	3830	235	6990	Qual	mg/kg .ifiers: NC	C,MSB		SW846	6010B	02/26/99	CR37R12
	3830	235	6900	Qual Dilut	mg/kg ifiers: NC ion Factor: vsis Time:	C,MSB 1		SW846	6010B	02/26/99	CR37R12
Arsenic											
	5.1 5.1	235 235	216 217		mg/kg mg/kg ion Factor: : sis Time: :		0.28		6010B 6010B	02/26/99 02/26/99	CR37R12 CR37R12
Cadmium											
			5.62 5.65		mg/kg mg/kg ion Factor: 1 sis Time: 1		0.48		6010B 6010B	02/26/99 02/26/99	CR37R13(CR37R133
Chromium											
			32.5 31.9		mg/kg mg/kg ion Factor: 1 sis Time: 1			SW846 SW846			CR37R133 CR37R134
Copper											
		29.4 29.4	42.7		mg/kg mg/kg ion Factor: 1 sis Time: 1			SW846 SW846			CR37R136 CR37R137
ron											
:	10700 1	117 1	11900 (Quali	mg/kg fiers: NC,	MSB	:	SW846	6010B	02/26/99	CR37R12G
:	L0700 1	17 1	L1000 (Quali Diluti	mg/kg fiers: NC, on Factor: 1 is Time: 2	MSB	:	SW846	6010B	02/26/99	CR37R12H

State Strephon Active Strephon State State Zip Code State State Zip Code State Zip 23 Steph 4 - 2 Steph 12-2 Steph 12-3 Steph 12-3 Steph 12-3 Steph 12-3 Steph 12-3 Steph 12-3 Steph 12-3	Project Manager Telephone Number (Area Code Site Contact Carrier/Waybill Number Carrier/Waybill Number Carrier/Waybill Number	The second states of the secon	Date Date 22-22-99 22-22-99 Lab Number more space is needed X X	Clear Control of Clear Conditions of Receipt
$\frac{1}{2}$ Shot 1/4 - 3 Identification				ssed if samples are retained
X Non-Hazard 🗌 Flammable 🗌 Skin Irritant 🗍 Poison B Turn Around Time Required 24 Hours 🗌 48 Hours 🗍 7 Days 🗍 14 Days 🗍 27 C Relinnuished Bv	Days Other	C Requirements (Specify)	(A lee may be asse Months longer than 3 mont	amples are n
1. rejuquished by UNOLLED Level 2. Relinquished By	Date Time Date Time	1. Received By 2. Received By		Date Time Date Time
3. Relinquished By Comments	Date	3. Accepted by La		Date Time $OZZY^{d_1}$

Q

DISTRIBUTION: WHITE - Stays with the Sample; CANARY - Returned to Client with Report; PINK - Field Copy

Quanterra	Glarts 245G	-22-99 Chain of Custody Number 2336			Sandid Instruction	Conditions of Receipt	A. A. C. C. C.	The prize the										A	1	longer than 3 months)		Date	Date Time	Date Time	010 116170
		rn Sud-	Fax Number 24 119 Lab Number	ab Contact Attach list if more space is needed		Containers & M/ OC	S, B, HOEN /JUZ HOEN IOH SONH		× × ×	X		×××		***	X	X	×			U Disposal By Lab U Archive For Months		1. Received By	2. Received By	3. Received BD/ /	AND IN
		Project Manager	Telephone Number (Acea Gode)	Site Contact	Carrier/Waybill Number	Matrix	Times Incertification Incertif					-35-			-55-				Sample Disposal		Doner Entra it	-7-y	Time	Date Time	
Custody Record	DUA.4124 0797	I Uns. Evelosinal Sunn	IIO Eizenliche	م آ		ପ Contract/Purchase Order/Quote No. ପ	C Sample I.D. No. and Description (Containers for each sample may be combined on one line) Date	FLECIOS SILON 18-1 2-22	8 FL CO10 8 51 # 18-2 1	#7-200107 Site# (8-3 11	or	FLOONOS STE#22-1 2-12	FLOODS She # 12-2 4	FLOCIOS Site # 23-3 1	RED 105 She # 24 / 2225	FLOOTOS Site # 26-2 4	FLOODS Site# 26-3 4		Possible Hazard Identification	e rianimatie La Sain milain. Le Poison B	24 Hours 3 48 Hours 7 Days 14 Days 21 Days	Relinquished By	2. Relinquished By	3. Relinquished By	C _i coments

UDISTRIBUTION: WHITE - Stays with the Sample; CANARY - Returned to Client with Report; PINK - Field Copy

uchain of Custody Record				erra
LUA-4124 0797			(tao	Turte Der Selo
HILLE Erclerice Second	Project Manager	Ernaute	Date Chain o	Chain of Custody Number 2338
MO Esendone	lumber (Area Code)/	Fax Number 9736 CEN/19	Lab Number	2 " 2
1 a very PY 33630	4	Lab Contact Ana More more	Analysis (Attach list if more space is needed)	
B	Carrier/Waybill Number		ोष	
Contract/Purchase Order/Quote No.	Matrix	Containers &		Special Instructions/ Conditions of Receipt
Containers for each sample I.D. No. and Description () (Containers for each sample may be combined on one line)	Lio2	#OSTH	370U	
FLOO108 SHO#27-1 2-22-99				Al, Ar Cd, Cr. Cue
700103 Sta# 27-2		X		Ph 2m He
TUDIOS SIA# 37-3 >		×		1
100				
10 MO# 20-				
20-11 # 20-C				
A <- nc # all (100 7)	*	×-		
FLOORDS SHAFOUN		×		
700008 54 400-2		×		
FLEOROD STE# 60-3				()
Docarbe Level of the second				
n mmable 🗌 Skin Irritant 🔲 Poison B	Sample Disposal Content Client	Disposal By Lab	(A lee may be assessed if samples are retained Months honorer than 3 months)	mples are retained
4 Days 🗌 21 Days	- T	Spe		
Xeless J	J-J2-99 Time	1. Received By	Date	Time
	Date Time	2. Received By	Date	. Time
thed By	Date Time	3. Received BA	Date Date	Date Time
CGarnents C Carents C Carents C C Contract C C C C C C C C C C C C C C C C C C C				
ONSTRIBUTION: WHITE - Stays with the Sample; CANARY - Returned to Client with Report; PINK - Field Copy	with Report; PINK - Field Copy			

Q

AIBUTION: WHITE - Stays with the Sample; CANARY - Returned to Client with Report; PINK - Field Copy

|--|

DISTRIBUTION: WHITE . Stays with the Sample; CANARY - Returned to Client with Report; PINK - Field Copy

					Quanterra	- Denver
		•~	\sim	Sample Receiving Checklist		
Lo	ot #:_	(U	B240185 Date/Time Received: 022499	0900	
Company Name & Sampling Site: US65 Floridu						
*Cooler #(s):						
	Tem	perati	ires:	36		
Unpacking & Labeling Check Points:						
NIA		No				Initials
	Q	í 🗅	1.	Cooler seals intact.	-	PK
	6		2.	Chain of custody present.	_	<u> </u>
		Ø	3.	Bottles broken and/or are leaking, comment if yes.	_	
PHOTOGRAPH BROKEN BOTTLES						
	Ø	a	4.	Containers labeled, comment if no.		
	Ø	ū	5.	pH of all samples checked and meet requirements, note exceptions.		
	ď	۵	6.	Chain of custody includes "received by" and "relinquished" by signatures, dates, and times.	_	
	Ø	Q	7.	Receipt date(s) > 48 hours past the collection date(s)? If yes, notify PA/PM.		
	Ø		8.	Chain of custody agrees with bottle count, comment if no.		
	ď		9.	Chain of custody agrees with labels, comment if no.		
đ			10.	VOA samples filled completely, comment if no.		
A	ū	Q	11.	VOA bottles preserved, check for labels.		
		ø	12.	Did samples require preservation with sodium thiosulfate?	_	
		Ø	13.	If yes to #12, did the samples contain residual chlorine?	_	
		Q r	14.	Sediment present in "D," dissolved, bottles.		
		Ø	15.	Are analyses with short holding times requested?		
Q	Q	Ø	16.	Is extra sample volume provided for MS, MSD or matrix duplicates?		
	Q	ď	17.	Multiphase samples present? If yes, comment below.		
		ď	18. /	Any subsampling for volatiles? If yes, list samples.		
				PHOTOGRAPH MULTIPHASE SAMPLES		
ď	C	a	19.	Clear picture taken, labeled, and stapled to project folder.		
ଘ				Subcontract COC signed and sent with samples to bottle prep?		
				Was sample labeling double checked by a second person?		V

Document any problems or discrepancies and the actions taken to resolve them on a Condition Upon Receipt Anomaly Report (CUR)


\QA\Forms\Sample Receiving\SR Checklist

,

2/10/99 Revision

.


OPEN-FILE REPORT 99-226



Albers Equal-Area Conic projection Standard Parallels 29° 30' and 45° 30', central meridian –83° 00'

> LOCATION OF PREVIOUS BOTTOM-SEDIMENT SAMPLING SITES (1982-1997), AND PRESENT BOTTOM-SEDIMENT SAMPLING SITES IN THE CALOOSAHATCHEE RIVER AND ESTUARY STUDY AREA, LEE COUNTY, FLORIDA, JULY 20-30, 1998






Base from U.S. Geological Survey digital data, 1:100,000, 1983 Albers Equal-Area Conic projection Standard Parallels 29° 30' and 45° 30', central meridian –83° 00'

DISTRIBUTION OF BERYLLIUM-7 IN BOTTOM SEDIMENT IN THE CALOOSAHATCHEE RIVER AND ESTUARY STUDY AREA, LEE COUNTY, FLORIDA, JULY 20-30, 1998

OPEN-FILE REPORT 99-226


Distribution of trace elements above Natural Range - Plate 3



Albers Equal Area Conic projection Standard Parallels 29° 30' and 45° 30', central meridian -83° 00'

> DISTRIBUTION OF TRACE ELEMENTS ABOVE NATURAL RANGE IN BOTTOM SEDIMENT IN THE CALOOSAHATCHEE RIVER AND ESTUARY STUDY AREA, LEE COUNTY, FLORIDA, JULY 20-30, 1998






Albers Equal-Area Conic projection

Standard Parallels 29° 30' and 45° 30', central meridian -83° 00'

DISTRIBUTION OF TOTAL ORGANIC CARBON IN BOTTOM SEDIMENT IN THE CALOOSAHATCHEE RIVER AND ESTUARY STUDY AREA, LEE COUNTY, FLORIDA, JULY 20-30, 1998




Base from U.S. Geological Survey digital data, 1:100,000, 1983 Albers Equal-Area Conic projection Standard Parallels 29° 30' and 45° 30', central meridian –83° 00'

> DISTRIBUTION OF DDT ENZYME-LINKED IMMUNOSORBENT ASSAY DATA IN BOTTOM SEDIMENT IN THE CALOOSAHATCHEE RIVER AND ESTUARY STUDY AREA, LEE COUNTY, FLORIDA, JULY 20-30, 1998

OPEN-FILE REPORT 99-226






Albers Equal-Area Conic projection

Standard Parallels 29° 30' and 45° 30', central meridian -83° 00'

DISTRIBUTION OF CHLORDANE ENZYME-LINKED IMMUNOSORBENT ASSAY DATA IN BOTTOM SEDIMENT IN THE CALOOSAHATCHEE RIVER AND ESTUARY STUDY AREA, LEE COUNTY, FLORIDA, JULY 20-30, 1998



Albers Equal-Area Conic projection Standard Parallels 29° 30' and 45° 30', central meridian –83° 00'

DISTRIBUTION OF CARCINOGENIC POLYCYCLIC AROMATIC HYDROCARBONS ENZYME-LINKED IMMUNOSORBENT ASSAY DATA IN BOTTOM SEDIMENT IN THE CALOOSAHATCHEE RIVER AND ESTUARY STUDY AREA, LEE COUNTY, FLORIDA, JULY 20-30, 1998