

# Identification of Genes Associated with Morphology in *Aspergillus niger* by Using Suppression Subtractive Hybridization

Ziyu Dai, Xingxue Mao, Jon K. Magnuson, and Linda L. Lasure\*

Chemical and Biological Processes Development Group, Process Science & Engineering Division,  
Pacific Northwest National Laboratory, Richland, Washington 99352

Received 19 September 2003/Accepted 29 December 2003

The morphology of citric acid production strains of *Aspergillus niger* is sensitive to a variety of factors, including the concentration of manganese ( $Mn^{2+}$ ). Upon increasing the  $Mn^{2+}$  concentration in *A. niger* (ATCC 11414) cultures to 14 ppb or higher, the morphology switches from pelleted to filamentous, accompanied by a rapid decline in citric acid production. The molecular mechanisms through which  $Mn^{2+}$  exerts effects on morphology and citric acid production in *A. niger* cultures have not been well defined, but our use of suppression subtractive hybridization has identified 22 genes responsive to  $Mn^{2+}$ . Fifteen genes were differentially expressed when *A. niger* was grown in media containing 1,000 ppb of  $Mn^{2+}$  (filamentous form), and seven genes were expressed in 10 ppb of  $Mn^{2+}$  (pelleted form). Of the 15 filament-associated genes, seven are novel and eight share 47 to 100% identity with genes from other organisms. Five of the pellet-associated genes are novel, and the other two genes encode a pepsin-type protease and polyubiquitin. All 10 genes with deduced functions are either involved in amino acid metabolism-protein catabolism or cell regulatory processes. Northern blot analysis showed that the transcripts of all 22 genes were rapidly enhanced or suppressed by  $Mn^{2+}$ . Steady-state mRNA levels of six selected filament-associated genes remained high during 5 days of culture in a filamentous state and remained low under pelleted growth conditions. The opposite behavior was observed for four selected pellet-associated genes. The full-length cDNA of the filament-associated clone, *Brsa-25*, was isolated. Antisense expression of *Brsa-25* permitted pelleted growth and increased citrate production at concentrations of  $Mn^{2+}$  that were higher than the parent strain could tolerate. These results suggest the involvement of the newly isolated genes in the regulation of *A. niger* morphology.

The morphology of filamentous fungi in fermentation processes is critical to maximum product output. The optimal morphology for the production of organic acids, enzymes, and secondary metabolites differs among fungi, but growth as small pellets is usually correlated with highly efficient fungal processes. For example, pelleted morphology is necessary for maximum production of citric acid by *Aspergillus niger* (9), itaconic acid by *Aspergillus terreus* (30), pravastatin precursor by *Penicillium citrinum* (17, 47), and certain heterologous proteins by *A. niger* (57). It has been reported that filamentous growth is preferable for penicillin production by *Penicillium chrysogenum* (49) and fumaric acid production by *Rhizopus arrhizus* (6). The ability to obtain and maintain a particular morphology is one of the key parameters in the development of productive fungal fermentations. Empirically determined process conditions, such as agitation, dissolved oxygen concentration, substrate (carbon) concentration, nitrogen, phosphorous, and micronutrient concentrations, pH, ionic strength, and inoculum concentration have all been demonstrated to have effects on morphology which differ among different fungi (5, 10, 22, 29, 53). Decreasing mass transfer limitations is a likely beneficial effect of fungi adopting a small-pellet morphology (less than approximately 1 mm in diameter) in submerged fermentations. The small pellets decrease the culture viscosity (47), which increases the efficiency of mixing and thus mass transfer. In the cited study, the product output declined precipitously at pellet

diameters greater than 1 mm. Michel et al. showed that oxygen concentration falls rapidly with the depth below the surface of pellets of *Phanerochaete chrysosporium* (31). The exact depth varied with the external oxygen concentration but did not exceed 0.8 mm. These studies imply that there is a practical upper limit for pellet size associated with high product output in aerobic bioprocesses (31). Despite the known benefits of proper fungal morphology, the molecular mechanisms involved in the regulation of the morphology of filamentous fungi in submerged culture remain inadequately defined. Knowledge of the genes and enzymes involved in the complex process of fungal morphology determination is a prerequisite for the application of genetic engineering to the control of morphology in fungi. It is our hypothesis that this preferred morphology in the filamentous fungi, though induced by various nutritional or environmental conditions, is controlled by common genetic factors. To test this hypothesis, we have chosen a citric acid-producing strain of *A. niger* as the model organism.

Industrial strains of *A. niger* are capable of growing on solutions in excess of 20% (wt/vol) glucose or sucrose and converting approximately 90% of the supplied carbohydrate to citric acid. These remarkable properties are the reason that *A. niger* has been used to produce citric acid for 80 years and is currently the primary source of commercial citric acid production (28). This complex bioprocess is known to depend on a variety of environmental factors, including the concentration of  $Mn^{2+}$  in the medium. The effects of  $Mn^{2+}$  on citric acid production, cell wall composition, and morphology have been examined by using biochemical approaches. Röhr and Kubicek

\* Corresponding author. Mailing address: Chemical and Biological Process Development, Pacific Northwest National Laboratory, 902 Battelle Blvd., P.O. Box 999, MSIN K2-12, Richland, WA 99352.

(41) found that *A. niger* produced reasonable amounts of citric acid only when the  $Mn^{2+}$  concentration in the culture medium was well below 1  $\mu M$  (55 ppb). Manganese deficiency leads to an increase in protein turnover, which ultimately leads to a high intracellular concentration of  $NH_4^+$  (23, 27). The high  $NH_4^+$  concentration prevents citrate-mediated feedback inhibition of glucose catabolism (15, 16), thus allowing citric acid accumulation. In addition,  $Mn^{2+}$  deficiency results in peculiar morphology development characterized by increased spore swelling and squat, bulbous hyphae (46). An analysis of cell wall compositions from cultures grown with or without adequate manganese revealed that  $Mn^{2+}$ -deficient cultures had increased amounts of chitin but decreased amounts of  $\beta$ -glucan and galactans (21). Despite the interest in the regulation of citric acid biosynthesis in *A. niger*, the molecular mechanisms responsible for the effects of manganese on morphology formation and citric acid production in liquid culture have not been studied in detail.

Suppression subtractive hybridization (SSH) is a method that utilizes a suppressive PCR to create cDNA libraries from which the cDNAs common to two different physiological states of an organism are subtracted, thus allowing the identification of genes differentially expressed in response to an experimental stimulus (13, 14, 54). The SSH method differs from earlier subtractive methods by including a normalization step that equalizes the relative abundance of cDNA within a target population. This modification enhances the probability of identifying the increased expression of low-abundance transcripts and represents a potential advantage over other methods for identifying differentially regulated genes, such as differential display reverse transcriptase PCR (26) and cDNA representation difference analysis (50). Here, we describe the application of SSH for the identification of genetic elements associated with pelleted and filamentous morphologies, observed as  $Mn^{2+}$ -induced and  $Mn^{2+}$ -suppressed genes in *A. niger*. The responses of the newly isolated genes to different developmental stages during the fermentation processes were examined by RNA blot analysis. The full-length *Brsa-25* gene was isolated, and its effects on *A. niger* morphology and citric acid production were examined by using antisense expression.

#### MATERIALS AND METHODS

**Strains and media.** *Escherichia coli* strains JM109 and DH5 $\alpha$  were used as hosts for routine cloning experiments. *Agrobacterium tumefaciens* AGL0, containing a Bo542 chromosomal background and a disabled helper-Ti plasmid pEHA101 (25), was used for the transformation of *A. niger*. *A. niger* strain ATCC 11414, obtained from the American Type Culture Collection (Rockville, Md.), was grown on potato dextrose agar plates at 30°C for culture maintenance and spore preparation. The cultures were incubated for 5 days, and the spores were harvested by washing with sterile 0.8% Tween 80 (polyoxyethylenesorbitan monooleate). Conidia were enumerated with a hemacytometer. Aliquots of the resulting spore suspension ( $10^9$  spores/ml) were used to inoculate culture tube or baffled-flask liquid cultures. The citric acid production (CAP) medium contained 140 g of glucose/liter, 3.1 g of  $NH_4NO_3$ /liter, 0.15 g of  $KH_2PO_4$ /liter, 0.15 g of NaCl/liter, 2.2 g of  $MgSO_4 \cdot 7H_2O$ /liter, 6.6 mg of  $ZnSO_4 \cdot 7H_2O$ /liter, and 0.1 mg of  $FeCl_3$ /liter adjusted to pH 2.0 with 4 M  $H_2SO_4$ . Cations were removed from the glucose solution by ion exchange on Dowex 50W-X8, 100/200-mesh, H cation exchange resin (Fisher Scientific, Pittsburgh, Pa.) prior to adding the other nutrient components. The manganese concentration in the medium was adjusted by the addition of appropriate volumes of a stock solution of  $MnCl_2 \cdot 4H_2O$  (10 mM). The  $Mn^{2+}$  concentration in the medium, before and after growth of *A. niger*, was determined by using a Hewlett-Packard 4500 series inductively coupled plasma mass spectrometer (ICP-MS) with a sub-part per billion detection limit

(Agilent Technologies, Palo Alto, Calif.). The samples and the manganese standard solutions were serially diluted to optimal mass ranges with ultrapure deionized water before being injected into the ICP-MS for measurement. Three replicates of each sample and standard were measured. Concentrations of manganese in the samples were calculated based on the signal response of the manganese standards.

**Culture methods.** Glass baffled flasks of 250 and 1,000 ml and 16- by 125-mm glass culture tubes were silanized with SigmaCote (Sigma, St. Louis, Mo.) to minimize leaching of metals. For citric acid production tests, *A. niger* was grown in 50 ml of CAP media containing 10 or 1,000 ppb of  $Mn^{2+}$  in 250-ml baffled flasks at 30°C and 250 rpm. Samples for citric acid analysis were taken at intervals. Small cultures for examining the effects of  $Mn^{2+}$  on morphology and citric acid production were grown in 16- by 125-mm culture tubes containing 2 ml of CAP medium and incubated at 30°C and 250 rpm for 3 days. The culture tubes were laid at an angle of approximately 20° against the platform of the shaker.

To produce sufficient biomass for RNA isolation, 12 1-liter baffled flasks containing 250 ml of CAP medium with 10 ppb of  $Mn^{2+}$  were used. Each flask was inoculated with  $10^6$  spores/ml and incubated for 12 h at 30°C and 250 rpm to obtain pelleted morphology, and then 1,000 ppb of  $Mn^{2+}$  was added to six of the flasks to induce filamentous growth. This procedure was replicated four times to obtain time points 20, 40, 60, and 120 min after manganese induction of filamentous growth. At each time point, the growth of fungal cultures was suspended by rapid cooling in an ice water bath. The biomass was immediately separated from the culture supernatant by centrifugation for 10 min at 4°C and  $15,000 \times g$ . The biomass was transferred from 500-ml centrifugation bottles to 50-ml centrifugation tubes, immediately frozen in liquid  $N_2$  for 5 min, and stored at -80°C. The biomass was also prepared from 50-ml cultures with or without 1,000 ppb of  $Mn^{2+}$  at different developmental stages (0.5 to 5 days) for examining the expression patterns of newly isolated genes. The biomass from these cultures was collected by centrifugation at  $9,500 \times g$  in 50-ml centrifuge tubes. The biomass of transgenic clones was prepared from cultures grown in 16- by 125-mm glass culture tubes. The biomass was collected by centrifugation in a 1.8-ml microcentrifuge tube at  $20,000 \times g$  and 4°C for 5 min.

**Citric acid measurements.** Citric acid concentrations were determined with an endpoint spectrophotometric enzyme assay (4). Five microliters of each culture supernatant was assayed (see above).

**RNA isolation.** Total RNA was isolated from *A. niger* according to the modified acid-guanidinium isothiocyanate phenol-chloroform extraction method described previously (7, 11). The total RNA concentration was quantified spectrophotometrically. Polyadenylated RNA was isolated from the total RNA with the Oligotex kit (QIAGEN, Valencia, Calif.).

**SSH.** The SSH procedure was performed with a PCR-Select cDNA subtraction kit (Clontech, Palo Alto, Calif.) as directed by the manufacturer, except a twofold-greater amount of "driver" cDNA was added to the first and second hybridizations. Starting material consisted of 2  $\mu g$  of an mRNA pool comprised of 25% of each mRNA preparation from the 20-, 40-, 60-, and 120-min  $Mn^{2+}$ -induced (filamentous morphology) cultures. The second mRNA pool was comprised of 25% of each mRNA preparation from the 20-, 40-, 60-, and 120-min non- $Mn^{2+}$ -induced (pellet morphology) cultures. For isolation of cDNAs associated with pellet morphology, the cDNA from non- $Mn^{2+}$ -induced *A. niger* cells was used as the "tester" and the cDNA from the  $Mn^{2+}$ -induced cells was used as the driver. For isolation of cDNAs associated with filamentous morphology, the cDNA from the  $Mn^{2+}$ -induced cells was used as the tester and the cDNA from the non- $Mn^{2+}$ -induced *A. niger* cells was used as the driver. One of two restriction endonucleases, RsaI or AluI, was used to digest the initial cDNA pools employed in the construction of the SSH cDNA libraries described above. Thus, four SSH libraries were prepared: *Arsa* designates pellet-associated genes from the SSH library generated with the RsaI-digested cDNA, *Brsa* designates filament-associated genes from the SSH library generated with the RsaI-digested cDNA, *Aalu* designates pellet-associated genes from the SSH library generated with the AluI-digested cDNA, and *Balu* designates filament-associated genes from the SSH library generated with the AluI-digested cDNA.

**Differential screening of SSH cDNA libraries and sequencing of SSH cDNA fragments from SSH.** The PCR products generated by SSH were cloned into the pGEM-T Easy vector (Promega, Madison, Wis.) to form the SSH cDNA libraries described above. The JM109 *E. coli* colonies containing the pGEM-T Easy vector with the SSH cDNA inserts from the four libraries were selected randomly and cultured overnight for plasmid DNA purification. Plasmid DNAs were purified and digested with the restriction endonuclease EcoRI. Two sets of EcoRI-digested DNA fragments were separated by gel electrophoresis and transferred to two separate nylon membranes. Alternatively, the intact plasmid DNAs were directly arrayed on two separate nylon membranes for differential screening. For differential screening of SSH cDNA clones associated with pelleted or filamentous




FIG. 1. Citric acid production by *A. niger* in the presence of 10 or 1,000 ppb of manganese. *A. niger* was cultured under citric acid-producing conditions with 10 and 1,000 ppb of  $Mn^{2+}$ . The conidia ( $10^6$  conidia/ml) were inoculated into 50 ml of CAP medium in 250-ml baffled flasks and incubated at 30°C and shaken at 250 rpm. The samples were harvested at different time points after 2 days of growth. Data are means of determinations from at least three independent fermentations.

tous morphology, the forward SSH cDNA radioactive probes were synthesized by randomly priming the same SSH cDNA used to construct the SSH cDNA library associated with pelleted or filamentous morphology by using [ $\alpha$ -<sup>32</sup>P]dCTP and the Klenow fragment of DNA polymerase I (Rediprime II DNA labeling system; Amersham Biosciences, Piscataway, N.J.) and the reverse SSH cDNA radioactive probes from the same SSH cDNAs used for construction of the SSH cDNA library associated with filamentous or pelleted morphology, respectively. DNA sequencing of the SSH cDNA clones of interest identified by differential screening was performed at Iowa State University by using BigDye terminator cycle sequencing kits, followed by analysis on an ABI Prism 377 DNA sequencer. The DNA sequences obtained were compared to the nonredundant NCBI nucleotide and protein databases by using BLAST (1), to the EMBL-EBI FUNGI nucleotide database and the Swiss-Prot database by using FASTA3 (36), and to the fungal (*Aspergillus nidulans*, *Neurospora crassa*, and *Magnaporthe grisea*) genome sequence databases at the Center for Genome Research of the Whitehead Institute via BLAST search.

**RNA blotting analysis.** Fifteen or twenty micrograms of total RNA was used for RNA blotting analysis as described by Dai et al. (11) with Zeta-probe blotting membranes (Bio-Rad, Richmond, Calif.). Hybridizations were performed at 65°C (8) to radioactive probes synthesized by random priming of an EcoRI fragment containing the SSH cDNA fragment with [ $\alpha$ -<sup>32</sup>P]dCTP and the Klenow fragment of DNA polymerase I (Amersham Biosciences). The blots were exposed to X-ray film at -30°C with intensifying screens. The film was developed and scanned by using an Epson Expression 800 scanner with a transparency unit (Epson America, Inc., Long Beach, Calif.), and the relative expression levels of mRNAs were quantified by using GelExpert software (NucleoTech, San Carlos, Calif.). The amount of 18S rRNA in each sample was also determined; as an internal control, the blots were stripped according to the manufacturer's instructions (Bio-Rad) and hybridized with a radiolabeled 18S rRNA probe. The relative abundance of each gene transcript was normalized to the amount of 18S rRNA at each time point and expressed as a percentage.

**Isolation of the full-length *Brsa-25* cDNA with RACE-PCR and the full-length gene by PCR.** The full-length *Brsa-25* clone was isolated by 5' and 3' rapid amplification of cDNA ends (RACE)-PCR. Poly(A)<sup>+</sup> RNA (1  $\mu$ g) pooled from different treatments described in the "SSH" section was used to synthesize anchored, double-stranded cDNA templates for amplification of the 5' and 3' ends of cDNA clones via a Marathon cDNA amplification kit (Clontech). The oligonucleotide FP-8 (5'-GGGTGGTGGAGATATCTGGGAAGT-3') and adapter primer (API) provided by the manufacturer were used for the isolation of the 5'-end fragment, while FP-7 (5'-GCAGTATATTCACACGCCAAGTCTCATC-3') and API were used for the isolation of the 3'-end fragment via RACE-PCR. The amplified fragments were cloned into pGEM-T Easy vectors (Promega), and three or four independent plasmids were sequenced. The sequences of the 5'- and 3'-end cDNA fragments were aligned with the sequence of the *Brsa-25* SSH cDNA fragment to verify that the newly isolated fragments belonged to the proper gene. The full-length cDNA clone of *Brsa-25* was amplified by PCR with the primer pair FP-37 (5'-CCTCTATTCTGTCTCCCTTC

GGCGAT-3') and FP-38 (5'-GACACCATCACAGACATATACAGAGA-3'). The fragment was cloned into the pGEM-T Easy vector to form pZD557 and was sequenced. Genomic DNA fragments for *Brsa-25* were isolated with the oligonucleotide pair FP-81 (5'-GGTTTCTTTATCCTGTCCGTATGCTG-3') and FP-82 (5'-CTGTGGAGTAGATGGGCACCTTGTAT-3'). The genomic fragments were cloned into the pGEM-T Easy vector and sequenced.

**Antisense expression vector and *Agrobacterium*-mediated transformation of *A. niger*.** The *Brsa-25* antisense expression vector was constructed by amplifying a 1,900-bp fragment containing the whole coding region and a portion of the terminator by using plasmid DNA (pZD557) as the template, high-fidelity DNA polymerase, and a primer pair designed to introduce BamHI and HpaI sites (in bold) at the 5' and 3' ends, respectively (FP-66, 5'-CAGGATCCCCTTATTC TGTCTCCCTTCGGCGAT-3'; FP-67, 5'-GGGTTAACGACACCATCACAGACATATACAGAGA-3'). The PCR fragment was first cloned into the PCR-Blunt II-TOPO vector (Invitrogen, Carlsbad, Calif.) to form pZD570. Second, the full-length cDNA fragment was excised with the restriction endonucleases BamHI and HpaI and ligated to the appropriately digested vector pZD567 (modified from pAN8-1 [39]) to form pZD574, in which the *Brsa-25* cDNA fragment in antisense orientation was under the control of the *gpdA* promoter and the *tpc* transcriptional terminator. Third, the fragment containing the *gpdA* promoter, the *Brsa-25* cDNA fragment in antisense orientation, and the *tpc* transcription terminator was excised with restriction endonucleases BglII and NdeI, treated with Klenow enzyme, and ligated to the SmaI fragment of pZD581 to form binary vector pZD586. The resulting construct was introduced into *Agrobacterium tumefaciens* strain AGL0 and transferred into *A. niger* cells based on the methods described by Piers et al. (38) and de Groot et al. (12).




FIG. 2. Effects of  $Mn^{2+}$  on *A. niger* morphological formation. The conidia ( $10^6$  conidia/ml) were inoculated into 2 ml of CAP medium supplemented with different amounts of  $Mn^{2+}$  in 16- by 125-mm silanized glass tubes that were positioned at about 15° from horizontal on the shaker platform. The cultures were incubated at 30°C with shaking at 250 rpm. The mycelia from each culture were observed microscopically after 3 days of growth to assess the effects of  $Mn^{2+}$  on development. All photos were taken at the same magnification ( $\times 75$ ).




FIG. 3. Effects of  $Mn^{2+}$  on citric acid production in *A. niger* cultures. *A. niger* conidia ( $10^6$  conidia/ml) were inoculated into 2.0 ml of CAP medium containing different concentrations of  $Mn^{2+}$  in 16- by 125-mm silanized glass tubes that were positioned about  $15^\circ$  from horizontal against the shaker platform. The cultures were incubated at  $30^\circ C$  and shaken at 250 rpm. Citric acid was measured in the culture medium after 3 days of growth.

## RESULTS AND DISCUSSION

**Detailed examination of the effects of  $Mn^{2+}$  on citric acid production and morphology of *A. niger*.** The effects of  $Mn^{2+}$  concentration on citric acid synthesis, uptake, and export have been examined in different *A. niger* strains (2, 32, 41, 45). However, a detailed analysis of the effects of  $Mn^{2+}$  on morphology formation and its linkage to citric acid production in *A. niger* cultures has not been performed. The effect of  $Mn^{2+}$  concentration on citric acid production by *A. niger* versus time was examined in baffled-flask cultures with 10 or 1,000 ppb of  $Mn^{2+}$ . Citric acid production dramatically increased after 48 h of growth in 10-ppb  $Mn^{2+}$  cultures, while it remained very low in 1,000-ppb  $Mn^{2+}$  cultures (Fig. 1). Cultures of *A. niger* in CAP medium with 10 ppb of  $Mn^{2+}$  produced about 35 g of citric acid/liter after 5 days of growth, exhibiting a sixfold increase from day two to day three. Similar citric acid accumulation patterns have been observed in other *A. niger* strains (43).

We further examined the effects of  $Mn^{2+}$  levels on the morphology formation and citric acid production of *A. niger* cells grown in glass culture tubes. Cultures with  $Mn^{2+}$  concentrations from 10 (background level) to 13 ppb had restricted hyphal growth (pellet morphology was maintained), but an increase of just 1 to 2 ppb in  $Mn^{2+}$  concentration dramatically enhanced the hyphal growth (Fig. 2). Citric acid concentrations in 3-day *A. niger* cultures were similar for  $Mn^{2+}$  concentrations from 10 to 13 ppb but decreased more than 70% in the 14- and 15-ppb  $Mn^{2+}$  cultures (Fig. 3). The effects of  $Mn^{2+}$  concentration on citric acid production have been reported previously (21, 44, 45), but in the present study, the effects of  $Mn^{2+}$  on *A. niger* morphology and citric acid production were examined concurrently. Low  $Mn^{2+}$  concentrations (10 to 13 ppb) also significantly suppressed the biomass accumulation (data not shown). The results provide the information on the proper culture conditions necessary for the isolation and characterization of genes that are responsive to  $Mn^{2+}$  concentration and likely to be involved in the control of morphology.

**Isolation of cDNA clones for  $Mn^{2+}$ -responsive morphology control genes in *A. niger*.** Previous observations (9, 21, 41) and the results reported here demonstrate that optimal citric acid production in *A. niger* cultures is associated with pelleted morphology. Fungal morphology formation is regulated by different factors, such as dissolved  $O_2$  concentration, agitation, substrate concentration, and the minerals in the culture media (10, 21, 55). In order to synchronize *A. niger* growth and minimize temporal variations in the mRNA pool, 12 shake flask cultures were grown under pelleted growth conditions (10 ppb of  $Mn^{2+}$ ) for the first 12 h. Then, 1,000 ppb of  $Mn^{2+}$  was added to six of the cultures to induce hyphal growth. The addition of 1,000 ppb of  $Mn^{2+}$  induced hyphal growth in most of the cells after just 40 min and in all of the cells by 120 min (Fig. 4).

Based on the conspicuous effect of  $Mn^{2+}$  concentration on




FIG. 4. Early induction of filamentous mycelia growth. Conidia ( $10^6$  conidia/ml) were inoculated into 250 ml of CAP medium in 1,000-ml baffled flasks and incubated at  $30^\circ C$  at 250 rpm. *A. niger* was precultured under citric acid production conditions for 12 h. Thereafter, 1,000 ppb of  $Mn^{2+}$  was added to half of the *A. niger* cultures for induction of filamentous growth while the other half of the *A. niger* cultures were maintained in CAP medium (10 ppb of  $Mn^{2+}$ ) for pelleted growth. The mycelia were harvested at different induction intervals for microscopic observation and RNA extraction. The photos in the left panels were taken from mycelia harvested from a citric acid-producing culture (pellet growth), and the ones in the right panels were taken from mycelia harvested from  $Mn^{2+}$ -inducing filamentous growth. All photos were taken at the same magnification ( $\times 75$ ). The labels to the right of the panels represent the length of induction.

TABLE 1. mRNA size and DNA sequence analysis of the SSH cDNAs differentially expressed in pellet or filamentous morphology cells

Clone type and name	Putative gene identification <sup>a</sup>	mRNA length (kb) <sup>b</sup>	cDNA obtained (bp) <sup>c</sup>
Filamentous morphology-associated			
<i>Balu-4</i>	<i>A. nidulans</i> G-protein $\beta$ -subunit ( <i>sfaD</i> ) gene (69.9%)	2.0	293
<i>Balu-42</i>	Unknown	1.2	606
<i>Balu-52</i>	Unknown	2.2	191
<i>Brsa-25</i>	Unknown	1.9	413
<i>Brsa-35</i>	Unknown	1.5	372
<i>Brsa-43</i>	<i>A. oryzae</i> peptidase ( <i>tppA</i> ) gene (47%)	2.1	348
<i>Brsa-47</i>	<i>N. crassa</i> inositol-1-phosphate synthase ( <i>MIPS</i> ) gene (76.7%)	1.8	153
<i>Brsa-48</i>	Unknown	1.0	350
<i>Brsa-61</i>	<i>A. nidulans</i> cobalamin-independent methionine synthase ( <i>metH/D</i> ) gene (89%)	2.2	235
<i>Brsa-62</i>	<i>A. nidulans</i> "pyridoxine synthesis" ( <i>pyroA</i> ) gene (60%)	2.1	525
<i>Brsa-64</i>	Unknown	1.0	198
<i>Brsa-109</i>	Unknown	1.9	272
<i>Brsa-112</i>	<i>N. crassa</i> ATP citrate lyase ( <i>acl1</i> ) (85%)	2.0	130
<i>Brsa-116</i>	<i>P. chrysogenum</i> homocitrate synthase ( <i>lys1</i> ) gene (80.6% in the first 200 bp)	1.5	515
<i>Brsa-118</i>	<i>S. cerevisiae</i> hydroxymethylglutaryl-CoA synthase ( <i>erg13</i> ) gene (55%)	2.0	299
Pellet morphology-associated			
<i>Arsa-7</i>	Unknown	1.5	619
<i>Arsa-10</i>	<i>Talaromyces emersonii</i> pepsin-type protease (54%)	1.7	404
<i>Arsa-27</i>	Unknown	1.2	550
<i>Aalu-37</i>	Unknown	0.6	305
<i>Arsa-43</i>	<i>Arthroderma benhamiae</i> polyubiquitin ( <i>ubi</i> ) gene (100%)	1.1	390
<i>Arsa-48</i>	Unknown	0.8	615
<i>Aalu-90</i>	Unknown	1.5	338

<sup>a</sup> The percentages of nucleotide sequence identity shown in parentheses were obtained by using BLAST to search the NCBI nonredundant database.

<sup>b</sup> The length of mRNAs was estimated based on the RNA ladder size.

<sup>c</sup> The length of SSH cDNA clones was determined by DNA sequencing with an automated ABI Prism 377 DNA sequencing system.

morphology (Fig. 2 and 4), we hypothesized that a set of genes may be involved in  $Mn^{2+}$ -inducible morphology changes in *A. niger* cultures. In order to isolate the few critical genes from the estimated 14,000 genes found in *A. niger* (G. Groot, H. Pel, N. van Peij, and A. van Ooyen, Abstr. Annu. Meet. Soc. Ind. Microbiol., 2002), the highly selective SSH method was employed. The SSH and Southern differential screening analysis suggested differential expression of 18 mRNAs in *A. niger* cells induced by 1,000 ppb of  $Mn^{2+}$  and of nine mRNAs in *A. niger* cells with the initial (10-ppb  $Mn^{2+}$ ) culture medium. Differential expression of 15 of the 18 genes represented by the cloned fragments for the high- $Mn^{2+}$  (1,000-ppb) cultures and seven of nine genes represented by the cloned fragments for the low- $Mn^{2+}$  (10-ppb) cultures were confirmed by Northern analysis (data not shown). These 15  $Mn^{2+}$ -enhanced clones had relatively low expression levels when *A. niger* cells were maintained at low- $Mn^{2+}$  conditions, while their expression was significantly enhanced upon the addition of 1,000 ppb of  $Mn^{2+}$ . In contrast, the seven  $Mn^{2+}$ -suppressed clones exhibited very high expression levels in low- $Mn^{2+}$  cultures and drastically decreased expression after a 40-min exposure to high  $Mn^{2+}$  concentrations. The full-length mRNAs for the 15  $Mn^{2+}$ -enhanced clones were between 1 and 2.2 kb and were between 0.6 and 1.7 kb for the seven  $Mn^{2+}$ -suppressed clones (Table 1).

**Sequence analysis of SSH cDNA clones.** The nucleotide sequences of the partial cDNA clones (ranging from 130 to 619 bp) were determined to gain insight into the functions of the encoded proteins. BLAST and FASTA3 analyses of the nucleotide and translated sequences of the cDNA clones against GenBank, the EMBL-EBI FUNGI nucleotide database, and the genome databases of *A. nidulans*, *N. crassa*, and *M. grisea* revealed that seven of the  $Mn^{2+}$ -enhanced (filament-associ-

ated) clones did not have significant similarity to known sequences. Eight of the clones possessed various degrees of identity to known genes from other organisms (Table 1). The gene products with homology to known proteins fell into one of two groups, those involved in signal transduction or those involved in amino acid synthesis or protein catabolism.

There are two genes associated with the signaling group. The translated sequence of *Balu-4* is 96% identical to the *A. nidulans* G-protein  $\beta$ -subunit gene, *sfaD*. This heterotrimeric G-protein component is known to be required for normal growth and repression of sporulation in *A. nidulans* (42, 58). The translated sequence of *Brsa-47* is 60% identical to the *Pichia pastoris* inositol-1-phosphate synthase (*ino1*) gene. This is the first enzyme on the biosynthetic pathway to inositol phosphates involved in intracellular signaling. For example, inositol-1,4,5-trisphosphate induces  $Ca^{2+}$  release, which stimulates hyphal tip growth in *N. crassa* (48).

The remaining six genes with putative functions fall into the amino acid synthesis or protein utilization group. The deduced amino acid sequence of *Brsa-43* shares 59% identity with the tripeptidyl peptidase A (*tppA*) of *Aspergillus oryzae*. Tripeptidyl peptidases have been well studied in mammalian systems, where they release N-terminal tripeptides from oligopeptides generated by different endopeptidases. The tripeptides are further degraded by other exopeptidases to release amino acids and dipeptides (52). Expression of *tppA* in *Streptomyces lividans* was enhanced during filamentous growth and suppressed during pelleted growth (59). The translated protein sequence of *Brsa-62* shares 86% identity with the *A. nidulans* *pyroA* gene product, an enzyme involved in pyridoxine biosynthesis, which is important for amino acid metabolism. The deduced amino acid sequence of *Brsa-112* shares 85% identity with that of the

*N. crassa acI1* gene product, ATP citrate lyase, which provides cytosolic acetyl coenzyme A (acetyl-CoA) for lipid synthesis and is crucial for the accumulation of substantial amounts of lipid in fungi (56). In a fungus from the same family as *N. crassa*, *Sordaria macrospora*, the ATP citrate lyase was found to be specifically induced at the beginning of the sexual cycle, thus producing the acetyl-CoA required for biosynthesis during fruiting body formation at later stages of sexual development (33). Clearly the expression of ATP citrate lyase during filamentous growth would lead to a net decrease in the production of citric acid. However, the relative contribution of this enzyme to decreasing citrate accumulation has not been quantified. The increased levels of the *acI1* transcript are also consistent with increased amino acid biosynthesis, as the oxaloacetate produced can be transaminated to L-aspartate and subsequently to other amino acids of the aspartate group. The clone *Brsa-116* shares 88% identity with the *P. chrysogenum lys1* gene encoding homocitrate synthase, the first step in lysine biosynthesis in fungi. *Brsa-118* shares 54% identity with the *Saccharomyces cerevisiae erg13* gene encoding hydroxymethylglutaryl-CoA synthase. This enzyme produces the first metabolite on the pathway to branched-chain amino acid synthesis, as well as sterol synthesis (though the latter is controlled at the level of hydroxymethylglutaryl-CoA reductase). The genes *acI1*, *erg13*, *ino1*, *lys1*, *pyroA*, and *tpaA* were found to be involved in cell growth and tissue development (3, 20, 35, 51). The deduced peptide sequence of the *Brsa-61* product is identical to that of the *A. nidulans metH/D* gene product. This gene encodes the cobalamin-independent methionine synthase, which is the enzyme responsible for methionine synthesis in eukaryotic organisms.

BLAST analysis of the Mn<sup>2+</sup>-suppressed clones (pellet associated) showed that five of seven clones had no significant homology to known sequences in GenBank and other databases. The deduced amino acid sequence of *Arsa-10* is 59% identical to *A. oryzae* aspergillopepsin O (*pepO*), an aspartic proteinase. The Northern blot analysis of *Arsa-10* showed high expression during early pelleted growth and suppression when *A. niger* switched to filamentous growth (see Fig. 6). Reichard et al. examined the aspergillopepsin PEP with immunofluorescence and found that it was mainly located in developing conidiophores of aspergilli, in submerged mycelia, and on the tips of growing aerial mycelia, whereas mature aerial hyphae and spores showed no immunofluorescence (40). The results suggest a role for such enzymes in the growth of hyphae and the development of conidiophores, and thus for the sporulation process in aspergilli.

Another gene, *Arsa-43*, is identical to the translated ubiquitin (*ubi*) gene of *Arthroderma benhamiae* (Table 1) (18). Ubiquitin is attached to other proteins via an isopeptide linkage formed by multiple enzymatic steps to form a polyubiquitinated protein. The attachment of ubiquitin through K48 or K29 marks the modified proteins for proteolysis by the 26S proteasome. Proteins modified by this process are generally tightly regulated proteins involved in the control of cellular processes, for example, meiosis in fission yeast (34). The attachment of ubiquitin through K63 occurs for proteins involved in other critical cellular processes, such as stress response in *S. cerevisiae* (37).




FIG. 5. Induction of *Balu-4*, *Brsa-25*, *Brsa-43*, *Brsa-47*, *Brsa-109*, and *Brsa-118* mRNA by 1,000 ppb of Mn<sup>2+</sup>. Twenty micrograms of total RNA used in SSH was subjected to denaturing gel electrophoresis and hybridized with radioactively labeled probes prepared from cDNA clone fragments. Autoradiographs of the RNA blotting are shown on the left. Relative RNA levels are plotted on the right. The percentage of the relative amount of mRNA estimated by gel blot intensities was calculated based on relative levels of 18S rRNA.

**Expression pattern of the Mn<sup>2+</sup>-responsive transcripts during early developmental stages.** In order to investigate the relative levels and temporal expression patterns of mRNA transcripts, Northern blot analyses were performed for a selection of 12 of the 22 genes potentially involved in Mn<sup>2+</sup>-responsive morphology switching. The 20-, 40-, and 120-min RNA pools for the Northern analyses were the same as those used for the SSH library construction. All of the filament-associated genes, except the *Brsa-43* and *Brsa-109* genes, had one transcript (Fig. 5). The levels of the transcripts for each gene were determined at different time points by densitometry of the Northern blots. The relative transcription levels for all

six genes at different time points both before and after addition of  $Mn^{2+}$  (induction of filamentous growth) were estimated based on the 18S rRNA amounts at each time point (Fig. 5, right panel). In the absence of  $Mn^{2+}$  induction (pelleted growth condition), the transcription of the six filament-associated genes remained relatively low. *Brsa-43* and *Brsa-47* transcription gradually increased throughout the time course of  $Mn^{2+}$  induction, while *Brsa-25* transcription reached a maximum within 40 min and maintained that level. The transcription of *Balu-4*, *Brsa-109*, and *Brsa-118* was transient, attaining peak expression at 40 min and declining thereafter (the transcript levels at 40 min were 326, 233, and 192% of the 18S rRNA transcript, respectively).

Similarly, RNA blotting analysis was used to examine the expression patterns of the pellet-associated ( $Mn^{2+}$ -suppressed) genes in cultures harvested 20, 40, and 120 min after the addition of 1,000 ppb of  $Mn^{2+}$  or without additional  $Mn^{2+}$  (Fig. 6). The relative transcript levels for the three time points with or without  $Mn^{2+}$  induction were estimated based on the amount of 18S rRNA for each time point (Fig. 6, right panel). The relative transcript levels of clones *Arsa-7*, *Aalu-37*, *Arsa-43*, and *Aalu-90* under pelleted growth conditions were at least 300% of the 18S rRNA transcript. The transcripts associated with these four clones decreased significantly 40 min after 1,000 ppb of  $Mn^{2+}$  was added to the 12-h culture. After 120 min, the transcript levels of the four genes (*Arsa-7*, *Aalu-37*, *Arsa-43*, and *Aalu-90*) were only 57, 26, 70, and 22% of the 18S rRNA levels, respectively. In contrast to the four genes above, *Arsa-10* and *Arsa-27* exhibited relatively low expression under pelleted growth conditions (10 ppb); however, their response to filamentous growth conditions (addition of 1,000 ppb of  $Mn^{2+}$ ) was similar to that of the other four highly transcribed, pellet-associated genes, i.e., transcript levels rapidly decreased.

**Expression patterns of the  $Mn^{2+}$ -responsive transcripts during the citrate production process.** Pelleted morphology and citric acid overproduction are associated physiological traits, as are filamentous morphology and a lack of citrate production. The entire developmental time course of growth and citric acid production in *A. niger* is completed in approximately 5 days. To evaluate the potential involvement of various morphology control genes with regard to citric acid production, the expression patterns of selected genes were examined at different growth stages extending to 5 days. Figure 7 shows the mRNA accumulation patterns for six filamentous morphology-associated genes (*Balu-4*, *Balu-42*, *Brsa-25*, *Brsa-47*, *Brsa-109*, and *Brsa-118*) from 1 to 5 days after the addition of 1,000 ppb of  $Mn^{2+}$  (noncitrate production condition [NCP]) (right panel of Fig. 7) and without addition of  $Mn^{2+}$  (citrate production condition [CP]) (left panel of Fig. 7). The transcription of all six genes was suppressed under pelleted growth (CP) conditions during the 5-day time course and dramatically enhanced under filamentous growth (NCP) conditions. The transcript levels of clone *Balu-4* (G-protein  $\beta$ -subunit) were lower than those of the other five selected clones during the NCP time course, while the *Balu-4* transcript was not even detected during the CP time course. The results shown in Fig. 5 and 7 suggest that *Balu-4* is indeed required for filamentous (NCP) growth and that  $Mn^{2+}$  only enhanced a transient expression of the *Balu-4* gene. This is consistent with previous observations for *A. nidulans* and *N. crassa* (42, 58). The suppressive effect of the




FIG. 6. Suppression of *Arsa-7*, *Arsa-10*, *Arsa-27*, *Aalu-37*, *Arsa-43*, and *Aalu-90* mRNA by 1,000 ppb of  $Mn^{2+}$ . Twenty micrograms of the total RNA pools was subjected to denaturing gel electrophoresis and hybridized with radioactively labeled probes prepared from the cloned cDNA fragments. Autoradiographs of the Northern blots are shown on the left. Relative RNA levels are plotted on the right. The percentages of relative amounts of mRNA were estimated by densitometry of bands and normalized to the relative amount of 18S rRNA.

G-protein  $\beta$ -subunit on vegetative growth of *Cryphonectria parasitica* was also observed on synthetic medium (19). This suggests that the G-protein  $\beta$ -subunit has dynamic effects on fungal growth and development. Clones *Balu-42* and *Brsa-109* maintained relatively high steady-state transcription levels over 4.5 days of NCP growth (Fig. 7). The transcription levels of these two genes (on the basis of rRNA levels) were at least four times greater than those of *Balu-4* and *Brsa-118* and at least two times greater than those of *Brsa-25* and *Brsa-47* during NCP growth (data not shown). Low transcription was observed for clones *Balu-42* and *Brsa-47* during the 5 days of pelleted (CP) growth. The *Brsa-25*, *Brsa-109*, and *Brsa-118* genes, like *Balu-4*, were specifically expressed under NCP




FIG. 7. RNA gel blot analysis of *Balu-4*, *Balu-42*, *Brsa-25*, *Brsa-47*, *Brsa-109*, and *Brsa-118* during citrate production (10 ppb; pelleted) and noncitrate production (1,000 ppb; filamentous) growth. The blots contained 15  $\mu$ g of total RNA prepared from the biomass of *A. niger* cells grown for 0.04, 0.08, 1, 1.5, 2, 3, 4, 4.5, or 5 days after an initial growth period of 12 h with 10 ppb of  $Mn^{2+}$ . All membranes were stripped and hybridized with an 18S rRNA probe to verify equivalent sample loading and for the estimation of the relative transcription of those genes.

growth conditions. Interestingly, *Brsa-25* and *Brsa-118* had relatively high transcription levels on the first day of NCP growth but decreased to undetectable levels by 1.5 days (Fig. 7). Transcription of *Brsa-25* and *Brsa-118* increased on day 3 of NCP growth, but thereafter, the transcription of *Brsa-25* increased further while *Brsa-118* decreased slightly. This suggests that both *Brsa-25* and *Brsa-118* are required during the rapid growth stage and also during the later vegetative growth stages that may be associated with certain physiological stresses.

Four pellet-associated clones were also examined during CP and NCP growth conditions. The clones *Arsa-7*, *Aalu-37*, and *Aalu-90* exhibited high levels of transcription during 5 days of pelleted (CP) growth (Fig. 8, left panel). The relative mRNA levels of these clones (normalized to the 18S rRNA amount)

gradually increased (data not shown). During filamentous (NCP) growth, clones *Aalu-37* and *Aalu-90* had very low transcript levels on the first day of NCP growth, followed by a gradual increase in transcription (Fig. 8). In contrast, the transcription of clone *Arsa-7* was suppressed during the first 3 days of NCP growth and rapidly increased thereafter. The transcription of *Arsa-48* exhibited a third pattern, increasing rapidly on the first day of pelleted (CP) growth and then gradually decreasing until day 3, followed by another increase during the later stages of CP growth (days 4 and 5) (Fig. 8). *Arsa-48* was suppressed during the entire 4.5 days of NCP growth. The results indicate that these genes not only respond to  $Mn^{2+}$  but are also responsive to other factors during filamentous growth.

**Isolation of the full-length *Brsa-25* cDNA and its genomic clone.** For the putative morphology control genes, isolation of the full-length genomic clone and untranslated regions would be of interest in the search for regulatory elements. In addition, for cDNA clones of unknown function, perhaps a full-length gene would reveal homologies not detected with the partial cDNA sequence. To this end, the SSH clone *Brsa-25* (unknown function) was selected for further characterization in this study. The 413-bp fragment of *Brsa-25* was used to design two gene-specific oligonucleotide primers. The 5' and 3' ends of the *Brsa-25* gene were isolated by using RACE-PCR. Sequence analysis confirmed the newly isolated 5' and 3' ends of *Brsa-25*. The full-length cDNA has a 1,797-bp open reading




FIG. 8. RNA gel blot analysis of *Arsa-7*, *Aalu-37*, *Arsa-48*, and *Aalu-90* during citrate production (10 ppb; pelleted) and noncitrate production (1,000 ppb; filamentous) growth. The blots contained 15  $\mu$ g of total RNA prepared from the biomass of *A. niger* cells grown for 0.04, 0.08, 1, 1.5, 2, 3, 4, 4.5, or 5 days after an initial growth period of 12 h with 10 ppb of  $Mn^{2+}$ . All membranes were stripped and hybridized with an 18S rRNA probe to verify equivalent sample loading and for the estimation of the relative transcription of those genes.






FIG. 9. Putative protein encoded by *Brsa-25*. (A) The diagram shows the *Brsa-25* gene structure containing six exons (rectangles) and five introns. ATG is the translation start codon and TAG is the translation stop codon. (B) Deduced amino acid sequence of *Brsa-25*. (C) Hydropathy plot of the predicted *Brsa-25* protein. The plot was constructed according to the method of Kyte and Doolittle (24), with a window of 11 amino acid residues.

frame, a 62-bp 5' untranslated region, and a 198-bp 3' untranslated region. The cDNA encodes a 598-amino-acid protein with an apparent molecular mass of 66.3 kDa, as shown in Fig. 9B. The hydropathy profile analysis of this protein, performed with the Kyte-Doolittle algorithm (24), indicated a highly hydrophobic nature and predicted 8 to 11 putative transmembrane domains (Fig. 9C). To determine the genomic structure and organization of *Brsa-25*, a 2.28-kb genomic DNA fragment was amplified by PCR with primers based on the two ends of the *Brsa-25* cDNA and was sequenced. This full-length gene was compared with the *Brsa-25* cDNA sequence. As shown in Fig. 9A, the *Brsa-25* gene consists of six exons and five introns. To ensure maximum likelihood of identifying homologous sequences, both the nonredundant database (GenBank) and focused-coverage databases (*N. crassa*, *A. nidulans*, and *Aspergillus fumigatus* genome sequence databases) were used for BLAST analyses. The amino acid sequence of *Brsa-25* showed 22% identity with the N amino acid transport system protein of *N. crassa*. The result of an NCBI conserved-domain search (RPS-BLAST) indicated that *Brsa-25* contained transmembrane regions that were 98.2% aligned to the conserved domain of the transmembrane amino acid transporter protein.

This suggests that *Brsa-25* may be an amino acid transporter or, at minimum, an integral membrane protein.

**Antisense expression of *Brsa-25* in *A. niger*.** To investigate the potential role of the *Brsa-25* gene in *A. niger* morphology control and citric acid production, an antisense expression vector was constructed (Fig. 10A). A construct containing only the *gpdA* promoter and *trpC* terminator was introduced into *A. niger* as a transgenic control. Ten transgenic controls and 15 antisense *A. niger* transformants were examined in CAP medium containing 15 ppb of  $Mn^{2+}$  (conditions normally leading to filamentous growth). Eleven of the antisense *Brsa-25* transformants of *A. niger* restricted the filamentous growth compared with the transgenic control in 60-h cultures (Fig. 10B). The citric acid concentration in 60-h cultures was determined and showed that citrate production in the antisense transformants increased an average of 30% versus the production of the transgenic control. Figure 11 shows the citric acid production in the 60-h cultures of the control and transgenic clones shown in Fig. 10C. The citric acid production in the cultures of antisense strains *Brsa-25-3* and *Brsa-25-5* was about 35% higher than that of the selected control, while the citric acid production in antisense strain *Brsa-25-8* was only 11% higher




FIG. 10. Effect of antisense expression of *Brsa-25* on *A. niger* morphology formation. (A) Diagram of the plasmid pZD570 containing the *Brsa-25* gene in antisense orientation. The *pGpdA* corresponds to the promoter of the glyceraldehyde-3-phosphate dehydrogenase of *A. nidulans*. *TrpC* is the *A. nidulans* *TrpC* transcription terminator. This plasmid also contains the *hph* gene of *E. coli*, which confers hygromycin resistance. (B) Microscopic observation of the morphology of the transgenic control (TAN-2811) containing the transgene expression vector with only the promoter (*pGpdA*) and terminator (*TrpC*) and the selected *Brsa-25* antisense transgenic line (antisense Brsa-25-3) after 60-h culture at 30°C and 250 rpm. (C) The RNA gel blot analysis of steady-state mRNA levels of *Brsa-25* in antisense suppression strains. Total RNA was isolated from 60-h cultures of wild-type *A. niger* (lane 1), transgenic control strains (lane 2), antisense strain Brsa-25-3, -5, and -8 (lanes 3 to 5). Twenty micrograms of total RNA was loaded on each lane and hybridized with the radioactive labeled probe of the *Brsa-25* SSH cDNA fragment. The same blot was stripped and hybridized with 18S rRNA for equivalent loading.

than that of the control. The citric acid production observed in these selected clones was inversely correlated to the levels of mRNA seen in the Northern blot analysis (Fig. 10C). The differences in citric acid production and mRNA depletion by different antisense strains of *Brsa-25* likely result from positional effects on transcription resulting from the integration

of the antisense cassette into different points in the genome. When the *Brsa-25* antisense transformants of *A. niger* were grown at even higher  $Mn^{2+}$  concentrations (20 ppb), the inhibition of filamentous growth and increase in citrate production became less pronounced (data not shown). The data are consistent with the gene *Brsa-25* having the expected effect on *A. niger* morphology in response to manganese. However, the retention of the sensitivity of morphology development to higher  $Mn^{2+}$  concentrations suggests that morphology control in *A. niger* is effected by multiple genes.

In summary, the differentially expressed genes that have a tentatively identified function can be assigned to two general categories: those involved in amino acid or protein metabolism (or cell growth) and those involved in cell regulation. The genes *erg13*, *lys1*, *metH/D*, *pyroA*, *acl1*, and *tpa4* that are induced by high  $Mn^{2+}$  levels and the gene *pepO* that is suppressed by  $Mn^{2+}$  belong in the amino acid metabolism category. The rapid hyphal growth associated with the switch to filamentous morphology observed upon induction by sufficient  $Mn^{2+}$  levels probably requires increased protein production, as well as degradation and utilization of proteins required for the maintenance of the pelleted growth state. The observed induction of genes involved in amino acid anabolism is consistent with this requirement. The expression patterns of *lys1* and *erg13* exhibited a rapid increase before decreasing, suggesting a transient high demand for de novo protein synthesis. The induced genes, *ino1*, *sfaD*, and *acl1*, and the repressed gene, *ubi*, belong in the category of cell regulation. One of the 13 genes without a known function or one of the tentatively identified cell regulatory genes may be a keystone gene that controls morphology. Alternatively, multiple genes acting in concert may be required for the observed effect on morphology and citric acid production. Encouragingly, the functional analysis of *Brsa-25* indicated that this "unknown" gene was indeed involved in the regulation of morphology formation. Further functional evaluation of the known and unknown genes may




FIG. 11. Suppression of filament-associated gene *Brsa-25* leads to enhanced citric acid production. Citric acid in the supernatants of 60-h test tube cultures was measured biochemically. The control was a transformed strain carrying the *pGpdA* promoter and *TrpC* terminator.

reveal their roles in the associated physiological properties of morphology control and citric acid production in *A. niger*.

#### ACKNOWLEDGMENTS

We thank Thomas W. Wietsma of the William R. Wiley Environmental Molecular Sciences Laboratory at PNNL for performing manganese analyses with an ICP-MS, Diane F. Hu and Katie S. Panther for RNA gel blot image scanning and densitometry, and Scott E. Baker for critical review of the manuscript. DNA sequencing was conducted by the DNA Sequencing and Synthesis Facility of Iowa State University.

Xingxue Mao received a visiting scholarship from Guangdong Academy of Agriculture Science, China. This work was supported by the PNNL Laboratory Directed Research and Development program. Pacific Northwest National Laboratory is operated by the Battelle Memorial Institute for the U.S. Department of Energy under contract DE-AC06-76RL01830.

#### REFERENCES

- Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Banik, A. K. 1976. Mineral nutrition of *Aspergillus niger* for citric acid production. *Folia Microbiol. (Prague)* **21**:139–143.
- Banuelos, O., J. Casqueiro, S. Gutierrez, and J. F. Martin. 2000. Overexpression of the *lys1* gene in *Penicillium chrysogenum*: homocitrate synthase levels, alpha-aminoacidic acid pool and penicillin production. *Appl. Microbiol. Biotechnol.* **54**:69–77.
- Bergmeyer, H. U. 1985. Metabolites 2: tri- and dicarboxylic acids, purines, pyrimidines and derivatives, coenzymes, inorganic compounds, p. 5–10. In *Citric acids*. VCH Publishers, Weinheim, Germany.
- Braun, S., and S. E. Vecht-Lifshitz. 1991. Mycelial morphology and metabolite production. *Trends Biotechnol.* **9**:63–68.
- Byrne, G. S., and O. P. Ward. 1989. Effect of nutrition on pellet formation by *Rhizopus arrhizus*. *Biotechnol. Bioeng.* **33**:912–914.
- Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* **162**:156–159.
- Church, G. M., and W. Gilbert. 1984. Genomic sequencing. *Proc. Natl. Acad. Sci. USA* **81**:1991–1995.
- Clark, D. S. 1962. Submerged citric acid fermentation of ferrocyanide treated beet molasses: morphology of pellets of *Aspergillus niger*. *Can. J. Microbiol.* **8**:133–136.
- Cui, Y. Q., R. C. Van Der Lans, and K. C. Luyben. 1997. Effect of agitation intensities on fungal morphology of submerged fermentation. *Biotechnol. Bioeng.* **55**:715–726.
- Dai, Z., B. S. Hooker, D. B. Anderson, and S. R. Thomas. 2000. Expression of *Acidothermus cellulolyticus* endoglucanase E1 in transgenic tobacco: biochemical characteristics and physiological effects. *Transgenic Res.* **9**:43–54.
- de Groot, M. J., P. Bundock, P. J. Hooykaas, and A. G. Beijersbergen. 1998. *Agrobacterium tumefaciens*-mediated transformation of filamentous fungi. *Nat. Biotechnol.* **16**:839–842.
- Diatchenko, L., Y. F. Lau, A. P. Campbell, A. Chenchik, F. Moqadam, B. Huang, S. Lukyanov, K. Lukyanov, N. Gurskaya, E. D. Sverdlov, and P. D. Siebert. 1996. Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proc. Natl. Acad. Sci. USA* **93**:6025–6030.
- Gurskaya, N. G., L. Diatchenko, A. Chenchik, P. D. Siebert, G. L. Khaspekov, K. A. Lukyanov, L. L. Vagner, O. D. Ermolaeva, S. A. Lukyanov, and E. D. Sverdlov. 1996. Equalizing cDNA subtraction based on selective suppression of polymerase chain reaction: cloning of Jurkat cell transcripts induced by phytohemagglutinin and phorbol-12-myristate-13-acetate. *Anal. Biochem.* **240**:90–97.
- Habison, A., C. P. Kubicek, and M. Roehr. 1983. Partial purification and regulatory properties of phosphofructokinase from *Aspergillus niger*. *Biochem. J.* **209**:669–676.
- Habison, A., C. P. Kubicek, and M. Rohr. 1979. Phospho-fructokinase as a regulatory enzyme in citric-acid producing *Aspergillus niger*. *FEMS Microbiol. Lett.* **5**:39–42.
- Hosobuchi, M., F. Fukui, H. Matsukawa, T. Suzuki, and H. Yoshikawa. 1993. Morphology control of preculture during production of ML-236B, a precursor of pravastatin sodium, by *Penicillium citrinum*. *J. Ferment. Bioeng.* **76**:476–481.
- Kano, R., Y. Nakamura, S. Watanabe, and A. Hasegawa. 1999. Repeated ubiquitin genes in *Trichophyton mentagrophytes*. *Curr. Microbiol.* **39**:302–305.
- Kasahara, S., and D. L. Nuss. 1997. Targeted disruption of a fungal G-protein beta subunit gene results in increased vegetative growth but reduced virulence. *Mol. Plant-Microbe Interact.* **10**:984–993.
- Katayama, S., N. Adachi, K. Takao, T. Nakagawa, H. Matsuda, and M. Kawamukai. 1995. Molecular cloning and sequencing of the *hcs* gene, which encodes 3-hydroxy-3-methylglutaryl coenzyme A synthase of *Schizosaccharomyces pombe*. *Yeast* **11**:1533–1537.
- Kisser, M., C. P. Kubicek, and M. Roehr. 1980. Influence of manganese on morphology and cell wall composition of *Aspergillus niger* during citric acid fermentation. *Arch. Microbiol.* **128**:26–33.
- Kossen, N. W. F. 2000. The morphology of filamentous fungi. *Adv. Biochem. Eng. Biotechnol.* **70**:1–33.
- Kubicek, C. P., W. Hampel, and M. Roehr. 1979. Manganese deficiency leads to elevated amino-acid pools in citric acid accumulating *Aspergillus niger*. *Arch. Microbiol.* **123**:73–80.
- Kyte, J., and R. F. Doolittle. 1982. A simple method for displaying the hydrophobic character of a protein. *J. Mol. Biol.* **157**:105–132.
- Lazo, G. R., P. A. Stein, and R. A. Ludwig. 1991. A DNA transformation-competent *Arabidopsis* genomic library in *Agrobacterium*. *Bio/Technology* **9**:963–967.
- Liang, P., and A. B. Pardee. 1992. Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* **257**:967–971.
- Ma, H., C. P. Kubicek, and M. Roehr. 1985. Metabolic effects of manganese deficiency in *Aspergillus niger*: evidence for increased protein degradation. *Arch. Microbiol.* **141**:266–268.
- Magnuson, J. K., and L. L. Lasure. Organic acid production by filamentous fungi. In J. Tkacz (ed.), *Advances in fungal bio/technology for industry, agriculture and medicine*, in press. Kluwer Academic Publishers, Norwell, Mass.
- Metz, B., and N. W. F. Kossen. 1977. The growth of molds in the form of pellets: a literature review. *Biotechnol. Bioeng.* **19**:781–800.
- Metz, B., N. W. F. Kossen, and J. C. van Suizdum. 1979. The rheology of mold suspensions. *Adv. Biochem. Eng.* **11**:103.
- Michel, F. C., Jr., E. A. Grulke, and C. A. Reddy. 1992. Determination of the respiration kinetics for mycelial pellets of *Phanerochaete chrysosporium*. *Appl. Environ. Microbiol.* **58**:1740–1745.
- Netik, A., N. V. Torres, J.-M. Riol, and C. P. Kubicek. 1997. Uptake and export of citric acid by *Aspergillus niger* is reciprocally regulated by manganese ions. *Biochim. Biophys. Acta* **1326**:287–294.
- Nowrousian, M., S. Masloff, S. Poggeler, and U. Kuck. 1999. Cell differentiation during sexual development of the fungus *Sordaria macrospora* requires ATP citrate lyase activity. *Mol. Cell. Biol.* **19**:450–460.
- Okazaki, K., H. Okayama, and O. Niwa. 2000. The polyubiquitin gene is essential for meiosis in fission yeast. *Exp. Cell Res.* **254**:143–152.
- Osmani, A. H., G. S. May, and S. A. Osmani. 1999. The extremely conserved *pyrA* gene of *Aspergillus nidulans* is required for pyridoxine synthesis and is required indirectly for resistance to photosensitizers. *J. Biol. Chem.* **274**:23565–23569.
- Pearson, W. R., and D. J. Lipman. 1988. Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* **85**:2444–2448.
- Pickart, C. M. 2000. Ubiquitin in chains. *Trends Biochem. Sci.* **25**:544–548.
- Piers, K. L., J. D. Heath, X. Liang, K. M. Stephens, and E. W. Nester. 1996. *Agrobacterium tumefaciens*-mediated transformation of yeast. *Proc. Natl. Acad. Sci. USA* **93**:1613–1618.
- Punt, P. J., and C. A. van den Hondel. 1992. Transformation of filamentous fungi based on hygromycin B and phleomycin resistance markers. *Methods Enzymol.* **216**:447–457.
- Reichard, U., M. Monod, and R. Ruchel. 1996. Expression pattern of aspartic proteinase antigens in *aspergilli*. *Mycoses* **39**:99–101.
- Röhr, M., and C. P. Kubicek. 1981. Regulatory aspects of citric acid fermentation by *Aspergillus niger*. *Process Biochem.* **16**:34–37.
- Rosen, S., J. H. Yu, and T. H. Adams. 1999. The *Aspergillus nidulans* *sfad* gene encodes a G protein beta subunit that is required for normal growth and repression of sporulation. *EMBO J.* **18**:5592–5600.
- Ruijter, G. J. G., H. Panneman, D.-B. Xu, and J. Visser. 2000. Properties of *Aspergillus niger* citrate synthase and effects of *citA* overexpression on citric acid production. *FEMS Microbiol. Lett.* **184**:35–40.
- Sanchez-Marroquin, A., R. Carreno, and M. Ledezma. 1970. Effect of trace elements on citric acid fermentation by *Aspergillus niger*. *Appl. Microbiol.* **20**:888–892.
- Schrefler, G., C. P. Kubicek, and M. Roehr. 1986. Inhibition of citric acid accumulation by manganese ions in *Aspergillus niger* mutants with reduced citrate control of phosphofructokinase. *J. Bacteriol.* **165**:1019–1022.
- Schweiger, L. B. 1961. Production of citric acid by fermentation. Miles Laboratories, Inc., Elkhart, Ind.
- Serizawa, N., M. Hosobuchi, and H. Yoshikawa. 1997. Biochemical and fermentation technological approaches to production of pravastatin, a HMG-CoA reductase inhibitor, p. 779–804. In *Drugs and pharmaceutical sciences*. Marcel Dekker, New York, N.Y.
- Silverman-Gavrila, L. B., and R. R. Lew. 2002. An IP(3)-activated Ca(2+) channel regulates fungal tip growth. *J. Cell Sci.* **115**:5013–5025.
- Smith, J. J., M. D. Lilly, and R. I. Fox. 1990. The effect of agitation on the morphology and penicillin production of *Penicillium chrysogenum*. *Biotechnol. Bioeng.* **35**:1011–1023.
- St. John, T. P., and R. W. Davis. 1979. Isolation of galactose-inducible DNA

- sequences from *Saccharomyces cerevisiae* by differential plaque filter hybridization. *Cell* **16**:443–452.
51. Sugiyama, M., and J. Nikawa. 2001. The *Saccharomyces cerevisiae* Isw2p-Itc1p complex represses INO1 expression and maintains cell morphology. *J. Bacteriol.* **183**:4985–4993.
  52. Tomkinson, B. 1999. Tripeptidyl peptidases: enzymes that count. *Trends Biochem. Sci.* **24**:355–359.
  53. Tucker, K. G., and C. R. Thomas. 1992. Mycelial morphology: the effect of spore inoculum level. *Biotechnol. Lett.* **14**:1071–1074.
  54. Von Stein, O. D., W. G. Thies, and M. Hofmann. 1997. A high throughput screening for rarely transcribed differentially expressed genes. *Nucleic Acids Res.* **25**:2598–2602.
  55. Wongwicharn, A., B. McNeil, and L. M. Harvey. 1999. Effect of oxygen enrichment on morphology, growth, and heterologous protein production in chemostat cultures of *Aspergillus niger* B1-D. *Biotechnol. Bioeng.* **65**:416–424.
  56. Wynn, J. P., A. bin Abdul Hamid, and C. Ratledge. 1999. The role of malic enzyme in the regulation of lipid accumulation in filamentous fungi. *Microbiology* **145**:1911–1917.
  57. Xu, J., L. Wang, D. Ridgway, T. Gu, and M. Moo-Young. 2000. Increased heterologous protein production in *Aspergillus niger* fermentation through extracellular proteases inhibition by pelleted growth. *Biotechnol. Prog.* **16**:222–227.
  58. Yang, Q., S. I. Poole, and K. A. Borkovich. 2002. A G-protein beta subunit required for sexual and vegetative development and maintenance of normal G alpha protein levels in *Neurospora crassa*. *Eukaryot. Cell* **1**:378–390.
  59. Yun, S. I., A. R. Yahya, M. Malten, D. Cossar, W. A. Anderson, J. M. Scharer, and M. Moo-Young. 2001. Peptidases affecting recombinant protein production by *Streptomyces lividans*. *Can. J. Microbiol.* **47**:1137–1140.