Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|------------------------------|-----------------------------|-----------------------------|---------------------------|---------------------------------|-----------------------------|-----------------------------|---|
| The birthwish and months and | | | Ra | ace | I | Hi | spanic Origin | 8 40,797 3 201,391 8 22,202 7 13.6 9 70.9 8 7.8 9 5 77 4 8 6 20.9 8 55.4 6 * 2 15.0 |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | |
| | | | | | Percent | | | |
| Total of all cities of 100,000 or more | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 101,130 904,790 60,921 | 53,250 633,144 34,175 | 41,787 207,831 22,928 | 543 4,802 588 | 5,550 59,013 3,230 | 23,202 251,210 22,355 | 30,258 382,793 12,118 | 40,797 201,391 22,202 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.4 78.1 5.3 | 6.6 80.7 4.4 | 13.5 70.9 7.8 | 7.5 68.9 8.4 | 7.2 79.4 4.3 | 6.4 72.5 6.5 | 6.7 86.9 2.8 | 13.6 70.9 7.8 |
| Abilene, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 156 1,301 111 | 123 1,188 100 | 29 81 9 | - 1 1 | 4 31 1 | 34 256 47 | 89 935 54 | 29 77 8 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.7 72.9 6.2 | 7.7 74.1 6.2 | 20.1 56.3 * | * * | * 88.6 * | 7.8 58.9 10.8 | 7.6 79.8 4.6 | 20.9 55.4 * |
| Akron, Ohio | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 384 2,877 103 | 215 2,061 51 | 158 763 50 | 2 5 - | 9 48 2 | 3 23 - | 212 2,039 51 | 158 761 50 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.1 83.8 3.0 | 9.2 88.8 2.2 | 15.0 73.2 4.8 | * * | * 77.4 * | * 88.5 * | 9.2 88.8 2.2 | 15.0 73.1 4.8 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origin | | | |
|---|---------------------|---------------------|----------------------|---------------------------|---------------------------------|---------------------|-----------------------------|-------------------------|
| Area, birthweight and prenatal care | İ | | Ra | ace | I | Hi | spanic Origin | (1) |
| Area, Brieffwerghe and prenaeur cure | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanio black |
| | | | | | Percent | | | |
| Albany, New York | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 138 940 76 | 53 536 23 | 80 360 46 | - 4 - | 5 40 7 | 10 64 2 | 34 361 12 | 72 337 41 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.3 81.2 6.6 | 8.2 86.9 3.7 | 15.4 74.4 9.5 | * * | * 76.9 * | 76.2 * | 8.3 90.3 * | 15.1 75.9 9.2 |
| Albuquerque, New Mexico | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 582 5,397 453 | 516 4,939 413 | 29 114 16 | 21 213 19 | 16 131 5 | 290 2,516 302 | 233 2,454 114 | 25 110 15 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.7 74.4 6.2 | 7.5 74.9 6.3 | 14.1 59.4 * | 7.1 75.0 * | 70.1 | 7.5 68.4 8.2 | 7.5 82.8 3.8 | 13.2 62.5 * |
| Alexandria city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 139 1,246 131 | 65 801 63 | 59 351 58 | - 2 1 | 15 92 9 | 21 215 38 | 43 587 25 | 59 350 58 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.8 72.6 7.6 | 6.0 76.5 6.0 | 10.5 64.9 10.7 | * * | * 73.0 * | 5.8 59.6 10.5 | 6.1 85.3 3.6 | 10.7 64.9 10.8 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|---------------------|---------------------|----------------------|-------------------------------|-----------------------------|---------------------|-------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | 1 | Ні | ispanic Origin | 1 (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | 13 87 12 * 68.5 * 39 110 29 19.6 55.8 14.7 |
| | | | | | Percent | | | |
| Allentown, Pennsylvania | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 166 1,239 75 | 148 1,103 55 | 16 99 14 | - 3 - | 2 34 6 | 55 335 25 | 95 780 32 | 87 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.0 80.1 4.8 | 10.3 82.0 4.1 | * 66.9 * | * * | * 68.0 * | 10.8 73.5 5.5 | 10.0 85.9 3.5 | 68.5 |
| Amarillo, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 282 2,195 159 | 238 2,029 129 | 40 113 29 | 1 13 - | 3 40 1 | 80 583 64 | 159 1,448 65 | 110 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.5 74.6 5.4 | 8.9 75.9 4.8 | 19.7 56.2 14.4 | * * | * 80.0 * | 8.7 64.2 7.0 | 9.0 81.8 3.7 | 55.8 |
| Anaheim, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 379 5,401 313 | 310 4,660 291 | 26 133 7 | - 12 - | 43 596 15 | 212 3,188 233 | 98 1,479 58 | 128 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.4 76.6 4.4 | 5.0 75.4 4.7 | 15.4 78.7 * | * * | 6.2 86.5 * | 4.8 71.9 5.3 | 5.6 84.5 3.3 | 16.0 78.5 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|---------------------------|--------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | R | ace | I | ні | spanic Origin | 2 31 20 207 5 4 8 12.7 1 85.2 3 * |
| Area, Direnweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | |
| | | | | | Percent | | | |
| Anchorage, coext. with Anchorage city, | Alaska | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 268 3,477 80 | 177 2,660 43 | 33 228 6 | 45 412 25 | 13 177 6 | 20 314 9 | 162 2,429 35 | 31 207 4 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.5 85.2 2.0 | 5.7 86.4 1.4 | 12.1 84.8 * | 8.7 80.3 4.9 | * 79.7 * | 5.1 79.9 * | 5.8 87.1 1.3 | 12.7 85.2 * |
| Ann Arbor, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 65 1,009 7 | 45 774 4 | 12 86 3 | - 3 - | 8 146 - | - 32 - | 42 679 4 | 10 78 3 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 4.9 94.0 * | 4.5 95.3 * | * 82.7 * | * * | * 95.4 * | 91.4 * | 5.0 95.8 * | * 81.3 * |
| Arlington, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 411 4,558 253 | 301 3,699 192 | 83 559 43 | - 13 3 | 27 287 15 | 80 777 104 | 220 2,895 87 | 83 555 43 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 82.6 4.6 | 6.8 83.2 4.3 | 11.5 77.4 6.0 | * * | 8.1 86.2 * | 7.0 68.2 9.1 | 6.7 88.4 2.7 | 11.5 77.3 6.0 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|----------------------|---------------------------|-----------------------------|---------------------|-------------------------|---|
| Area birthusiaht and avanatal gave | | | R | ace | I | H: | ispanic Origin | . (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black 34 215 41 10.2 66.6 12.7 |
| | | | | | Percent | | | |
| Arlington, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 164 2,102 162 | 116 1,693 106 | 35 215 41 | - 3 - | 13 191 15 | 51 504 65 | 65 1,189 41 | 215 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.0 79.6 6.1 | 5.4 81.8 5.1 | 10.4 66.6 12.7 | * * * | * 78.0 * | 6.4 66.1 8.5 | 4.9 90.8 3.1 | 66.6 |
| Atlanta, Georgia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 869 6,511 391 | 127 2,139 80 | 729 4,249 301 | 10 | 13 113 10 | 35 541 58 | 92 1,602 22 | 4,225 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.7 81.3 4.9 | 5.2 88.4 3.3 | 13.2 78.1 5.5 | * * * | * 79.6 * | 4.7 75.0 8.0 | 5.4 94.3 1.3 | 78.2 |
| Aurora, Colorado | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 429 3,516 204 | 311 2,855 155 | 86 456 26 | - 26 6 | 32 179 17 | 60 478 83 | 251 2,393 73 | 439 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.9 82.5 4.8 | 9.0 83.5 4.5 | 14.6 78.8 4.5 | * 68.4 * | 13.9 78.9 * | 7.8 63.0 10.9 | 9.3 89.4 2.7 | 14.3 79.1 4.7 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|-----------------------|-------------------------------|------------------|---------------------|-------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | 1 | Hi | ispanic Origin | n (1) |
| Area, Direnweight and prehacar care | All races | White | Black | American Indian (2) | Asian/Pacific | Hispanic | Non-Hispanio white | 152 908 76 12.1 73.3 6.1 84 375 28 14.3 70.8 5.3 |
| | | | | | Percent | | | |
| Austin, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 693 8,230 430 | 496 6,854 342 | 157 930 78 | 1 24 - | 39 422 10 | 252 2,952 286 | 249 3,930 58 | 908 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.8 81.5 4.3 | 5.9 82.2 4.1 | 12.2 73.3 6.2 | * 88.9 * | 8.3 91.1 * | 6.1 72.0 7.0 | 5.8 91.9 1.4 | 73.3 |
| Bakersfield, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 487 4,666 307 | 381 4,097 273 | 84 378 28 | 6 42 - | 16 149 6 | 203 2,011 176 | 177 2,098 97 | 375 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.1 73.4 4.8 | 6.4 73.6 4.9 | 14.2 70.7 5.2 | * 82.4 * | * 74.9 * | 6.5 68.7 6.0 | 6.3 78.9 3.6 | 70.8 |
| Baltimore city, Maryland | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,484 6,647 486 | 274 2,389 87 | 1,199 4,161 398 | 3 24 - | 8 73 1 | 12 64 6 | 262 2,314 79 | 4,134 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 14.3 77.6 5.7 | 9.2 88.6 3.2 | 16.5 72.3 6.9 | * 80.0 * | * 83.9 * | 66.7 * | 9.2 89.6 3.1 | 16.5 72.4 6.9 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|--------------------|---------------------|-------------------------------|-----------------------------|----------------|-------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black 385 1,850 173 14.3 68.7 6.4 103 730 34 12.1 85.4 4.0 |
| | | | | | Percent | | | |
| Baton Rouge, Louisiana | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 537 3,940 205 | 139 1,946 28 | 386 1,858 173 | - 5 - | 12 131 4 | 5 82 1 | , - | 1,850 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.7 78.8 4.1 | 6.5 90.6 1.3 | 14.3 68.8 6.4 | * * * | * 89.1 * | * 86.3 * | 6.5 90.8 1.3 | 68.7 |
| Beaumont, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 146 1,534 51 | 43 741 15 | 103 730 34 | - 3 - | - 60 2 | 8 144 7 | 35 596 8 | 730 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.5 88.9 3.0 | 5.4 92.5 * | 12.1 85.4 4.0 | * * | * 89.6 * | * 86.2 * | 5.5 94.2 * | 85.4 |
| Berkeley, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 62 890 7 | 30 576 3 | 26 189 4 | - 4 - | 6 121 - | 6 129 1 | 22 434 2 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.4 92.8 * | 4.9 94.6 * | 11.8 85.9 * | * * | * 96.0 * | * 93.5 * | 4.8 95.0 * | 12.5 86.9 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|-----------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | 1 (1) |
| area, pirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Birmingham, Alabama | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 468 3,111 197 | 74 944 39 | 389 2,129 154 | - 3 - | 5 35 4 | 4 28 5 | | 389 2,127 154 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.3 75.5 4.8 | 6.7 85.0 3.5 | 13.0 71.8 5.2 | * * | 81.4 * | * 59.6 * | 6.6 86.2 3.1 | 13.0 71.8 5.2 |
| Boise City, Idaho | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 168 2,291 74 | 156 2,200 69 | 2 17 1 | 2 10 - | 8 64 4 | 5 114 8 | 1,991 | 2 17 1 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.9 84.5 2.7 | 5.7 85.0 2.7 | * * | * * | * 75.3 * | * 70.4 * | 5.9 86.1 2.5 | * * * |
| Boston, Massachusetts | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 671 5,847 312 | 265 3,205 103 | 372 2,203 190 | - 6 2 | 34 433 17 | 129 1,149 57 | 158 2,348 60 | 340 1,876 174 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.0 79.4 4.2 | 6.8 84.1 2.7 | 12.2 73.6 6.3 | * * | 6.2 79.7 * | 8.6 80.1 4.0 | 5.8 86.3 2.2 | 12.9 71.8 6.7 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|--------------------------|--------------------------|-------------------------|---------------------------|------------------------------|-----------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | spanic Origin | 92 503 39 11.8 76.3 5.9 961 4,392 634 12.3 64.7 9.3 |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | |
| | | | | | Percent | | | |
| Bridgeport, Connecticut | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 217 1,506 86 | 112 927 41 | 92 512 41 | 4 4 - | 9 63 4 | 74 525 26 | 37 399 16 | 503 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.6 79.8 4.6 | 8.3 82.0 3.6 | 11.5 76.0 6.1 | * * | * 80.8 * | 9.2 80.9 4.0 | 6.7 83.5 * | 76.3 |
| Bronx borough, Bronx county, New York | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 2,255 13,692 1,740 | 1,085 7,462 852 | 1,112 5,737 842 | 10 2 | 58 483 44 | 1,033 7,213 922 | 166 1,437 111 | 4,392 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.6 66.8 8.5 | 8.7 68.4 7.8 | 11.0 64.7 9.5 | * * | 7.3 68.8 6.3 | 8.4 66.6 8.5 | 7.8 74.5 5.8 | 64.7 |
| Brooklyn borough, Kings county, New Yor | rk | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 3,607 24,957 2,997 | 1,456 13,612 1,299 | 1,947 9,333 1,403 | 1 18 1 | 203 1,994 294 | 785 6,089 773 | 747 7,910 564 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.8 70.4 8.5 | 6.9 72.8 6.9 | 11.7 67.1 10.1 | * * | 6.7 71.1 10.5 | 8.0 69.0 8.8 | 6.3 75.9 5.4 | 11.8 66.7 10.3 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|---------------------|---------------------|----------------------|---------------------------|-----------------------------|---------------------|-------------------------|--------------------------------------|
| Area, birthweight and prenatal care | | | R | ace | I | Hi | spanic Origin | (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | 248 1,029 246 13.6 61.0 14.6 444 3 |
| | | | | | Percent | | | |
| Buffalo, New York | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 468 3,326 439 | 190 2,091 159 | 273 1,166 268 | 1 17 5 | 4 52 7 | 26 246 37 | 35 603 28 | 1,029 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.8 72.4 9.6 | 7.1 80.3 6.1 | 13.5 61.9 14.2 | * * | * 65.8 * | 7.7 75.0 11.3 | 4.9 86.9 4.0 | 61.0 |
| Cedar Rapids, Iowa | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 102 1,558 27 | 96 1,464 23 | 6 44 3 | - 3 - | - 47 1 | 2 32 2 | 94 1,433 21 | 44 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.8 89.3 1.5 | 5.9 90.3 1.4 | * 74.6 * | * * | * 83.9 * | 76.2 * | 5.9 90.6 1.3 | |
| Charlotte, North Carolina | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 763 7,307 222 | 341 4,775 71 | 395 2,244 145 | - 13 - | 27 275 6 | 38 294 17 | 303 4,482 55 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.0 86.8 2.6 | 6.6 92.4 1.4 | 13.5 77.1 5.0 | * * | 8.3 84.6 * | 8.8 68.1 * | 6.4 94.6 1.2 | 13.6 77.2 5.0 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|--------------------------|--------------------------|--------------------------|---------------------------|---------------------------|------------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | spanic Origin | |
| Area, Diffilweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanio black |
| | | | | | Percent | | | |
| Chattanooga, Tennessee | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 266 1,708 125 | 104 1,073 38 | 155 601 85 | - 3 1 | 7 31 1 | 1 13 5 | 103 1,055 34 | 601 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.4 75.3 5.5 | 7.8 82.4 2.9 | 16.4 65.6 9.3 | * * | * 70.5 * | * * * | 7.9 83.0 2.7 | 65.6 |
| Chesapeake city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 182 2,375 68 | 90 1,745 18 | 88 569 48 | 1 7 - | 3 54 2 | 1 42 1 | 89 1,704 17 | 568 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.7 87.8 2.5 | 4.8 92.6 * | 11.7 75.5 6.4 | * * | * 91.5 * | * 85.7 * | 4.8 92.8 * | 75.5 |
| Chicago, Illinois | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5,573 38,020 3,474 | 1,889 21,700 1,326 | 3,513 14,810 2,022 | 11 46 11 | 160 1,464 115 | 1,098 12,622 959 | 803 9,147 374 | 14,732 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.6 74.2 6.8 | 6.6 78.3 4.8 | 15.8 68.4 9.3 | * 69.7 * | 8.1 78.4 6.2 | 6.2 73.7 5.6 | 7.3 85.7 3.5 | 15.8 68.4 9.3 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origin | n of mother | | |
|---|-----------------------|---------------------|----------------------|-------------------------------|-------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | R | ace | | Hi | spanic Origin | (1) |
| Area, bitchweight and prenacar care | All races | White | Black | American Indian (2) | Asian/Pacific | Hispanic | Non-Hispanic white | Non-Hispanio black |
| | | | | | Percent | | | |
| Chula Vista, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 143 2,345 144 | 115 1,946 128 | 13 134 7 | - 10 2 | 15 255 7 | 79 1,450 115 | 35 507 13 | 13 130 7 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 4.8 79.6 4.9 | 4.6 78.5 5.2 | 81.2 * | * * | * 87.9 * | 4.1 76.2 6.0 | 5.9 85.9 * | * 82.3 * |
| Cincinnati, Ohio | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 589 4,685 469 | 216 2,573 132 | 370 2,026 333 | - 9 - | 3 77 4 | 7 38 5 | 209 2,541 128 | 370 2,021 333 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.2 81.0 8.1 | 7.5 89.0 4.6 | 13.2 72.5 11.9 | * * * | * 89.5 * | * 82.6 * | 7.3 89.1 4.5 | 13.3 72.5 11.9 |
| Cleveland, Ohio | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,083 6,620 578 | 358 3,251 175 | 712 3,273 402 | 1 11 - | 12 85 1 | 49 441 38 | 310 2,800 138 | 709 3,245 400 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.7 74.2 6.5 | 8.7 79.8 4.3 | 14.1 69.2 8.5 | * * | * 80.2 * | 7.9 71.6 6.2 | 8.9 81.3 4.0 | 14.2 69.1 8.5 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|---------------------|---------------------------|------------------------------|--------------------|--|-------------------------|
| The birth weight and marked and | | | Ra | ace | | Hi | 360 4,067 132 7.6 86.1 2.8 84 860 36 | 1 (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | | Non-Hispanic black |
| | | | | | Percent | | | |
| Colorado Springs, Colorado | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 543 5,353 239 | 432 4,712 188 | 88 412 41 | 2 53 1 | 21 176 9 | 74 665 56 | 4,067 | 87 399 41 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.5 83.7 3.7 | 7.8 84.7 3.4 | 16.1 75.5 7.5 | * 81.5 * | 9.8 81.9 * | 8.5 76.3 6.4 | 86.1 | 16.4 75.7 7.8 |
| Columbus, Georgia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 277 1,833 158 | 90 915 39 | 185 885 119 | - 1 - | 2 32 - | 6 55 4 | 860 | 185 884 118 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.5 70.6 6.1 | 7.3 75.9 3.2 | 13.5 65.7 8.8 | * * | * 74.4 * | 67.9 * | 76.4 | 13.5 65.7 8.8 |
| Columbus, Ohio | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,006 9,155 508 | 562 6,574 295 | 418 2,291 196 | 3 27 1 | 23 263 16 | 8 137 8 | | 418 2,276 187 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.2 85.9 4.8 | 7.4 89.1 4.0 | 13.9 78.1 6.7 | * 84.4 * | 7.1 85.4 * | * 79.7 * | 7.5 89.4 3.9 | 14.0 78.3 6.4 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|--------------------------|------------------------|---------------------|---------------------------|------------------------------|---------------------|-------------------------|--------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | spanic Origin | 11 49 3 84.5 * 26 134 10 |
| Area, birthweight and prehatar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | |
| | | | | | Percent | | | |
| Concord, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 118 1,391 75 | 89 1,194 52 | 11 53 3 | - 6 - | 18 138 20 | 23 343 28 | 66 854 24 | 49 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.3 78.9 4.3 | 5.6 79.2 3.4 | * 85.5 * | * * | * 74.6 10.8 | 4.5 69.4 5.7 | 6.1 83.9 2.4 | * 84.5 * |
| Corpus Christi, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 333 3,459 190 | 297 3,258 178 | 26 140 10 | 2 8 1 | 8 53 1 | 198 2,065 152 | 99 1,200 26 | 134 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.3 79.4 4.4 | 6.9 79.7 4.4 | 13.2 72.5 * | * * | * 85.5 * | 6.9 75.3 5.5 | 6.9 88.6 1.9 | 72.0 |
| Dallas, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,776 16,182 1,099 | 1,027 11,711 785 | 696 3,912 289 | 5 46 5 | 48 513 20 | 686 7,093 629 | 342 4,634 154 | 696 3,899 289 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.0 76.0 5.2 | 6.4 76.4 5.1 | 12.5 74.1 5.5 | 73.0 * | 7.3 81.3 3.2 | 6.6 71.1 6.3 | 6.1 86.2 2.9 | 12.5 74.1 5.5 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | İ | | | Race and | Hispanic origin | n of mother | | |
|---|---------------------|---------------------|----------------------|---------------------------|-------------------|---------------------|-----------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | spanic Origin | C Non-Hispanic black 180 850 152 13.7 64.6 11.6 156 724 67 15.5 73.7 6.8 36 172 |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific | Hispanic | Non-Hispanio white | 180 850 152 13.7 64.6 11.6 15.5 73.7 6.8 |
| | | | | | Percent | | | |
| Dayton, Ohio | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 295 2,102 210 | 115 1,236 56 | 180 853 153 | - 2 1 | - 11 - | 1 22 4 | 114 1,217 53 | 850 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.4 73.9 7.4 | 7.7 82.2 3.7 | 13.6 64.5 11.6 | * * | * * | * 68.8 * | 7.7 82.4 3.6 | 64.6 |
| Denver, coext. with Denver city, Colora | ıdo | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 901 6,753 597 | 695 5,704 503 | 164 763 68 | 11 64 13 | 31 222 13 | 366 2,546 398 | 334 3,192 105 | 724 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.8 74.6 6.6 | 9.0 74.9 6.6 | 15.4 73.6 6.6 | 62.7 * | 10.3 75.0 * | 9.1 63.8 10.0 | 9.0 87.3 2.9 | 73.7 |
| Des Moines, Iowa | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 261 2,800 165 | 200 2,470 133 | 43 209 24 | 10 2 | 18 111 6 | 12 139 44 | 174 2,199 85 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 81.3 4.8 | 6.6 82.9 4.5 | 14.3 72.6 8.3 | * * | * 70.7 * | * 52.1 16.5 | 6.7 85.7 3.3 | 14.2 70.2 9.4 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|------------------------|---------------------|-----------------------|---------------------------|------------------------------|---------------------|-------------------------|----------------------------|
| | | | Ra | ace | | Hi | spanic Origin | nic Non-Hispanic black |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | |
| | | | | | Percent | | | |
| Detroit, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 2,116 10,545 964 | 208 1,908 112 | 1,884 8,461 838 | 5 20 5 | 19 156 9 | 38 427 21 | 142 1,372 77 | 8,369 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 12.8 71.2 6.5 | 7.7 78.2 4.6 | 13.9 69.6 6.9 | 69.0 * | * 77.6 * | 5.4 71.4 3.5 | 7.8 81.2 4.6 | 69.6 |
| District of Columbia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,196 4,541 832 | 148 1,382 129 | 1,034 3,070 692 | 2 2 1 | 12 87 10 | 62 418 69 | 88 956 58 | 3,059 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 14.3 64.6 11.8 | 7.2 77.4 7.2 | 16.7 60.0 13.5 | * * | * 70.2 * | 8.0 60.8 10.0 | 7.1 87.5 5.3 | 60.0 |
| Durham, North Carolina | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 290 2,129 89 | 82 1,048 9 | 198 970 79 | - 5 - | 10 106 1 | 10 136 4 | 72 915 5 | 967 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.5 84.7 3.5 | 7.3 93.5 * | 15.5 76.3 6.2 | * * | * 90.6 * | * 85.0 * | 7.5 94.9 * | 15.5 76.3 6.2 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|-----------------------|-----------------------|----------------------|---------------------------|---------------------------------|-----------------------|-------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | | H: | ispanic Origin | 1 (1) Non-Hispanic black Non-Hispanic black |
| Area, Dirthweight and prematar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| El Monte, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 214 2,459 97 | 189 2,124 89 | - 9 1 | - 2 - | 25 324 7 | 182 2,050 84 | 7 78 6 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.9 79.8 3.1 | 7.0 78.4 3.3 | * * | * * | 7.0 90.8 * | 7.0 78.6 3.2 | * 72.9 * | * |
| El Paso, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 961 7,759 1,533 | 914 7,503 1,514 | 31 166 15 | 2 9 2 | 14 81 2 | 825 6,537 1,440 | 92 975 76 | 157 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.1 62.9 12.4 | 7.0 62.7 12.6 | 12.0 67.5 * | * * | * 81.0 * | 7.0 61.0 13.4 | 6.5 76.8 6.0 | 68.6 |
| Elizabeth, New Jersey | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 154 1,316 183 | 72 926 93 | 76 343 89 | 1 1 - | 5 46 1 | 56 687 76 | 18 263 25 | 311 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.9 69.3 9.6 | 5.4 71.6 7.2 | 13.2 62.3 16.2 | * * | * 86.8 * | 5.6 69.3 7.7 | * 78.3 7.4 | 14.0 61.5 16.2 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|-----------------------------|---------------------|-------------------------|--------------------|
| Area, birthweight and prenatal care | | | R | ace | | Hi | ispanic Origin | . (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Erie, Pennsylvania | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 155 1,349 78 | 118 1,183 52 | 35 154 25 | 1 1 - | 1 11 1 | 5 47 7 | 113 1,137 45 | 152 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.2 78.6 4.5 | 7.7 82.0 3.6 | 10.7 60.2 9.8 | * * | * * * | * 59.5 * | 7.8 83.2 3.3 | 60.3 |
| Escondido, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 156 2,071 203 | 139 1,914 195 | 6 41 2 | - 16 - | 11 100 6 | 68 964 158 | 71 956 36 | 37 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.7 75.4 7.4 | 5.4 74.9 7.6 | * 78.8 * | * * | * 82.6 * | 4.7 66.2 10.9 | 6.4 86.5 3.3 | 80.4 |
| Eugene, Oregon | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 106 1,273 83 | 98 1,183 77 | 5 19 1 | 2 20 1 | 1 51 4 | 4 71 8 | 94 1,112 69 | 19 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.4 77.0 5.0 | 6.3 76.8 5.0 | * * * | * 83.3 * | * 77.3 * | 69.6 * | 6.5 77.3 4.8 | * * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origin | n of mother | | |
|---|---------------------|--------------------|---------------------|---------------------------|-------------------|----------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | Ні | spanic Origin | 1 (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Evansville, Indiana | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 142 1,547 60 | 117 1,359 43 | 25 173 14 | - 1 1 | | 1 17 - | 116 1,344 43 | 25 172 14 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.6 82.4 3.2 | 7.3 84.4 2.7 | 9.9 71.2 * | * * | * * * | * * | 7.3 84.4 2.7 | 10.0 71.1 * |
| Fayette, coext. with Lexington-Fayette | , Kentucky | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 280 2,734 107 | 222 2,308 63 | 55 387 43 | - - - | 3 39 1 | 3 39 6 | 219 2,268 57 | 55 383 43 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.5 85.4 3.3 | 8.2 88.2 2.4 | 10.0 72.1 8.0 | * * | * 86.7 * | * 73.6 * | 8.2 88.5 2.2 | 10.1 71.9 8.1 |
| Flint, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 370 2,476 104 | 132 1,162 36 | 237 1,298 68 | 1 7 - | - 9 - | 2 32 2 | 130 1,111 34 | 236 1,297 68 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 12.5 83.5 3.5 | 9.9 87.4 2.7 | 14.8 80.4 4.2 | * * | * * * | * 84.2 * | 10.2 87.4 2.7 | 14.7 80.4 4.2 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|---------------------------------|---------------------|-------------------------|------------------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | Danic Non-Hispanic black 109 |
| Area, birthweight and prehatar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | |
| | | | | | Percent | | | |
| Fort Lauderdale, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 391 3,179 116 | 130 1,825 41 | 259 1,307 75 | - 7 - | 2 40 - | 23 371 14 | 1,462 | 256 1,296 74 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.8 82.9 3.0 | 6.2 89.2 2.0 | 14.1 75.3 4.3 | * * * | * 85.1 * | 5.1 85.9 * | 90.0 | 14.1 75.3 4.3 |
| Fort Wayne, Indiana | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 311 2,793 132 | 206 2,281 75 | 100 456 55 | 1 12 - | 4 44 2 | 22 125 10 | 2,141 | 98 449 53 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.9 83.2 3.9 | 7.6 86.6 2.8 | 14.5 69.6 8.4 | * * * | * 81.5 * | 12.4 72.3 * | 87.6 | 14.5 69.8 8.2 |
| Fort Worth, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 753 6,986 622 | 472 5,318 461 | 260 1,432 145 | 1 18 1 | 20 218 15 | 247 2,277 343 | 225 3,043 119 | 260 1,429 145 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.0 74.8 6.7 | 6.6 74.8 6.5 | 13.5 74.5 7.5 | * * | 6.9 75.7 * | 7.0 64.2 9.7 | 6.3 85.3 3.3 | 13.5 74.5 7.6 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|---------------------------------|---------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | (1) |
| Area, bitthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanio black |
| | | | | | Percent | | | |
| Fremont, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 180 2,854 56 | 79 1,633 31 | 16 122 - | - 9 1 | 85 1,090 24 | 30 466 11 | 51 1,177 20 | 16 120 - |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.7 91.6 1.8 | 4.4 91.3 1.7 | * 93.8 * | * * | 7.1 91.9 2.0 | 5.6 87.8 * | 4.0 92.8 1.6 | 93.8 |
| Fresno, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 661 7,580 332 | 467 5,753 259 | 95 618 40 | 4 63 5 | 95 1,146 28 | 313 3,538 207 | 157 2,250 56 | 93 605 37 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.9 79.7 3.5 | 6.5 80.5 3.6 | 11.8 77.2 5.0 | * 70.8 * | 6.4 77.3 1.9 | 6.8 76.9 4.5 | 6.0 86.9 2.2 | 11.8 77.3 4.7 |
| Fullerton, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 91 1,639 66 | 77 1,387 58 | 7 55 4 | - 6 - | 7 191 4 | 35 679 36 | 44 712 22 | 6 53 4 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 4.5 82.2 3.3 | 4.5 81.9 3.4 | * 78.6 * | * * | * 85.3 * | 3.9 75.4 4.0 | 5.5 89.2 2.8 | * 79.1 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|---------------------|--------------------|----------------------|---------------------------|-----------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | 1 | Hi | spanic Origin | (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Garden Grove, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 156 2,449 95 | 86 1,604 75 | 3 32 3 | - 11 1 | 67 802 16 | 59 1,004 58 | 27 605 17 | 3 32 3 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.2 81.4 3.2 | 4.2 78.6 3.7 | 74.4 * | * * | 7.3 88.1 * | 4.4 75.2 4.3 | 3.8 85.1 * | * 74.4 * |
| Garland, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 232 2,554 126 | 181 2,113 96 | 36 260 25 | - 11 - | 15 170 5 | 70 690 63 | 111 1,425 33 | 36 257 25 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.7 76.4 3.8 | 6.4 77.1 3.5 | 9.2 68.2 6.6 | * * | * 81.7 * | 6.4 66.0 6.0 | 6.3 83.8 1.9 | 9.3 68.4 6.6 |
| Gary, Indiana | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 264 1,279 199 | 26 226 26 | 238 1,049 172 | - 3 1 | - 1 - | 7 40 8 | 18 182 19 | 238 1,044 172 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 12.9 62.8 9.8 | 7.8 70.2 8.1 | 14.0 61.5 10.1 | * * | * * | * 56.3 * | * 73.7 * | 14.0 61.4 10.1 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|------------------------------|--------------------|-------------------------|-------------------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | spanic Origin | 20 113 9 12.1 70.6 * |
| Area, Dirthweight and prehacar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | |
| | | | | | Percent | | | |
| Glendale, Arizona | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 260 2,932 201 | 231 2,711 182 | 21 116 9 | 4 35 6 | 4 70 4 | 85 714 83 | 147 2,006 99 | 113 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.9 78.8 5.4 | 6.7 79.6 5.3 | 12.3 69.9 * | 59.3 * | * 79.5 * | 8.1 68.9 8.0 | 6.1 84.2 4.2 | 70.6 |
| Glendale, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 136 2,061 45 | 102 1,670 38 | 4 20 1 | - 4 - | 30 367 6 | 43 727 24 | 56 936 14 | 18 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.7 86.7 1.9 | 5.3 86.8 2.0 | * 80.0 * | * * | 7.1 87.0 * | 4.9 83.1 2.7 | 5.4 89.8 * | * * |
| Grand Rapids, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 355 2,901 140 | 209 2,323 85 | 138 507 52 | 3 18 - | 5 53 3 | 40 339 22 | 168 1,965 59 | 134 499 51 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.6 71.6 3.5 | 6.7 75.4 2.8 | 15.6 58.6 6.0 | * * | * 68.8 * | 7.4 63.6 4.1 | 6.6 78.1 2.3 | 15.4 58.5 6.0 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|---------------------------------|---------------------|-------------------------|-------------------------|
| Amon high horight and property gave | | | R | ace | Ι | Hi | spanic Origin | . (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Greensboro, North Carolina | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 317 2,508 74 | 106 1,343 22 | 196 1,062 46 | 3 14 1 | 12 89 5 | 5 61 6 | 101 1,291 16 | 196 1,057 46 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.9 86.4 2.5 | 7.2 91.2 1.5 | 15.1 82.0 3.6 | * * * | * 77.4 * | * 70.9 * | 7.2 92.4 * | 15.2 81.9 3.6 |
| Hampton city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 175 1,650 91 | 63 968 28 | 110 640 60 | - 5 - | 2 37 3 | 2 39 2 | 61 933 26 | 110 636 60 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.4 79.0 4.4 | 5.6 86.0 2.5 | 12.1 70.3 6.6 | * * | * 75.5 * | 81.3 | 5.6 86.3 2.4 | 12.1 70.2 6.6 |
| Hartford, Connecticut | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 271 1,528 59 | 137 790 42 | 130 713 16 | - 1 - | 4 24 1 | 115 624 39 | 21 162 4 | 129 699 15 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 12.0 85.2 3.3 | 10.5 82.4 4.4 | 14.1 88.5 * | * * | * 88.9 * | 10.8 81.7 5.1 | 9.1 85.3 * | 14.5 88.6 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|--------------------|----------------------|---------------------------|---------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | Ι | Hi | ispanic Origin | (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Hayward, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 175 2,360 73 | 105 1,549 51 | 40 333 13 | 1 18 - | 29 460 9 | 56 927 32 | 645 | 38 321 13 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.2 86.1 2.7 | 5.6 85.4 2.8 | 10.2 85.8 * | * * | 5.4 88.6 * | 5.0 84.8 2.9 | 6.7 86.6 * | 10.0 85.6 * |
| Hialeah, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 152 2,120 30 | 141 2,029 27 | 9 73 3 | 2 5 - | 13 | 124 1,685 24 | 378 | 5 43 2 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.2 86.1 1.2 | 6.0 86.4 1.1 | * 76.8 * | * * | * * | 6.3 85.4 1.2 | 90.0 | * 76.8 * |
| Hollywood, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 190 1,660 134 | 112 1,207 64 | 73 390 68 | - 18 2 | 5 45 - | 31 345 24 | | 68 371 65 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.0 79.8 6.4 | 7.6 83.4 4.4 | 12.8 69.8 12.2 | * * | * 83.3 * | 6.9 77.9 5.4 | 8.3 85.5 4.2 | 12.5 69.6 12.2 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|--------------------------|--------------------------|-----------------------|---------------------------|---------------------------------|------------------------|-------------------------|-------------------------------|
| Area hirthweight and avenue large | | | Ra | ace | 1 | Hi | spanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | 1,242 7,254 457 12.7 77.1 4.9 |
| | | | | | Percent | | | |
| Honolulu, Hawaii | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 340 4,170 128 | 62 1,113 11 | 8 121 2 | 3 32 2 | 267 2,904 113 | 23 340 5 | 56 969 11 | 116 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.0 88.3 2.7 | 5.1 93.2 * | * 87.1 * | * 88.9 * | 7.6 86.6 3.4 | 5.8 88.5 * | 5.4 93.6 * | |
| Houston, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 3,226 31,917 1,552 | 1,850 22,673 1,042 | 1,247 7,294 461 | 2 36 1 | 127 1,914 48 | 1,259 14,820 890 | 591 7,894 158 | 7,254 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.7 80.0 3.9 | 6.3 80.3 3.7 | 12.6 77.1 4.9 | * 83.7 * | 5.7 87.3 2.2 | 6.1 76.0 4.6 | 6.7 90.0 1.8 | 77.1 |
| Huntington Beach, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 148 2,345 52 | 132 2,080 50 | 1 18 - | 1 7 - | 14 240 2 | 35 545 23 | 93 1,532 27 | 17 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.6 88.9 2.0 | 5.6 88.4 2.1 | * * | * * | * 94.1 * | 5.1 80.3 3.4 | 5.6 91.7 1.6 | * * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|------------------------|---------------------|---------------------|-------------------------------|------------------|-----------------|-------------------------|---------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Ні | ispanic Origin | |
| Area, Birthweight and prenatar care | All races | White | Black | American Indian (2) | Asian/Pacific | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Huntsville, Alabama | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 165 1,864 81 | 57 1,189 27 | 105 612 52 | 1 4 - | 2 59 2 | 3 27 2 | 56 1,163 26 | 612 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 83.7 3.6 | 4.3 90.6 2.1 | 12.5 73.1 6.2 | * * | * 81.9 * | 73.0 | 4.4 90.9 2.0 | 73.3 |
| Independence, Missouri | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 108 1,243 56 | 102 1,191 53 | 5 32 1 | 1 8 - | 12 2 | 7 40 3 | 95 1,152 50 | 31 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.1 83.3 3.8 | 7.1 84.2 3.7 | * 69.6 * | * * | * * * | * 81.6 * | 6.8 84.3 3.7 | * 68.9 * |
| Indianapolis, Indiana | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,242 10,088 439 | 732 7,826 202 | 499 2,125 230 | 2 9 - | 9 128 7 | 15 193 19 | 717 7,625 182 | 2,120 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.3 79.8 3.5 | 7.6 85.0 2.2 | 14.1 65.2 7.1 | * * | * 81.0 * | 64.3 * | 7.7 85.6 2.0 | 14.1 65.3 7.1 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| 1 | | | | Race and | d Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|-----------------------------|--------------------|-------------------------|-----------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | H | ispanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Inglewood, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 205 2,337 113 | 115 1,606 86 | 83 694 21 | 2 4 - | 5 33 6 | 114 1,566 81 | 3 48 5 | 81 685 21 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.8 78.8 3.8 | 5.5 78.5 4.2 | 9.5 80.1 2.4 | * * | 67.3 * | 5.6 78.4 4.1 | * 81.4 * | 9.4 80.2 2.5 |
| Irvine, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 81 1,571 15 | 53 1,172 13 | 3 59 1 | 1 5 - | 24 335 1 | 11 181 6 | 43 992 7 | 3 58 1 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 4.8 94.1 * | 4.2 93.7 * | * 85.5 * | * * | 6.9 97.7 * | 85.4 * | 4.1 95.2 * | * 86.6 * |
| Irving, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 222 2,416 182 | 166 1,938 143 | 39 250 29 | 1 7 - | 16 221 10 | 59 801 61 | 106 1,138 82 | 39 248 29 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.9 77.3 5.8 | 6.4 76.9 5.7 | 11.4 75.5 8.8 | * * | * 84.4 * | 5.0 69.8 5.3 | 7.5 82.8 6.0 | 11.5 75.6 8.8 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origin | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|---------------------------------|--------------------|-----------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | spanic Origin | n (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanio black |
| | | | | | Percent | | | |
| Jackson, Mississippi | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 443 2,370 154 | 63 584 17 | 376 1,764 137 | 2 4 - | 2 18 - | - 6 - | 63 574 17 | 376 1,764 137 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 14.2 76.1 4.9 | 9.5 88.8 * | 15.4 72.6 5.6 | * * | * * * | * * | 9.7 88.9 * | 15.4 72.6 5.6 |
| Jacksonville, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 914 8,672 323 | 412 5,790 145 | 475 2,629 172 | 2 17 - | 25 236 6 | 11 240 12 | 403 5,582 133 | 473 2,597 172 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.2 82.7 3.1 | 5.9 87.3 2.2 | 12.5 74.3 4.9 | * * | 7.8 80.8 * | * 81.4 * | 6.0 87.6 2.1 | 12.6 74.2 4.9 |
| Jersey City, New Jersey | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 457 2,564 269 | 143 1,196 106 | 260 897 128 | 3 7 - | 51 464 35 | 107 806 71 | 46 496 39 | 245 774 121 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.4 72.9 7.7 | 8.1 74.8 6.6 | 16.3 68.4 9.8 | * * | 8.0 77.9 5.9 | 8.8 73.2 6.4 | 6.6 78.7 6.2 | 17.2 67.2 10.5 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | ispanic Origin | (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Kansas City, Kansas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 225 1,946 117 | 103 1,225 58 | 118 661 52 | - 8 - | 4 52 7 | 22 261 25 | 81 961 31 | 118 655 52 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.1 79.4 4.8 | 6.8 81.2 3.8 | 13.6 76.6 6.0 | * * | * 74.3 * | 5.8 69.6 6.7 | 7.2 85.1 2.7 | 13.7 76.4 6.1 |
| Kansas City, Missouri | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 665 5,903 283 | 291 3,951 112 | 360 1,804 159 | 1 21 4 | 13 127 8 | 27 377 16 | 261 3,562 96 | 360 1,797 159 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.1 84.6 4.1 | 6.4 89.1 2.5 | 14.1 76.7 6.8 | * 65.6 * | * 81.4 * | 5.3 81.8 * | 6.4 89.9 2.4 | 14.1 76.7 6.8 |
| Knoxville, Tennessee | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 259 2,226 48 | 189 1,809 27 | 66 368 21 | - 6 - | 4 43 - | 2 19 - | 187 1,790 27 | 66 368 21 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.1 88.2 1.9 | 9.2 90.1 1.3 | 14.0 80.3 4.6 | * * | * 82.7 * | * * | 9.3 90.3 1.4 | 14.0 80.3 4.6 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | į | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|-----------------------------|---------------------|-------------------------|---------------|
| Area, birthweight and prenatal care | - | | Ra | ace | I | Hi | ispanic Origin | n (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | 13.3 85.5 4.0 |
| | | | | | Percent | | | |
| Lakewood, Colorado | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 126 1,441 49 | 117 1,362 46 | - 15 1 | 1 13 - | 8 51 2 | 16 209 15 | 101 1,151 31 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.7 88.5 3.0 | 7.6 88.4 3.0 | * * | * * * | * 89.5 * | 77.1 * | 7.9 90.8 2.4 | * |
| Lansing, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 221 2,031 114 | 140 1,488 88 | 73 453 21 | - 10 - | 8 80 5 | 16 209 18 | 119 1,181 54 | 447 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.3 86.8 4.9 | 8.2 87.8 5.2 | 13.5 85.3 4.0 | * * | * 79.2 * | * 82.9 * | 9.0 90.4 4.1 | 85.5 |
| Laredo, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 271 2,965 666 | 267 2,946 664 | - 1 2 | - - - | 4 18 - | 265 2,886 660 | 2 60 4 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.7 62.1 13.9 | 5.6 62.0 14.0 | * * * | * * | * * * | 5.7 61.7 14.1 | * 80.0 * | * * * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origin | n of mother | | (1) Non-Hispanic black 154 829 82 12.9 71.9 7.1 |
|---|---------------------|---------------------|----------------------|---------------------------|---------------------------------|---------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | spanic Origin | n (1) |
| Area, bitchweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | |
| | | | | | Percent | | | |
| Las Vegas, Nevada | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 874 8,948 564 | 631 7,396 443 | 158 842 82 | 10 76 6 | 75 634 33 | 193 2,035 278 | 437 5,360 159 | 829 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.5 79.3 5.0 | 6.7 80.2 4.8 | 13.1 72.1 7.0 | * 71.7 * | 9.2 80.4 4.2 | 6.2 68.3 9.3 | 6.9 86.0 2.6 | 71.9 |
| Lincoln, Nebraska | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 218 2,672 67 | 194 2,473 53 | 14 64 5 | 2 21 4 | 8 114 5 | 8 92 5 | 186 2,360 48 | 13 62 5 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.1 86.4 2.2 | 6.9 87.5 1.9 | 66.0 * | * 65.6 * | * 82.0 * | 74.8 * | 6.9 88.0 1.8 | 65.3 * |
| Little Rock, Arkansas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 307 2,213 275 | 98 1,204 73 | 203 955 193 | 3 8 1 | 3 46 8 | 5 35 10 | 94 1,168 64 | 202 953 191 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.4 78.0 9.7 | 6.7 86.4 5.2 | 14.1 69.4 14.0 | * * | * 79.3 * | 57.4 * | 6.7 87.6 4.8 | 14.1 69.5 13.9 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|--------------------------|--------------------------|---------------------|---------------------------|------------------------------|--------------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Ні | spanic Origin | (1) |
| Area, Direnweight and prenatar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanio black |
| | | | | | Percent | | | |
| Livonia, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 70 1,069 10 | 69 1,037 10 | - 4 - | - 3 - | 1 25 - | 3 17 - | 63 994 10 | - 4 - |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.0 95.4 * | 6.1 95.4 * | * * | * * | 96.2 * | * * | 5.8 95.5 * | * * |
| Long Beach, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 625 7,201 337 | 336 5,000 236 | 176 1,188 60 | 2 28 3 | 111 985 38 | 219 3,441 188 | 120 1,598 53 | 174 1,168 58 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.9 80.4 3.8 | 5.4 81.0 3.8 | 11.5 78.9 4.0 | * 80.0 * | 8.8 79.6 3.1 | 4.9 77.9 4.3 | 6.5 88.2 2.9 | 11.6 79.1 3.9 |
| Los Angeles, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 4,640 56,112 2,345 | 3,339 46,021 1,957 | 962 5,966 279 | 14 93 5 | 325 4,032 104 | 2,659 36,595 1,803 | 708 9,630 159 | 931 5,772 271 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.6 80.1 3.3 | 5.7 79.7 3.4 | 12.8 80.0 3.7 | * 74.4 * | 6.9 86.1 2.2 | 5.6 77.2 3.8 | 6.6 90.6 1.5 | 12.8 80.0 3.8 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|---------------------|---------------------------|-----------------------------|---------------------|-------------------------|-------------------------|
| Amon histhyroight and amonatal gave | | | Ra | ace | | Hi | ispanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Louisville, Kentucky | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 614 5,809 178 | 310 4,068 77 | 299 1,635 99 | 13 | 5 93 2 | 8 66 2 | , | 297 1,626 99 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.1 85.8 2.6 | 6.9 90.6 1.7 | 13.9 76.0 4.6 | * * | * 83.8 * | * 85.7 * | 6.8 90.7 1.7 | 13.9 75.9 4.6 |
| Lowell, Massachusetts | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 125 1,114 88 | 84 874 50 | 8 65 7 | 1 4 - | 32 171 31 | 30 193 34 | | 7 48 4 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.7 68.9 5.4 | 7.1 74.1 4.2 | * 67.7 * | * * | 9.5 51.2 9.3 | 9.9 63.3 11.1 | | * 71.6 * |
| Lubbock, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 263 2,489 142 | 211 2,236 123 | 45 206 19 | 2 9 - | 5 38 - | 97 819 94 | 113 1,387 30 | 45 203 18 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.1 78.8 4.5 | 7.3 79.8 4.4 | 14.6 68.4 * | * * | * 88.4 * | 7.7 67.5 7.7 | 7.0 89.0 1.9 | 14.8 68.4 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|--------------------------|-----------------------|----------------------|-------------------------------|------------------------------|---------------------|-----------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | spanic Origin | 188 1,074 55 14.2 81.1 4.2 |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | |
| | | | | | Percent | | | |
| Macon, Georgia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 234 1,656 63 | 46 567 8 | 188 1,074 55 | - 1 - | - 14 - | - 13 - | 46 554 8 | 1,074 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.9 84.5 3.2 | 7.5 92.2 * | 14.2 81.1 4.2 | * * | * * * | * * | 7.6 92.2 * | 81.1 |
| Madison, Wisconsin | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 176 2,279 50 | 136 1,926 28 | 32 184 13 | 2 12 2 | 6 157 7 | 8 96 9 | 129 1,832 19 | 184 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.8 87.9 1.9 | 6.5 91.5 1.3 | 12.2 70.0 * | * * | 76.2 * | * 72.7 * | 6.5 92.6 * | 70.5 |
| Manhattan borough, New York county, Ne | ew York | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,822 12,574 1,237 | 1,065 8,417 603 | 636 2,754 550 | - 8 - | 121 1,395 84 | 633 4,024 541 | 475 4,669 173 | 507 1,932 413 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.1 73.8 7.3 | 8.3 78.4 5.6 | 12.2 62.9 12.6 | * * | 5.9 73.5 4.4 | 8.5 66.1 8.9 | 7.8 87.7 3.2 | 14.0 63.0 13.5 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|-----------------------|---------------------------|-----------------------------|---------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | H: | ispanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Memphis, Tennessee | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,357 8,058 875 | 202 2,608 106 | 1,135 5,275 756 | 1 9 - | 19 166 13 | 9 129 28 | 193 2,489 78 | 1,132 5,267 754 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 12.2 73.6 8.0 | 6.7 86.5 3.5 | 14.5 68.4 9.8 | * * | * 76.9 * | * 63.2 13.7 | 88.2 | 14.5 68.4 9.8 |
| Mesa, Arizona | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 407 5,177 274 | 365 4,879 240 | 15 105 9 | 16 104 22 | 11 89 3 | 103 1,098 145 | 265 3,797 97 | 15 102 9 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.3 80.4 4.3 | 6.0 81.2 4.0 | * 77.8 * | * 56.2 11.9 | * 79.5 * | 6.3 68.1 9.0 | 85.9 | * 77.9 * |
| Mesquite, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 119 1,506 48 | 86 1,262 39 | 26 179 8 | - 2 - | 7 63 1 | 14 218 14 | 1,045 | 26 177 8 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.5 84.8 2.7 | 5.6 84.5 2.6 | 12.0 85.2 * | * * | * 88.7 * | 78.4 * | 5.8 85.9 2.1 | 12.1 85.1 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|------------------------|---------------------|----------------------|---------------------------|------------------------------|---------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | . (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | 776 4,254 419 12.7 70.5 6.9 684 3,432 492 13.2 66.3 9.5 |
| | | | | | Percent | | | |
| Miami, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,424 11,502 677 | 593 6,852 242 | 812 4,472 427 | 3 36 5 | 16 142 3 | 529 5,895 207 | 102 1,213 46 | 4,254 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.5 77.2 4.5 | 7.1 82.2 2.9 | 12.7 70.6 6.7 | * 59.0 * | * 80.2 * | 7.2 80.8 2.8 | 7.1 85.8 3.3 | 70.5 |
| Milwaukee, Wisconsin | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,104 8,369 706 | 374 4,551 179 | 686 3,453 494 | 8 77 5 | 36 288 28 | 106 1,001 83 | 269 3,578 98 | 3,432 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.8 74.5 6.3 | 6.8 83.4 3.3 | 13.2 66.3 9.5 | * 76.2 * | 7.8 61.9 6.0 | 7.4 70.3 5.8 | 6.6 87.7 2.4 | 66.3 |
| Minneapolis, Minnesota | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 479 3,710 444 | 229 2,544 189 | 191 853 166 | 26 104 34 | 33 209 55 | 34 180 48 | 185 1,924 137 | 846 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.3 69.3 8.3 | 6.6 78.1 5.8 | 12.3 59.8 11.6 | 10.9 49.5 16.2 | 6.1 45.5 12.0 | 8.5 47.6 12.7 | 7.2 79.4 5.7 | 12.4 59.8 11.6 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|---------------------------------|--------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | R | ace | 1 | Hi | ispanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | 234 1,173 163 13.6 68.0 9.5 |
| | | | | | Percent | | | |
| Mobile, Alabama | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 366 2,580 216 | 127 1,360 49 | 234 1,176 163 | - 2 - | 5 42 4 | 1 21 3 | 126 1,343 46 | 1,173 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.0 77.9 6.5 | 8.3 88.9 3.2 | 13.5 68.1 9.4 | * * | * 77.8 * | * 72.4 * | 8.3 89.2 3.1 | 68.0 |
| Modesto, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 260 3,062 125 | 198 2,668 100 | 23 121 8 | 6 48 4 | 33 225 13 | 82 1,154 61 | 117 1,533 44 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.6 78.6 3.2 | 5.9 79.4 3.0 | 15.0 79.6 * | * 76.2 * | 10.1 70.1 * | 5.2 74.0 3.9 | 6.4 83.8 2.4 | 16.2 80.3 * |
| Montgomery, Alabama | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 352 2,468 177 | 86 1,212 30 | 262 1,213 145 | - 3 - | 4 40 2 | 1 26 3 | 86 1,191 28 | 261 1,209 145 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.8 76.0 5.5 | 6.4 90.2 2.2 | 14.0 65.5 7.8 | * * | * 83.3 * | * 78.8 * | 6.5 90.4 2.1 | 14.0 65.4 7.8 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|-----------------------------|--------------------|-------------------------|------------------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | spanic Origin | (1) |
| Area, Birthweight and prehatar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | C Non-Hispanic black 56 348 16 |
| | | | | | Percent | | | |
| Moreno Valley, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 188 1,891 87 | 114 1,389 60 | 58 359 16 | - 8 - | 16 135 11 | 69 807 43 | 45 591 15 | 348 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.8 79.8 3.7 | 6.5 80.4 3.5 | 12.4 77.5 * | * * * | * 78.9 * | 6.4 76.9 4.1 | 6.5 85.7 * | 77.9 |
| Nashville-Davidson, Tennessee | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 745 7,141 238 | 396 4,940 103 | 324 1,996 123 | - 13 - | 25 192 12 | 25 258 24 | 371 4,688 79 | 1,991 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.2 89.2 3.0 | 7.3 91.5 1.9 | 13.6 84.3 5.2 | * * | 11.1 86.1 * | 7.4 76.8 7.1 | 7.3 92.5 1.6 | 84.3 |
| New Haven, Connecticut | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 188 1,257 78 | 67 685 25 | 114 527 49 | 1 1 2 | 6 44 2 | 35 340 18 | 32 344 9 | 109 474 44 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.4 76.7 4.8 | 7.5 82.9 3.0 | 13.5 70.1 6.5 | * * | * 78.6 * | 7.3 78.9 * | 7.8 88.0 * | 14.5 70.9 6.6 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|---|--------------------|--------------------------|---------------------------|-----------------------------|--------------------------|--------------------------|--------------------------|
| The birth wint and months and | All races White 10,708 5,092 77,623 45,628 8,110 3,917 8.7 7.3 72.1 74.7 7.5 6.4 ster 2,947 1,246 729 1,246 170 12.5 7.5 ster 60.0 65.3 | Ra | ace | | Hi | Hispanic Origin (1) | | |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| New York, New York | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 77,623 | 45,628 | 4,733 23,853 3,318 | 6 83 9 | 877 8,059 866 | 3,181 23,809 3,083 | 2,142 23,607 1,179 | 4,264 20,547 2,859 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 72.1 | 74.7 | 11.6 67.7 9.4 | 76.1 * | 7.0 71.2 7.6 | 7.8 67.1 8.7 | 6.7 82.5 4.1 | 12.2 67.9 9.4 |
| Newark, New Jersey | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 2,947 | 1,246 | 483 1,652 553 | - 4 - | 2 45 6 | 132 1,029 137 | 21 271 34 | 476 1,596 546 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | | | 15.9 56.4 18.9 | * * | 67.2 * | 7.9 63.2 8.4 | 5.3 72.3 9.1 | 16.2 56.5 19.3 |
| Newport News city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 281 2,574 115 | 112 1,591 23 | 163 907 88 | - 5 1 | 6 71 3 | 1 88 2 | 111 1,508 21 | 163 901 87 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.8 80.4 3.6 | 6.1 86.8 1.3 | 12.7 70.9 6.9 | * * | * 84.5 * | * 81.5 * | 6.4 87.2 1.2 | 12.8 70.9 6.8 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|---------------------|---------------------------|-----------------------------|---------------------|-------------------------|---|
| Area, birthweight and prenatal care | | | R | ace | I | Hi | ispanic Origin | (1) |
| area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | 220 1,271 190 11.2 64.9 9.7 308 1,995 62 12.7 84.4 2.6 |
| | | | | | Percent | | | |
| Norfolk city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 378 3,019 260 | 137 1,623 64 | 223 1,276 191 | 1 9 2 | 17 111 3 | 14 104 10 | 1,526 | 1,271 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.3 74.4 6.4 | 7.0 84.0 3.3 | 11.3 64.9 9.7 | * * * | * 78.2 * | * 78.2 * | 7.0 84.3 3.0 | 64.9 |
| Oakland, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 558 5,438 155 | 158 2,495 71 | 320 2,037 65 | 3 30 1 | 77 876 18 | 99 1,581 52 | | 1,995 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.7 87.0 2.5 | 5.5 88.7 2.5 | 12.9 84.4 2.7 | * 81.1 * | 7.6 88.5 * | 5.3 87.0 2.9 | | 84.4 |
| Oceanside, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 220 2,683 258 | 172 2,218 210 | 28 247 27 | - 20 5 | 20 198 16 | 81 798 163 | 93 1,431 48 | 241 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.0 74.0 7.1 | 5.7 74.2 7.0 | 8.3 73.3 8.0 | * 66.7 * | 7.5 74.4 * | 6.3 62.0 12.7 | 5.4 83.4 2.8 | 8.5 73.3 7.9 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|-----------------------------|---------------------|-------------------------|-------------------------|
| Two hinthusiaht and proposal gave | | | Ra | ace | Ι | Hi | ispanic Origin | 1 (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Oklahoma City, Oklahoma | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 614 4,533 287 | 383 3,449 162 | 189 744 90 | 23 169 27 | 19 171 8 | 62 347 32 | 321 3,110 129 | 189 742 90 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.5 78.2 5.0 | 7.3 81.9 3.8 | 13.4 66.7 8.1 | 7.1 62.6 10.0 | * 85.5 * | 6.4 70.7 6.5 | 7.5 83.4 3.5 | 13.5 66.6 8.1 |
| Omaha, Nebraska | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 386 4,681 179 | 276 3,859 110 | 98 691 59 | 3 44 7 | 9 87 3 | 27 396 35 | 245 3,418 74 | 98 687 59 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.9 84.3 3.2 | 6.2 87.0 2.5 | 10.4 73.2 6.3 | 71.0 * | * 78.4 * | 4.9 72.3 6.4 | 6.4 89.1 1.9 | 10.4 73.2 6.3 |
| Ontario, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 194 2,560 215 | 168 2,284 196 | 21 172 7 | 2 10 2 | 3 94 10 | 130 1,728 165 | 40 551 28 | 20 170 7 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.4 71.7 6.0 | 5.2 71.0 6.1 | 9.8 80.4 * | * * | * 74.6 * | 5.2 68.8 6.6 | 5.8 79.9 4.1 | 9.5 81.0 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|---------------------|-------------------------------|-----------------------------|--------------------|-------------------------|---|
| Dura historiaht and manatal and | | | Ra | ace | | Hi | spanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black 1 29 1 87.9 * 13.4 73.7 7.2 |
| | | | | | Percent | | | |
| Orange, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 92 1,773 98 | 88 1,613 94 | 1 31 1 | - 7 1 | 3 122 2 | 52 771 82 | 36 844 12 | 29 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 4.1 80.0 4.4 | 4.3 79.5 4.6 | * 88.6 * | * * | * 85.3 * | 4.7 69.8 7.4 | 3.9 91.1 * | * 87.9 * |
| Orlando, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 550 4,915 239 | 293 3,430 111 | 234 1,270 123 | 3 11 - | 20 204 5 | 92 961 36 | 203 2,480 75 | 1,258 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.2 83.3 4.1 | 7.4 87.3 2.8 | 13.4 73.8 7.1 | * * | 8.4 85.7 * | 8.0 83.6 3.1 | 7.2 88.9 2.7 | 73.7 |
| Orleans, coext. with New Orleans city, | Louisiana | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 939 5,777 462 | 82 1,244 33 | 837 4,355 420 | - 4 - | 20 174 9 | 8 141 6 | 74 1,119 28 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 12.4 76.3 6.1 | 6.0 90.5 2.4 | 14.0 72.8 7.0 | * * | 9.4 82.1 * | * 82.9 * | 6.1 91.6 2.3 | 14.0 72.7 7.0 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|--------------------|-------------------|---------------------------|-----------------------------|--------------------|-------------------------|--------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | 3 7 3 39 3 1 1 |
| area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Overland Park, Kansas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 121 1,781 16 | 110 1,688 14 | 7 39 1 | - 8 - | 4 46 1 | 4 46 1 | 1,623 | 39 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.5 95.6 * | 6.2 95.9 * | * 92.9 * | * * | * 92.0 * | 92.0 | 6.1 96.0 * | 92.9 |
| Oxnard, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 198 3,087 98 | 176 2,772 85 | 11 107 1 | 1 14 1 | 10 194 11 | 153 2,292 71 | 20 363 11 | 92 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.3 82.9 2.6 | 5.2 82.9 2.5 | * 84.9 * | * * | * 82.6 * | 5.5 81.9 2.5 | 4.9 89.0 * | |
| Pasadena, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 221 2,158 74 | 144 1,611 56 | 51 289 14 | - 3 - | 26 255 4 | 90 968 45 | 54 649 11 | 278 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.6 84.5 2.9 | 7.5 84.5 2.9 | 14.0 80.3 * | * * | 9.1 89.5 * | 7.4 80.1 3.7 | 7.6 91.4 * | 14.0 80.6 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|---------------------|--------------------|----------------------|-------------------------------|-----------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | ispanic Origin | (1) |
| Area, Direnweight and prenatar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Pasadena, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 171 2,055 90 | 166 2,005 89 | 3 24 - | - 4 1 | 2 22 - | 98 1,197 68 | 68 809 21 | 3 21 - |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.5 80.4 3.5 | 6.5 80.4 3.6 | * 82.8 * | * * | * 81.5 * | 6.1 76.9 4.4 | 7.1 86.2 2.2 | 84.0 |
| Paterson, New Jersey | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 280 2,123 231 | 144 1,432 99 | 127 635 129 | 1 2 - | 8 54 3 | 118 1,236 92 | 30 263 19 | 122 569 117 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.2 71.3 7.8 | 7.4 74.9 5.2 | 12.5 64.2 13.0 | * * | * 75.0 * | 7.0 74.3 5.5 | 8.5 75.8 * | 13.3 63.8 13.1 |
| Peoria, Illinois | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 144 1,578 81 | 65 1,115 35 | 76 418 46 | - 3 - | 3 42 - | 1 40 2 | 65 1,079 33 | 75 414 46 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 81.3 4.2 | 5.1 87.6 2.7 | 12.3 67.4 7.4 | * * | * 95.5 * | * 81.6 * | 5.3 87.9 2.7 | 12.2 67.2 7.5 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|--------------------------|--------------------------|-------------------------|-------------------------------|------------------------|-----------------------|-------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | | Ні | ispanic Origin | n (1) |
| Area, birthweight and prenatar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | 1,709 6,822 1,360 14.9 63.2 12.6 |
| | | | | | Percent | | | |
| Philadelphia, coext. with Phila. city, | Pennsylvani | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 2,637 15,682 1,881 | 799 7,997 434 | 1,728 6,930 1,373 | 3 31 7 | 107 724 67 | 268 1,752 178 | 545 6,342 269 | 6,822 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.6 72.1 8.7 | 8.0 82.6 4.5 | 14.8 63.3 12.5 | * 66.0 * | 9.5 68.5 6.3 | 10.3 70.0 7.1 | 7.3 86.7 3.7 | 63.2 |
| Phoenix, Arizona | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,636 16,271 2,061 | 1,402 14,767 1,813 | 151 779 124 | 46 371 89 | 37 354 35 | 681 6,304 1,385 | 726 8,494 437 | 766 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.0 70.6 8.9 | 6.7 71.2 8.7 | 13.0 67.3 10.7 | 6.7 54.6 13.1 | 7.7 74.8 7.4 | 6.3 59.3 13.0 | 7.0 83.4 4.3 | 67.4 |
| Pittsburgh, Pennsylvania | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 427 3,867 204 | 166 2,448 68 | 251 1,259 131 | 1 7 1 | 9 153 4 | 3 42 2 | 161 2,395 66 | 249 1,251 127 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.1 84.4 4.5 | 6.1 90.5 2.5 | 14.3 74.1 7.7 | * * | * 90.0 * | * 82.4 * | 6.0 90.7 2.5 | 14.3 74.2 7.5 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|---------------------|---------------------------|-----------------------------|---------------------|-------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | ispanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black 12 108 5 83.7 * 83.7 * 82 490 41 12.7 |
| | | | | | Percent | | | |
| Plano, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 177 2,660 51 | 142 2,270 44 | 13 111 5 | 1 4 - | 21 275 2 | 20 234 23 | 2,034 | 108 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.9 91.5 1.8 | 5.6 91.5 1.8 | 84.1 * | * * | 7.0 94.2 * | 5.9 72.2 7.1 | | 83.7 |
| Pomona, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 214 2,428 183 | 165 2,119 159 | 34 168 13 | - 3 1 | 15 138 10 | 153 1,863 138 | 263 | 161 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.4 73.1 5.5 | 5.7 72.9 5.5 | 13.7 68.6 * | * * | * 83.1 * | 5.9 71.9 5.3 | * 80.4 6.4 | 68.2 |
| Portland, Oregon | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 442 5,800 283 | 312 4,794 199 | 82 496 41 | 3 65 8 | 45 445 35 | 38 450 47 | 274 4,350 152 | 490 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.2 81.0 4.0 | 5.4 82.7 3.4 | 12.6 76.1 6.3 | * 69.9 * | 7.3 72.4 5.7 | 5.9 70.6 7.4 | 5.3 84.2 2.9 | 12.7 76.0 6.4 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|--------------------------|--------------------------|---------------------|-------------------------------|-----------------------------|------------------------------------|-------------------------|---|
| Area hinthweight and property gare | | | Ra | ace | I | 2 2 19 14 1 - * * * * * * * * 6.1 | | . (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | 124 610 75 13.5 66.2 8.1 |
| | | | | | Percent | | | |
| Portsmouth city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 175 1,242 93 | 48 609 17 | 124 612 75 | 1 2 - | 19 | 14 | 47 599 17 | 610 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.6 75.3 5.6 | 6.9 87.2 * | 13.5 66.2 8.1 | * * | * | * | 6.9 87.4 * | 66.2 |
| Providence, Rhode Island | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 209 1,760 46 | 128 1,315 25 | 63 298 15 | 4 29 2 | 118 | 696 | 54 534 6 | 232 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.9 83.6 2.2 | 6.8 86.0 1.6 | 12.9 78.8 * | 70.7 | 75.2 | 84.5 | 7.9 89.0 * | 81.7 |
| Queens borough, Queens county, New Yor | rk | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 2,581 21,428 1,980 | 1,163 12,029 1,074 | 943 5,420 463 | 5 40 6 | 470 3,939 437 | 660 5,857 814 | 502 6,129 278 | 906 5,141 436 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.9 73.3 6.8 | 6.4 74.1 6.6 | 11.8 74.1 6.3 | * 70.2 * | 7.4 69.8 7.7 | 6.3 65.0 9.0 | 6.4 84.8 3.8 | 12.0 74.1 6.3 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|--------------------|---------------------|---------------------------|---------------------------------|--------------------|---|----------------|
| Area, birthweight and prenatal care | | | Ra | ace | T | Hi | Hispanic Origin (1) Non-Hispanic (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | | |
| | | | | | Percent | | | |
| Raleigh, North Carolina | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 346 3,457 123 | 140 2,289 43 | 194 1,015 76 | 1 4 1 | 11 149 3 | 10 166 16 | 2,126 | 1,012 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.6 86.6 3.1 | 5.6 91.9 1.7 | 14.5 76.8 5.7 | * * * | * 87.6 * | 72.5 | 93.8 | 76.7 |
| Rancho Cucamonga, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 112 1,474 52 | 88 1,232 46 | 12 131 4 | 1 6 - | 11 105 2 | 41 446 26 | 794 | 128 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.3 83.3 2.9 | 5.9 83.3 3.1 | * 81.4 * | * * | * 86.1 * | 7.1 77.7 4.5 | 86.8 | 81.0 |
| Reno, Nevada | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 245 2,522 103 | 213 2,244 80 | 12 60 6 | 5 57 6 | 15 161 11 | 54 623 31 | 1,628 | 59 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.1 84.8 3.5 | 8.0 85.6 3.1 | 75.0 * | * 79.2 * | * 80.1 * | 6.5 78.8 3.9 | 8.6 88.5 2.7 | * 74.7 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|---------------------------------|---------------------|-------------------------|---------------------|
| Area, birthweight and prenatal care | | | Ra | ace | 1 | Hi | spanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Richmond city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 412 2,256 177 | 81 803 24 | 328 1,417 152 | - 4 - | 3 32 1 | 11 35 4 | 71 772 21 | 1,414 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 14.0 77.2 6.1 | 9.0 89.1 2.7 | 16.5 71.6 7.7 | * * | * 82.1 * | 64.8 * | 8.3 90.7 2.5 | 71.6 |
| Riverside, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 358 4,537 350 | 288 3,978 305 | 50 334 28 | 3 33 2 | 17 192 15 | 155 2,163 211 | 139 1,824 85 | 320 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.1 77.3 6.0 | 5.6 77.6 6.0 | 11.0 74.1 6.2 | * 68.8 * | * 79.3 * | 5.2 73.0 7.1 | 6.4 84.2 3.9 | 74.2 |
| Rochester, New York | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 429 2,870 254 | 176 1,665 105 | 247 1,134 143 | - 6 1 | 6 65 5 | 43 374 41 | 122 1,093 59 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.5 76.5 6.8 | 8.1 81.9 5.2 | 13.7 69.9 8.8 | * * | * 75.6 * | 7.8 73.5 8.1 | 8.9 84.7 4.6 | 14.2 69.1 9.3 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|---------------------|---------------------------|-----------------------------|---------------------|-------------------------|-------------------------|
| Area hirthweight and avanatal gave | | | Ra | ace | I | Hi | ispanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Rockford, Illinois | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 230 1,883 128 | 149 1,535 82 | 75 304 44 | - 1 - | 6 43 2 | 17 188 19 | 132 1,347 63 | 75 304 44 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.7 71.7 4.9 | 7.3 75.4 4.0 | 14.0 57.6 8.3 | * * | 74.1 * | * 57.8 * | 7.7 78.7 3.7 | 14.0 57.6 8.3 |
| Sacramento, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 753 7,484 661 | 375 4,979 424 | 235 1,214 132 | 8 70 15 | 135 1,221 90 | 141 1,680 211 | 246 3,370 221 | 230 1,171 128 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.0 70.1 6.2 | 5.4 72.5 6.2 | 13.1 68.4 7.4 | * 58.8 * | 6.9 63.7 4.7 | 5.3 63.9 8.0 | 5.6 77.4 5.1 | 13.3 68.4 7.5 |
| Salem, Oregon | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 136 1,949 104 | 119 1,805 100 | 5 24 - | 5 35 2 | 7 85 2 | 32 338 57 | 87 1,468 43 | 5 21 - |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.5 78.9 4.2 | 5.2 78.9 4.4 | * 77.4 * | * 74.5 * | * 81.7 * | 5.5 58.9 9.9 | 5.1 85.5 2.5 | * 77.8 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|------------------------|------------------------|---------------------|-------------------------------|------------------------------|------------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | ispanic Origin | (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Salinas, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 161 2,347 191 | 146 2,190 182 | 4 29 3 | 1 5 - | 10 123 6 | 116 1,632 171 | 31 569 12 | 4 27 3 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 4.9 71.2 5.8 | 4.7 70.8 5.9 | 76.3 * | * * | 78.3 * | 4.7 66.9 7.0 | 4.6 84.8 * | 77.1 * |
| Salt Lake City, Utah | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 467 4,976 288 | 415 4,598 229 | 16 64 12 | 12 54 13 | 24 260 34 | 100 597 98 | 314 4,015 132 | 16 50 11 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 79.5 4.6 | 7.3 81.2 4.0 | * 54.7 * | * 50.5 * | 6.4 69.5 9.1 | 9.6 58.0 9.5 | 6.7 86.1 2.8 | * 55.6 * |
| San Antonio, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,589 18,066 560 | 1,391 16,612 506 | 161 1,084 49 | 2 34 - | 35 336 5 | 1,042 11,422 433 | 348 5,219 73 | 160 1,060 48 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 84.7 2.6 | 7.1 85.1 2.6 | 11.5 78.7 3.6 | 82.9 * | 9.2 88.9 * | 7.5 82.4 3.1 | 6.1 91.4 1.3 | 11.7 78.8 3.6 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|------------------------|----------------------|---------------------|---------------------------|---------------------------------|---------------------|-------------------------|-------------------------|
| | | | R | ace | | Hi | Ispanic Origin | 1 (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| San Bernardino, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 350 3,027 339 | 240 2,382 276 | 91 509 48 | 21 2 | 19 115 13 | 150 1,625 211 | 91 773 66 | 90 493 47 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.7 68.5 7.7 | 6.6 68.0 7.9 | 12.4 70.6 6.7 | * 63.6 * | * 69.7 * | 5.8 65.6 8.5 | 8.6 74.0 6.3 | 12.7 70.4 6.7 |
| San Diego, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 1,229 15,690 975 | 786 11,791 790 | 222 1,503 93 | 9 69 5 | 212 2,327 87 | 387 5,537 671 | 399 6,315 113 | 216 1,461 91 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.3 80.6 5.0 | 5.3 80.6 5.4 | 11.2 78.0 4.8 | 73.4 * | 7.5 83.2 3.1 | 4.9 71.1 8.6 | 5.7 91.1 1.6 | 11.3 78.1 4.9 |
| San Francisco, coext. with San Fran. c | ity, Califor | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 563 7,214 213 | 238 3,871 111 | 134 681 42 | 1 13 2 | 190 2,649 58 | 106 1,439 75 | 131 2,427 37 | 133 675 41 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.7 87.0 2.6 | 5.3 87.2 2.5 | 14.9 77.8 4.8 | * * | 6.4 89.4 2.0 | 5.6 77.2 4.0 | 5.1 94.2 1.4 | 15.0 78.1 4.7 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|----------------------|---------------------|---------------------|-------------------------------|-----------------------------|---------------------|-------------------------|-----------------------------------|
| Area, birthweight and prenatal care | ŀ | | Ra | ace | I | Hi | ispanic Origin | c Non-Hispani black 61 483 20 |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | |
| | | | | | Percent | | | |
| San Jose, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 965 13,222 575 | 626 9,408 423 | 62 494 20 | 6 54 2 | 271 3,266 130 | 396 5,355 358 | 230 4,073 67 | 61 483 20 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.9 83.5 3.6 | 5.5 83.7 3.8 | 10.4 84.2 3.4 | 73.0 * | 6.6 82.9 3.3 | 5.6 78.4 5.2 | 5.1 91.5 1.5 | 10.5 84.4 3.5 |
| Santa Ana, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 488 7,088 397 | 448 6,546 377 | 6 62 3 | - 8 1 | 34 472 16 | 411 5,982 366 | 37 572 9 | 6 59 3 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.3 77.9 4.4 | 5.3 77.5 4.5 | * 75.6 * | * * | 5.9 83.2 * | 5.2 76.7 4.7 | | * 74.7 * |
| Santa Clarita, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9 193 2 | 7 169 2 | 1 6 - | - - - | 1 18 - | - 35 1 | 7 134 1 | 1 6 - |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | * 91.5 * | * 90.9 * | * * | * * | * * * | * 77.8 * | * 95.0 * | * * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | 2 8 1 54 4 2 2 0 93.1 * 6 208 9 1,066 6 75 3 0 68.6 5 4.8 |
|---|---------------------|--------------------|---------------------|-------------------------------|---------------------------------|--------------------|-------------------------|---|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | |
| | | | | | Percent | | | |
| Santa Rosa, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 128 2,081 43 | 107 1,895 34 | 8 57 2 | 3 30 3 | 10 99 4 | 35 591 20 | 72 1,301 14 | 54 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.3 89.2 1.8 | 4.9 89.8 1.6 | 93.4 * | 62.5 * | * 86.1 * | 4.6 82.8 2.8 | 5.0 93.0 * | 93.1 |
| Savannah, Georgia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 301 2,002 117 | 87 899 41 | 208 1,068 75 | - 3 - | 6 32 1 | 1 31 5 | 86 869 36 | 1,066 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.2 75.2 4.4 | 8.1 84.6 3.9 | 13.2 68.6 4.8 | * * | * 80.0 * | 72.1 * | 8.3 85.0 3.5 | 68.6 |
| Scottsdale, Arizona | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 119 1,872 39 | 114 1,784 36 | 2 18 - | 2 28 2 | 1 42 1 | 11 201 23 | 103 1,586 14 | 18 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.8 91.1 1.9 | 5.8 91.3 1.8 | * * | * 77.8 * | 91.3 * | * 72.0 8.2 | 6.1 94.5 * | * * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | 89 467 61 11.7 74.2 9.7 305 1,281 179 16.3 68.3 9.5 |
|---|---------------------|---------------------|---------------------|-------------------------------|-----------------------------|--------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | ispanic Origin | |
| Area, Dirthweight and prenatal Care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | |
| | | | | | Percent | | | |
| Seattle, Washington | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 433 4,983 308 | 255 3,608 149 | 94 502 64 | 8 74 20 | 76 799 75 | 21 259 35 | 217 3,169 105 | 467 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.2 84.8 5.2 | 5.4 88.9 3.7 | 11.2 74.9 9.6 | * 64.9 17.5 | 5.8 77.3 7.3 | 4.7 73.4 9.9 | 5.6 90.3 3.0 | 74.2 |
| Shreveport, Louisiana | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 401 2,471 202 | 90 1,155 19 | 305 1,281 179 | 2 4 - | 4 31 4 | 2 24 1 | 88 1,132 18 | 1,281 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 12.6 77.6 6.3 | 7.1 91.2 * | 16.3 68.3 9.5 | * * | * 81.6 * | * 77.4 * | 7.1 91.4 * | 68.3 |
| Simi Valley, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 83 1,520 15 | 80 1,400 13 | - 11 - | - 6 - | 3 103 2 | 26 378 6 | 54 1,007 7 | |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.0 92.5 * | 5.3 92.7 * | * * | * * | * 91.2 * | 6.0 87.3 * | 5.1 94.7 * | * * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|--------------------|---------------------|-------------------------------|------------------------------|-----------------|-------------------------|--------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | H | ispanic Origin | . (1) |
| Area, Dirthweight and prematar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black |
| | | | | | Percent | | | |
| Sioux Falls, South Dakota | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 106 1,535 34 | 95 1,457 24 | 5 28 3 | 2 25 5 | 4 25 2 | 6 24 2 | 88 1,432 22 | 28 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.9 86.1 1.9 | 5.7 87.6 1.4 | * 63.6 * | * 58.1 * | * 75.8 * | 66.7 * | 5.4 88.0 1.4 | 63.6 |
| South Bend, Indiana | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 180 1,558 103 | 99 1,182 54 | 81 343 48 | - 2 - | 31 1 | 16 153 18 | 83 1,028 36 | 342 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.8 76.7 5.1 | 6.8 81.7 3.7 | 14.6 62.7 8.8 | * * * | * 86.1 * | 68.6 * | 6.8 84.1 2.9 | 62.8 |
| Spokane, Washington | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 208 2,697 91 | 190 2,497 82 | 6 58 3 | 6 73 3 | 6 69 3 | 4 56 3 | 183 2,436 78 | 57 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.4 87.8 3.0 | 6.3 88.3 2.9 | * 87.9 * | * 79.3 * | * 80.2 * | * 78.9 * | 6.3 88.5 2.8 | * 87.7 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | d Hispanic origi | n of mother | | |
|---|---------------------|---------------------|----------------------|-------------------------------|-----------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | ispanic Origin | (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanio black |
| | | | | | Percent | | | |
| Springfield, Illinois | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 162 1,479 72 | 113 1,212 38 | 48 245 34 | - 2 - | 1 20 - | 1 17 2 | 111 1,195 36 | 48 244 34 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.1 83.7 4.1 | 8.1 87.8 2.8 | 12.8 67.3 9.3 | * * | * 95.2 * | * * | 8.1 88.0 2.7 | 12.8 67.2 9.4 |
| Springfield, Massachusetts | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 194 1,525 166 | 132 1,194 108 | 59 294 51 | - 1 - | 3 36 7 | 66 451 60 | 67 744 52 | 58 293 47 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.6 67.7 7.4 | 7.8 71.0 6.4 | 11.5 58.0 10.1 | * * | * 58.1 * | 8.8 60.4 8.0 | 7.0 78.4 5.5 | 11.6 59.4 9.5 |
| Springfield, Missouri | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 143 1,952 59 | 134 1,858 53 | 6 48 4 | - 12 - | 3 34 2 | 1 29 4 | 133 1,828 49 | 6 48 4 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.3 86.2 2.6 | 6.2 86.7 2.5 | * 76.2 * | * * | 79.1 * | 64.4 * | 6.3 87.3 2.3 | 76.2 * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|-----------------------|---------------------|----------------------|-------------------------------|------------------------------|--------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | R | ace | I | Hi | ispanic Origin | 11.6 12.3 17. 2,349 19. 321 10. 7. 9.1 11.6 12.8 11.6 11.6 11.8 |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | |
| | | | | | Percent | | | |
| St. Louis city, Missouri | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 652 4,117 379 | 125 1,664 49 | 514 2,354 321 | - 5 - | 13 94 9 | 2 62 - | 123 1,607 49 | 514 2,349 321 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.1 73.9 6.8 | 6.4 87.3 2.6 | 13.7 66.6 9.1 | * * * | * 74.6 * | * 82.7 * | 6.6 87.5 2.7 | 13.7 66.5 9.1 |
| St. Paul, Minnesota | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 316 2,966 336 | 186 2,208 140 | 78 386 71 | 5 35 8 | 47 337 117 | 21 230 27 | 164 1,924 108 | 77 371 70 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.9 69.6 7.9 | 6.3 79.4 5.0 | 11.3 62.9 11.6 | * 59.3 * | 5.3 41.8 14.5 | 5.3 62.0 7.3 | 6.7 81.9 4.6 | 11.6 62.8 11.8 |
| St. Petersburg, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 332 2,749 140 | 165 1,932 71 | 151 720 65 | 1 4 1 | 15 93 3 | 7 104 4 | 158 1,830 67 | 151 717 65 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.7 80.3 4.1 | 7.3 85.3 3.1 | 14.6 69.8 6.3 | * * | * 75.6 * | * 80.0 * | 7.4 85.7 3.1 | 14.7 69.7 6.3 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|--------------------|---------------------|-------------------------------|-----------------------------|--------------------|-------------------------|-------------------------------------|
| Area, birthweight and prenatal care | ŀ | | R | ace | I | Hi | ispanic Origin | n (1) |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | 10.4 67.6 11.6 79.6 8.3 |
| | | | | | Percent | | | |
| Stamford, Connecticut | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 137 1,421 79 | 92 1,125 46 | 37 227 25 | - 1 1 | 8 68 7 | 22 221 23 | 64 848 23 | 221 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 81.8 4.5 | 6.6 85.8 3.5 | 10.4 67.0 7.4 | * * | * 80.0 * | 6.2 64.8 6.7 | 6.7 93.4 2.5 | 67.0 |
| Staten Island borough, Richmond county | New York | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 443 4,972 156 | 323 4,108 89 | 95 609 60 | - 7 - | 25 248 7 | 70 626 33 | 252 3,462 53 | 578 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.5 90.1 2.8 | 6.8 91.9 2.0 | 11.4 80.1 7.9 | * * | 8.4 89.2 * | 8.8 84.6 4.5 | 6.4 93.4 1.4 | 79.6 |
| Sterling Heights, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 79 1,352 15 | 72 1,285 14 | 2 11 1 | - 2 - | 5 54 - | - 21 - | 69 1,179 13 | 11 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.1 91.5 * | 4.9 91.5 * | * * * | * * | * 94.7 * | 91.3 * | 5.2 91.5 * | * * |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | | |
|---|---------------------|---------------------|---------------------|-------------------------------|-----------------------------|---------------------|-----------------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | Ra | ace | Ι | Hi | ispanic Origin | .n (1) | |
| Area, Dirthweight and prenatal Care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanic black | |
| | | | | | Percent | | | | |
| Stockton, California | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 388 3,560 307 | 225 2,421 221 | 76 398 37 | 6 26 6 | 81 715 43 | 141 1,378 163 | 1,098 | 75 383 36 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 69.9 6.0 | 6.4 71.3 6.5 | 12.8 69.5 6.5 | 63.4 * | 7.2 65.7 4.0 | 6.5 65.9 7.8 | 6.3 79.8 4.2 | 13.1 69.5 6.5 | |
| Sunnyvale, California | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 117 1,826 47 | 65 1,130 30 | 2 60 2 | 2 5 - | 48 631 15 | 22 384 13 | 749 | 2 56 2 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.4 87.1 2.2 | 4.9 86.9 2.3 | * 83.3 * | * * | 6.5 87.8 * | 4.5 81.4 * | | * 83.6 * | |
| Syracuse, New York | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 245 1,786 132 | 109 1,139 63 | 124 561 63 | 9 26 - | 3 60 6 | 13 99 12 | 1,027 | 123 540 62 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 10.1 74.7 5.5 | 7.5 79.7 4.4 | 14.7 67.2 7.5 | * 57.8 * | * 73.2 * | * 68.3 * | 7.3 80.7 3.8 | 15.2 67.3 7.7 | |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|--------------------|---------------------|-------------------------------|-----------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | Hi | Hispanic Origin | |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Tacoma, Washington | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 224 2,368 99 | 147 1,775 50 | 50 302 22 | 8 65 9 | 19 226 18 | 10 167 10 | 126 1,565 32 | 47 294 20 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.1 80.7 3.4 | 6.4 83.1 2.3 | 11.8 78.9 5.7 | 64.4 * | * 72.2 * | 69.9 * | 6.4 85.1 1.7 | 11.8 79.7 5.4 |
| Tallahassee, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 151 1,697 42 | 56 995 9 | 89 634 33 | - 1 - | 6 67 - | 5 47 1 | 51 948 8 | 89 633 33 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.9 89.1 2.2 | 5.3 94.0 * | 11.4 81.6 4.2 | * * * | 97.1 * | * 87.0 * | 5.1 94.4 * | 11.5 81.6 4.3 |
| Tampa, Florida | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 596 5,612 229 | 301 3,944 92 | 279 1,499 132 | 2 17 1 | 14 152 4 | 93 1,052 41 | 214 2,902 52 | 270 1,482 131 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.7 81.9 3.3 | 6.7 87.6 2.0 | 13.0 69.9 6.2 | * * | * 83.5 * | 7.3 83.6 3.3 | 6.5 89.0 1.6 | 12.7 69.9 6.2 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | | |
|---|-----------------------|--------------------|---------------------|-------------------------------|-----------------------------|--------------------|-------------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | Ra | ace | | Hispanic Origi | | gin (1) | |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black | |
| | | | | | Percent | | | | |
| Tempe, Arizona | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 134 1,702 90 | 109 1,502 75 | 10 49 4 | 5 45 5 | 10 106 6 | 34 371 46 | 75 1,131 29 | 10 48 4 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.4 81.2 4.3 | 5.9 81.8 4.1 | 74.2 * | 66.2 * | 84.1 | 6.2 67.9 8.4 | 5.8 87.5 2.2 | 73.8 * | |
| Thousand Oaks, California | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 78 1,264 14 | 72 1,165 12 | 1 10 - | 2 7 - | 3 82 2 | 18 289 9 | 54 876 3 | 1 10 - | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 5.7 92.4 * | 5.7 92.2 * | * * | * * * | * 94.3 * | * 83.8 * | 5.9 95.4 * | * * * | |
| Toledo, Ohio | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 419 4,265 163 | 263 3,289 76 | 152 907 81 | 1 11 4 | 3 58 2 | 24 243 16 | 238 3,051 61 | 151 897 81 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.3 84.1 3.2 | 7.0 87.7 2.0 | 12.3 73.3 6.5 | * * | * 89.2 * | 7.5 75.9 * | 6.9 88.6 1.8 | 12.4 73.5 6.6 | |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|-----------------------------|---------------------|-------------------------|-----------------------|
| Area, birthweight and prenatal care | | | Ra | ace | T | Hi | spanic Origin | n (1) |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black |
| | | | | | Percent | | | |
| Topeka, Kansas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 166 1,760 29 | 122 1,482 21 | 40 228 7 | 2 30 1 | 2 20 - | 15 156 5 | 107 1,301 16 | 40 223 7 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.4 89.4 1.5 | 7.5 91.0 1.3 | 14.2 80.9 * | * 88.2 * | * 83.3 * | * 81.3 * | 7.6 92.1 * | 14.5 80.8 * |
| Torrance, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 151 2,050 47 | 100 1,386 36 | 14 87 5 | - 13 - | 37 564 6 | 50 646 20 | 53 754 15 | 12 83 4 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.5 88.3 2.0 | 6.4 88.3 2.3 | * 77.0 * | * * | 5.9 90.5 * | 6.6 85.0 2.6 | 6.4 91.4 * | * 77.6 * |
| Tucson, Arizona | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 617 6,181 719 | 538 5,696 655 | 39 204 27 | 23 140 23 | 17 141 14 | 282 2,781 502 | 269 2,970 161 | 37 190 25 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.1 70.9 8.3 | 6.7 71.1 8.2 | 12.8 66.7 8.8 | 10.7 65.4 10.7 | * 75.8 * | 6.5 63.8 11.5 | 7.2 79.6 4.3 | 13.0 66.2 8.7 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|---------------------------|------------------------------|--------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | Ra | ace | | Hi | spanic Origin | (1) |
| Area, Dirthweight and prehatar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanio black |
| | | | | | Percent | | | |
| Tulsa, Oklahoma | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 486 4,404 355 | 302 3,348 203 | 154 729 100 | 18 237 45 | 12 90 7 | 31 270 40 | 272 3,084 163 | 153 727 100 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.2 75.5 6.1 | 7.0 79.1 4.8 | 13.4 64.5 8.8 | * 67.7 12.9 | * 76.9 * | 7.0 61.5 9.1 | 7.0 81.1 4.3 | 13.4 64.6 8.9 |
| Vallejo, California | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 135 1,168 148 | 41 521 65 | 47 285 50 | 1 3 - | 46 359 33 | 17 203 20 | 26 329 45 | 46 280 50 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.4 66.1 8.4 | 5.1 66.3 8.3 | 9.8 61.7 10.8 | * * | 8.9 70.9 6.5 | * 62.7 6.2 | 5.3 68.3 9.3 | 9.8 61.7 11.0 |
| Virginia Beach city, Virginia | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 474 5,460 173 | 303 4,206 100 | 149 991 65 | - 11 - | 22 252 8 | 25 236 4 | 278 3,983 97 | 149 981 64 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.3 84.6 2.7 | 6.3 87.8 2.1 | 11.2 74.4 4.9 | * * | 7.0 80.3 * | 8.7 82.2 * | 6.1 88.1 2.1 | 11.3 74.4 4.9 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|---------------------|-------------------------------|------------------------------|-----------------------------|-------------------------|---------------------|
| Area, birthweight and prenatal care | | | Ra | ace | I | ні | spanic Origin | (1) |
| Area, Diffilweight and prematar care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Non-Hispanic White | Non-Hispanic black | |
| | | | | | Percent | | | |
| Waco, Texas | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 186 1,513 201 | 138 1,200 131 | 48 292 70 | - - - | - 21 - | 54 376 73 | 85 826 59 | 47 289 69 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.0 65.3 8.7 | 7.8 67.7 7.4 | 9.2 55.8 13.4 | * * * | * 95.5 * | 7.8 54.6 10.6 | 7.8 76.0 5.4 | 9.1 55.9 13.3 |
| Warren, Michigan | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 117 1,576 34 | 114 1,506 31 | 2 24 2 | - 4 - | 1 42 1 | 1 26 - | 103 1,393 27 | 2 22 2 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 6.4 88.6 1.9 | 6.5 89.1 1.8 | * 77.4 * | * * | * 79.2 * | 96.3 * | 6.4 89.2 1.7 | * 75.9 * |
| Waterbury, Connecticut | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 160 1,136 88 | 115 910 65 | 44 206 23 | - 3 - | 1 17 - | 40 233 39 | 75 672 28 | 35 198 21 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 9.4 72.2 5.6 | 8.7 73.8 5.3 | 12.8 65.2 7.3 | * * | * * | 8.9 58.3 9.8 | 8.5 80.5 3.4 | 11.0 66.9 7.1 |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | | |
|---|-----------------------|---------------------|---------------------|---------------------------|------------------------------|--------------------|-------------------------|-------------------------|--|
| Area, birthweight and prenatal care | | | Ra | ace | | H: | ispanic Origin | in (1) | |
| Area, Dirthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanio white | Non-Hispanic black | |
| | | | | | Percent | | | | |
| Wichita, Kansas | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 425 4,965 163 | 291 4,115 108 | 117 590 45 | 3 28 5 | 14 232 5 | 39 360 34 | 252 3,757 71 | 117 590 45 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 7.2 84.1 2.8 | 6.1 86.0 2.3 | 14.7 74.0 5.6 | 68.3 * | * 82.9 * | 7.6 70.7 6.7 | 5.9 87.9 1.7 | 14.7 74.0 5.6 | |
| Winston-Salem, North Carolina | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 301 2,271 76 | 122 1,352 32 | 177 888 44 | - 3 - | 2 28 - | 21 244 19 | 101 1,108 13 | 177 888 44 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 11.5 87.4 2.9 | 8.1 90.7 2.1 | 16.4 82.6 4.1 | * * * | 90.3 * | 6.8 80.0 * | 8.5 93.5 * | 16.4 82.6 4.1 | |
| Worcester, Massachusetts | | | | | | | | | |
| Number | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 187 1,857 79 | 154 1,567 57 | 20 156 16 | - 7 - | 13 127 6 | 43 365 15 | 115 1,226 41 | 16 132 15 | |
| Percent | | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.3 82.8 3.5 | 8.3 84.3 3.1 | 9.3 72.6 * | * * | * 78.4 * | 9.4 80.2 * | 8.0 85.8 2.9 | * 70.2 * | |

Table 1-25. Number and Percent of Births of Low Birthweight, Number and Percent of Women Beginning Prenatal Care in First Trimester, and Number and Percent Receiving Late or No Care by Race and Hispanic Origin of Mother for Cities of 100,000 or More: United States, 1996

| | | | | Race and | l Hispanic origi | n of mother | | |
|---|---------------------|---------------------|----------------------|-------------------------------|----------------------------------|---------------------|-------------------------|-------------------------|
| Area, birthweight and prenatal care | | | R | ace | | Ні | Hispanic Origin | |
| Area, birthweight and prenatal care | All races | White | Black | American Indian (2) | Asian/Pacific Islander | Hispanic | Non-Hispanic white | Non-Hispanio black |
| | | | | | Percent | | | |
| Yonkers, New York | | | | | | | | |
| Number | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 244 1,832 408 | 149 1,392 282 | 78 331 103 | 1 1 1 | 16 108 22 | 80 510 174 | 66 869 111 | 70 306 99 |
| Percent | | | | | | | | |
| Low birthweight Care beginning in first trimester Late or no care | 8.6 68.7 15.3 | 7.3 71.1 14.4 | 13.3 61.5 19.1 | * * * | * 63.9 13.0 | 8.8 58.7 20.0 | 5.8 79.8 10.2 | 12.9 61.4 19.9 |

⁻ Quantity zero.

^{*} Figure does not meet standards of reliability or precision; based on fewer than 20 births in the numerator.

⁽¹⁾ Data not shown for non-Hispanic births of other races and births with origin not stated.

⁽²⁾ Includes Aleuts and Eskimos.