

Table V-1
CERVIX UTERI CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1973-1999
WITH UP TO THREE JOINPOINTS BY RACE AND AGE

	<u>All Races</u>		<u>Whites</u>		<u>Blacks</u>	
	<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>
<u>SEER Cancer Incidence^a</u>						
Age - All Ages	1973-81	-4.6★	1973-82	-5.0★	1973-93	-4.3★
	1981-99	-1.3★	1982-90	0.4	1993-99	-0.5
			1990-99	-2.5★		
Age - Under 50	1973-78	-6.4★	1973-81	-4.9★	1973-87	-5.2★
	1978-99	-0.8★	1981-89	1.5	1987-99	-1.3
			1989-99	-1.5★		
Age - 50 and Over	1973-82	-4.7★	1973-83	-5.4★	1973-99	-4.0★
	1982-99	-2.0★	1983-92	-0.7		
			1992-99	-4.3★		
<u>U.S. Cancer Mortality^b</u>						
Age - All Ages	1973-82	-4.6★	1973-82	-4.7★	1973-79	-5.4★
	1982-97	-1.6★	1982-99	-1.5★	1979-99	-2.9★
	1997-99	-5.4				
Age - Under 50	1973-79	-5.9★	1973-79	-5.3★	1973-76	-10.7
	1979-97	-0.8★	1979-99	-0.5★	1976-99	-3.0★
	1997-99	-6.3				
Age - 50 and Over	1973-82	-4.8★	1973-83	-4.9★	1973-99	-3.3★
	1982-99	-2.2★	1983-99	-2.0★		

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

★ The APC is significantly different from zero (p<.05).

Table V-2
CERVIX UTERI CANCER (Invasive)

SEER INCIDENCES AND U.S. DEATH^â RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence [§]			US Death ^â		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-SPECIFIC RATES, 1995-99</u>						
AGE AT DIAGNOSIS/DEATH:						
0-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	1.7	1.8	-	0.2	0.2	0.4
25-29	7.3	7.8	5.9	0.8	0.8	1.1
30-34	12.0	12.0	12.7	1.8	1.8	2.7
35-39	14.0	13.8	12.7	2.9	2.7	4.7
40-44	14.9	13.4	20.7	4.1	3.7	7.0
45-49	16.5	14.8	24.6	5.1	4.6	8.5
50-54	14.7	12.8	18.0	5.8	5.3	10.2
55-59	14.3	11.7	25.6	6.1	5.3	11.8
60-64	15.1	12.9	24.7	6.9	5.9	14.3
65-69	14.3	12.1	28.1	7.2	6.2	15.8
70-74	13.2	11.6	27.1	7.4	6.3	19.0
75-79	14.9	12.1	39.0	8.2	7.1	19.8
80-84	13.1	9.3	44.9	9.2	7.9	23.6
85+	13.4	10.9	41.7	11.9	10.0	33.4
<u>AGE-ADJUSTED RATES, 1995-99</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	9.0	8.1	13.6	3.1	2.8	6.2
Under 65	8.2	7.6	10.6	2.4	2.1	4.2
65 and over	13.9	11.4	34.2	8.3	7.1	20.7
All ages (world std.) [⊕]	7.1	6.5	10.1	2.3	2.0	4.4

§ SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.
 â NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.
 ⊕ Rates are per 100,000 and are age-adjusted to the world standard population.
 - Statistic not shown. Rate based on less than 25 cases for the time interval.

Table V-3
CERVIX UTERI CANCER (Invasive)

SEER INCIDENCES AND U.S. DEATH RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
SEER INCIDENCE RATES									
YEAR OF DIAGNOSIS:									
1973	17.2	11.7	31.6	15.5	10.3	29.2	36.7	24.1	69.7
1974	15.2	10.1	28.6	13.9	9.3	25.9	30.0	15.5	68.0
1975	14.8	9.7	28.2	13.2	8.8	24.8	32.8	17.5	73.0
1976	14.2	9.5	26.7	12.9	8.7	23.7	29.4	16.0	64.4
1977	13.0	8.4	25.1	11.6	7.5	22.6	28.2	15.9	60.4
1978	12.5	8.3	23.6	11.3	7.7	20.9	24.1	12.4	55.0
1979	12.7	8.7	23.4	10.9	7.5	19.8	28.8	16.0	62.4
1980	12.2	8.0	23.1	10.9	7.2	20.6	23.4	13.4	49.6
1981	10.8	7.3	19.7	9.5	6.6	17.2	23.9	13.6	50.8
1982	10.6	7.1	19.8	9.4	6.7	16.5	21.9	10.2	52.5
1983	10.5	7.3	18.8	9.6	6.8	17.0	18.8	11.9	37.0
1984	11.0	8.0	18.9	10.0	7.4	16.6	21.0	12.4	43.3
1985	10.2	7.1	18.5	9.1	6.7	15.4	18.9	9.4	43.8
1986	10.8	7.9	18.6	9.7	7.3	16.0	18.4	10.6	38.9
1987	10.0	7.1	17.6	9.0	6.9	14.5	18.8	8.7	45.1
1988	10.6	7.9	17.7	9.5	7.4	15.1	18.9	10.4	40.9
1989	10.7	7.9	18.0	9.9	7.8	15.7	15.6	8.0	35.5
1990	10.7	8.0	17.6	10.0	7.7	15.8	16.7	9.6	35.2
1991	10.0	7.0	17.9	9.2	6.8	15.6	16.1	8.9	35.0
1992	10.0	7.5	16.5	9.4	7.2	15.2	14.0	9.9	24.6
1993	9.7	7.1	16.4	9.1	7.0	14.7	13.7	8.8	26.7
1994	9.5	6.9	16.2	8.7	6.7	13.8	14.1	6.9	32.9
1995	8.8	6.7	14.4	7.9	6.4	11.8	13.6	7.8	28.9
1996	9.6	7.4	15.5	8.7	7.0	13.0	14.1	8.1	29.9
1997	9.2	7.0	14.7	8.3	6.7	12.2	13.8	8.2	28.6
1998	9.2	7.1	14.5	8.3	6.7	12.5	13.3	9.1	24.2
1999	8.0	6.3	12.5	7.4	6.2	10.5	13.3	8.0	27.4
1995-99	9.0	6.9	14.3	8.1	6.6	12.0	13.6	8.2	27.7
U.S. DEATH RATES									
YEAR OF DEATH:									
1973	6.2	2.7	15.5	5.3	2.2	13.4	15.6	6.8	38.7
1974	6.0	2.6	15.1	5.2	2.1	13.3	14.1	6.3	34.4
1975	5.6	2.4	13.9	4.7	1.9	12.1	13.3	5.7	33.3
1976	5.4	2.2	13.9	4.7	1.9	12.1	12.9	4.9	33.7
1977	5.0	2.0	12.7	4.2	1.6	11.0	12.2	5.0	31.1
1978	4.8	2.1	12.0	4.2	1.8	10.4	11.6	4.8	29.6
1979	4.6	1.9	11.6	3.9	1.6	10.2	10.6	4.5	26.7
1980	4.5	2.0	11.0	3.8	1.6	9.5	10.8	4.6	26.9
1981	4.3	1.8	10.9	3.7	1.6	9.4	9.9	3.7	26.0
1982	4.1	1.8	10.2	3.5	1.5	8.7	9.8	3.6	25.9
1983	4.0	1.8	9.9	3.4	1.5	8.4	9.8	3.9	25.2
1984	3.9	1.8	9.5	3.4	1.5	8.1	9.3	3.9	23.3
1985	3.8	1.7	9.4	3.3	1.5	7.9	9.0	3.4	23.7
1986	3.8	1.8	9.2	3.2	1.5	7.8	9.2	4.0	22.9
1987	3.6	1.7	8.6	3.1	1.5	7.4	8.2	3.6	20.3
1988	3.6	1.7	8.6	3.1	1.5	7.2	8.2	3.0	21.7
1989	3.6	1.7	8.5	3.0	1.5	7.1	8.6	3.3	22.6
1990	3.7	1.8	8.6	3.2	1.6	7.4	7.8	3.4	19.2
1991	3.5	1.7	8.2	3.0	1.5	7.0	7.7	3.2	19.4
1992	3.5	1.7	8.2	3.0	1.5	6.9	8.1	3.4	20.4
1993	3.4	1.7	8.0	3.0	1.5	6.8	7.7	3.2	19.7
1994	3.4	1.7	7.9	3.0	1.5	7.0	6.8	2.9	16.9
1995	3.3	1.6	7.5	2.8	1.5	6.4	7.0	3.0	17.7
1996	3.2	1.6	7.6	2.9	1.5	6.6	6.3	2.5	16.0
1997	3.2	1.6	7.2	2.8	1.5	6.3	6.5	2.9	16.2
1998	3.0	1.6	6.7	2.7	1.5	5.8	6.0	2.6	14.9
1999	2.9	1.4	6.8	2.6	1.3	5.9	5.5	1.9	14.9
1995-99	3.1	1.6	7.2	2.8	1.4	6.2	6.2	2.6	15.9

SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.
 NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.
 Statistic not shown. Rate based on less than 25 cases for the time interval.

Table V-4
CERVIX UTERI CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>5-YR RELATIVE SURVIVAL RATES</u>									
YEAR OF DIAGNOSIS:									
1960-63 ^a	-	-	-	58	-	-	47	-	-
1970-73 ^a	-	-	-	64	-	-	61	-	-
1974-76 ^b	69.1	81.4	58.1	69.8	82.3	58.7	63.8	75.3	53.8
1977-79 ^b	68.4	78.0	59.7	69.5	79.8	60.6	62.2	71.6	52.7
1980-82 ^b	67.7	79.5	56.1	68.2	80.8	55.8	61.3	72.1	51.1
1983-85 ^b	68.8	78.4	57.9	70.6	80.4	58.8	60.4	69.8	50.5
1986-88 ^b	68.9	78.6	56.7	71.7	81.4	58.4	55.5	63.7	47.7
1989-91 ^b	70.8	78.1	60.7	72.3	80.6	60.1	62.6	66.9	57.5
1992-98 ^b	70.7 \bar{U}	79.6	57.9	72.1 \bar{U}	81.9	57.2	59.9	63.1 \bar{U}	55.4
1992-98 ^b									
ALL STAGES	70.7	79.6	57.9	72.1	81.9	57.2	59.9	63.1	55.4
LOCALIZED	92.2	94.5	87.0	92.6	95.5	85.0	87.1	86.4	89.3
REGIONAL	50.6	54.4	47.6	51.3	56.3	47.2	41.0	37.7	44.1
DISTANT	15.2	21.8	11.0	16.5	24.9	11.5	7.5	12.6 \uparrow	3.0
UNSTAGED	52.0	67.6	36.4	53.3	71.3	36.3	49.8 \uparrow	55.7 \uparrow	40.1 \uparrow
<u>STAGE DISTRIBUTION (%) 1992-98^b</u>									
ALL STAGES									
Number of cases	7,594	4,376	3,218	5,663	3,367	2,296	1,097	606	491
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	54	66	39	56	67	40	46	54	36
REGIONAL	32	24	42	30	23	41	35	31	41
DISTANT	8	5	11	8	5	11	9	6	11
UNSTAGED	7	5	9	6	5	8	10	9	12
<u>5-YR RELATIVE SURVIVAL RATES, 1992-98^b</u>									
AGE AT DIAGNOSIS:									
<45	82.4	-	-	84.4	-	-	67.8	-	-
45-54	66.8	-	-	68.2	-	-	49.7	-	-
55-64	63.8	-	-	62.7	-	-	59.9 \uparrow	-	-
65-74	56.9	-	-	57.2	-	-	54.1 \uparrow	-	-
75+	41.4	-	-	38.7	-	-	50.2 \uparrow	-	-
Under 65	75.2	-	-	76.9	-	-	61.9	-	-
65 and over	50.9	-	-	50.1	-	-	52.7	-	-

^a

^b

Rates are based on End Results data from a series of hospital registries and one population-based registry. Rates are from the SEER 9 areas. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients into 1999.

The difference in rates between 1974-76 and 1992-98 is statistically significant ($p < .05$).

The standard error of the survival rate is between 5 and 10 percentage points.

The standard error of the survival rate is greater than 10 percentage points.

Statistic could not be calculated.

\bar{U}

\uparrow

\uparrow

-

Table V-5
CERVIX UTERI CANCER (Invasive)

INCIDENCE, DEATH, AND SURVIVAL RATES*

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1975	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Incidence																					
SEER ^a	14.8	12.2	10.8	10.6	10.5	11.0	10.2	10.8	10.0	10.6	10.7	10.7	10.0	10.0	9.7	9.5	8.8	9.6	9.2	9.2	8.0
Mortality																					
U.S. ^b	5.6	4.5	4.3	4.1	4.0	3.9	3.8	3.8	3.6	3.6	3.6	3.7	3.5	3.5	3.4	3.4	3.3	3.2	3.2	3.0	2.9
SEER ^a	4.6	3.5	3.7	3.5	3.4	3.3	3.3	3.1	2.9	3.1	3.3	2.8	2.9	2.8	2.6	3.0	2.4	2.5	2.2	2.5	2.0
Relative Survival Rates (SEER ^a)																					
1-year	86.5	86.9	86.6	87.3	89.2	88.4	87.0	86.1	87.5	87.5	89.2	88.3	88.9	87.5	89.3	87.9	89.7	88.5	89.0	88.3	
2-year	78.4	77.3	77.6	75.7	78.7	79.0	76.0	76.9	77.8	79.1	80.7	79.8	79.7	78.5	79.6	80.4	81.7	81.1	80.1		
3-year	73.8	73.3	73.6	70.1	74.3	74.3	71.3	71.9	72.5	75.5	75.9	75.6	74.4	74.4	75.3	76.4	78.4	76.6			
4-year	70.9	70.4	70.3	67.8	71.2	71.4	68.7	69.1	69.5	73.6	72.9	73.3	71.0	70.6	72.5	73.6	75.5				
5-year	68.9	68.8	68.0	66.2	69.9	69.6	66.9	66.5	67.4	72.6	71.1	72.2	68.9	68.2	70.6	71.9					
6-year	68.0	67.7	66.8	63.9	67.8	67.5	66.0	65.9	66.5	70.7	69.8	70.8	67.3	66.8	70.2						
7-year	67.5	65.4	64.4	63.4	66.6	66.1	64.6	64.8	65.8	70.0	68.6	70.0	66.5	65.8							
8-year	66.5	64.6	63.0	62.8	65.1	64.8	63.6	63.9	65.6	69.2	67.4	69.0	65.4								
9-year	66.5	63.7	62.8	62.6	65.1	63.6	62.7	63.9	65.3	68.7	66.4	66.9									
10-year	65.1	62.5	62.8	62.4	64.7	63.5	62.0	63.6	65.2	67.8	65.5										
11-year	64.5	62.0	62.8	62.3	63.9	62.8	61.7	63.1	64.7	67.5											
12-year	64.2	61.2	62.4	61.7	62.8	62.4	60.8	62.7	64.1												
13-year	63.1	61.2	62.3	60.4	62.6	61.7	60.2	62.7													
14-year	62.9	60.6	60.7	60.0	62.2	61.3	60.2														
15-year	62.6	60.6	59.5	59.2	61.4	60.1															
16-year	61.2	59.6	59.5	58.5	60.7																
17-year	60.4	58.9	58.5	57.0																	
18-year	60.2	57.2	57.9																		
19-year	59.9	56.6																			
20-year	59.7																				
21-year	59.4																				
22-year	58.8																				
23-year	58.1																				
24-year	57.0																				

^a Rates are from the SEER 9 areas.

^b Rates are from the NCHS public use data file.

* Incidence and death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

Survival rates are relative rates expressed as percents.

Table V-6
CERVIX UTERI CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1997-1999

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	0.82
10	0.00	0.04	0.17	0.82
20	0.04	0.17	0.34	0.82
30	0.13	0.30	0.45	0.79
40	0.17	0.32	0.47	0.66
50	0.16	0.31	0.42	0.50
60	0.16	0.28	0.34	0.36

Lifetime Risk of Being Diagnosed = 0.82%
 Lifetime Risk of Dying = 0.29%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.05	0.76
10	0.00	0.05	0.18	0.77
20	0.05	0.18	0.34	0.77
30	0.14	0.30	0.43	0.72
40	0.16	0.30	0.44	0.59
50	0.14	0.28	0.37	0.44
60	0.14	0.24	0.30	0.31

Lifetime Risk of Being Diagnosed = 0.76%
 Lifetime Risk of Dying = 0.26%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	1.03
10	0.00	0.04	0.16	1.05
20	0.04	0.16	0.37	1.05
30	0.12	0.33	0.52	1.02
40	0.22	0.41	0.60	0.92
50	0.20	0.40	0.58	0.74
60	0.22	0.42	0.52	0.59

Lifetime Risk of Being Diagnosed = 1.03%
 Lifetime Risk of Dying = 0.54%

Table V-7
CERVIX UTERI CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidenceⁱ

RACE/ETHNICITY	Rate 1992-1999	Trend 1992-1999
	Rate per 100,000 persons	EAPC (%)
	Females	Females
All Races	10.2	-2.3★
White	9.6	-2.2★
White Hispanic	18.5	-4.3★
White Non-Hispanic	8.0	-2.0★
Black	13.3	-1.9★
Asian/Pacific Islander	11.7	-3.9
Amer Ind/Alaska Nat	7.7	-
Hispanic [⊗]	17.5	-4.3★

Mortality[❖]

RACE/ETHNICITY	Rate 1992-1999	Trend 1992-1999
	Rate per 100,000 persons	EAPC (%)
	Females	Females
All Races	3.2	-2.7★
White	2.8	-2.1★
White Hispanic	4.1	-4.4★
White Non-Hispanic	2.7	-1.9★
Black	6.7	-4.9★
Asian/Pacific Islander	3.1	-1.9
Amer Ind/Alaska Nat	3.3	-7.1
Hispanic [⊗]	3.8	-4.4★

- The EAPC is the Estimated Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ⁱ Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- [⊗] Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics does not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Oklahoma, New York, and New Hampshire.
- [❖] Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- [★] The EAPC is significantly different from zero (p<.05).

Table V-8
CERVIX UTERI CANCER (Invasive)AGE-ADJUSTED SEER INCIDENCE RATES§ AND SEER DEATH RATESä
By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES§, 1995-99</u>									
REGISTRY									
11 registries	9.8	7.4	16.3	9.2	7.3	14.2	13.1	7.9	26.7
9 registries	9.0	6.9	14.3	8.1	6.6	12.0	13.6	8.2	27.7
San Francisco-Oakland	8.7	6.5	14.4	8.1	6.7	11.7	10.0	6.9	18.3
Connecticut	8.8	6.4	15.1	8.2	6.1	13.7	13.8	7.8	29.6
Metropolitan Detroit	10.0	7.5	16.8	8.3	6.5	12.9	15.5	9.9	30.3
Hawaii	10.1	7.4	17.0	8.8	6.8	14.1	-	-	-
Iowa	8.9	8.0	11.4	8.7	7.9	11.0	-	-	-
New Mexico	9.3	7.1	15.1	8.9	7.0	14.0	-	-	-
Seattle-Puget Sound	7.5	6.4	10.6	7.2	6.3	9.4	-	-	-
Utah	7.3	6.5	9.6	7.3	6.6	9.3	-	-	-
Metropolitan Atlanta	10.8	7.1	20.6	8.5	6.3	14.5	15.7	7.8	36.6
San Jose-Monterey	9.5	6.5	17.4	8.7	6.3	15.1	-	-	-
Los Angeles	12.7	8.9	22.4	13.2	9.9	21.7	11.9	7.0	24.9
<u>SEER DEATH RATESä, 1995-99</u>									
REGISTRY									
11 registries	2.6	1.2	6.3	2.4	1.2	5.6	4.7	2.1	11.5
9 registries	2.3	1.1	5.5	2.1	1.0	5.0	4.3	1.9	10.7
San Francisco-Oakland	2.0	0.9	4.7	1.7	0.8	4.0	3.9	-	-
Connecticut	2.2	1.1	4.9	2.0	1.0	4.5	4.9	-	-
Metropolitan Detroit	2.6	1.3	6.1	2.1	1.0	5.0	4.5	2.2	10.5
Hawaii	1.9	-	4.5	-	-	-	-	-	-
Iowa	2.4	1.3	5.3	2.3	1.3	5.2	-	-	-
New Mexico	2.9	1.2	7.2	2.8	1.2	7.2	-	-	-
Seattle-Puget Sound	2.3	1.1	5.6	2.3	1.2	5.2	-	-	-
Utah	1.8	1.0	4.0	1.8	1.0	3.9	-	-	-
Metropolitan Atlanta	2.8	0.9	7.8	2.0	-	5.6	5.3	1.5	15.3
San Jose-Monterey	2.7	1.1	6.9	2.4	1.1	5.8	-	-	-
Los Angeles	3.6	1.7	8.5	3.4	1.7	7.8	5.5	2.7	12.7

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.
 § SEER 9 areas. Incidence rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.
 ä NCHS public use data file. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.
 - Statistic could not be calculated or populations are inadequate.

Table V-9
CERVIX UTERI CANCER (Invasive)AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^ä RATES BY STATE, 1995-99

All Races, Females

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
TOTAL U.S.	3.1	0.02							
High Five States									
District of Columbia	5.3 ^{⊕*}	0.59	(01)	71.0					
Delaware	4.9 ^{⊕*}	0.50	(02)	58.1					
West Virginia	4.3 ^{⊕*}	0.29	(03)	38.7					
Kentucky	4.2 ^{⊕*}	0.20	(04)	35.5					
Mississippi	4.1 ^{⊕*}	0.24	(05)	32.3					
Low Five States									
Alaska	2.0 [*]	0.44	(47)	-35.5					
Minnesota	2.0 ^{⊕*}	0.13	(48)	-35.5					
Hawaii	1.9 ^{⊕*}	0.25	(49)	-38.7					
South Dakota	1.9 ^{⊕*}	0.32	(50)	-38.7					
Utah	1.8 ^{⊕*}	0.21	(51)	-41.9					
Alabama	3.3	0.17	(19)	6.5	Montana	2.4 [*]	0.32	(38)	-22.6
Alaska	2.0 [*]	0.44	(47)	-35.5	Nebraska	2.8	0.25	(32)	-9.7
Arizona	2.5 ^{⊕*}	0.15	(35)	-19.4	Nevada	3.4	0.29	(17)	9.7
Arkansas	3.9 [*]	0.24	(09)	25.8	New Hampshire	2.4 [*]	0.28	(39)	-22.6
California	3.0	0.06	(28)	-3.2	New Jersey	3.3	0.12	(20)	6.5
Colorado	2.2 ^{⊕*}	0.15	(45)	-29.0	New Mexico	2.9	0.26	(30)	-6.5
Connecticut	2.2 ^{⊕*}	0.15	(46)	-29.0	New York	3.3	0.08	(22)	6.5
Delaware	4.9 ^{⊕*}	0.50	(02)	58.1	North Carolina	3.4	0.13	(14)	9.7
District of Columbia	5.3 ^{⊕*}	0.59	(01)	71.0	North Dakota	2.9	0.42	(31)	-6.5
Florida	3.4	0.09	(15)	9.7	Ohio	3.1	0.10	(24)	0.0
Georgia	3.4	0.14	(13)	9.7	Oklahoma	3.6 [*]	0.20	(10)	16.1
Hawaii	1.9 ^{⊕*}	0.25	(49)	-38.7	Oregon	2.5 ^{⊕*}	0.17	(37)	-19.4
Idaho	2.7	0.31	(33)	-12.9	Pennsylvania	3.1	0.09	(26)	0.0
Illinois	3.4	0.10	(16)	9.7	Rhode Island	3.0	0.33	(29)	-3.2
Indiana	3.3	0.15	(18)	6.5	South Carolina	3.9 ^{⊕*}	0.20	(07)	25.8
Iowa	2.4 ^{⊕*}	0.18	(41)	-22.6	South Dakota	1.9 ^{⊕*}	0.32	(50)	-38.7
Kansas	2.4 ^{⊕*}	0.19	(40)	-22.6	Tennessee	3.9 ^{⊕*}	0.16	(08)	25.8
Kentucky	4.2 ^{⊕*}	0.20	(04)	35.5	Texas	3.5 [⊕]	0.09	(12)	12.9
Louisiana	4.1 ^{⊕*}	0.19	(06)	32.3	Utah	1.8 ^{⊕*}	0.21	(51)	-41.9
Maine	3.1	0.30	(23)	0.0	Vermont	3.6 [*]	0.48	(11)	16.1
Maryland	3.1	0.15	(25)	0.0	Virginia	3.0	0.13	(27)	-3.2
Massachusetts	2.4 ^{⊕*}	0.12	(42)	-22.6	Washington	2.3 ^{⊕*}	0.13	(44)	-25.8
Michigan	2.6 ^{⊕*}	0.10	(34)	-16.1	West Virginia	4.3 ^{⊕*}	0.29	(03)	38.7
Minnesota	2.0 ^{⊕*}	0.13	(48)	-35.5	Wisconsin	2.5 ^{⊕*}	0.14	(36)	-19.4
Mississippi	4.1 ^{⊕*}	0.24	(05)	32.3	Wyoming	2.4 [*]	0.44	(43)	-22.6
Missouri	3.3	0.15	(21)	6.5					

^ä NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.


SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

^{*} Absolute percent difference between state rate and total U.S. rate is 15% or more.

[⊕] Difference between state rate and total U.S. rate is statistically significant (p<=.0002).


SEER Incidence and U.S. Death Rates⁺ Cervix Uteri By Race




⁺ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program.

SEER Incidence and U.S. Death Rates⁺ Cervix Uteri, By Age and Race


White - Under 50 Years of Age




Black - Under 50 Years of Age



White - Ages 50 and Over



Black - Ages 50 and Over



• Incidence × Mortality

⁺ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program.