

Smoking-Attributable Deaths, 1999

State	Deaths per 100,000
All States	295.5
Alabama	326.6
Alaska	289.6
Arizona	283.5
Arkansas	343.3
California	261.8
Colorado	248.1
Connecticut	255.9
Delaware	313.9
District of Columbia	250.4
Florida	283.1
Georgia	333.9
Hawaii	174.3
Idaho	254.6
Illinois	303.1
Indiana	341.4
Iowa	265.6
Kansas	269.8
Kentucky	387.1
Louisiana	314.1
Maine	305.2
Maryland	283.3
Massachusetts	261.2
Michigan	299.0
Minnesota	231.1
Mississippi	367.8
Missouri	345.3
Montana	294.7
Nebraska	264.2
Nevada	414.3
New Hampshire	291.9
New Jersey	243.7
New Mexico	264.7
New York	252.1
North Carolina	301.1
North Dakota	227.7
Ohio	317.6
Oklahoma	320.1
Oregon	278.6
Pennsylvania	270.2
Rhode Island	283.1
South Carolina	315.6
South Dakota	230.2
Tennessee	346.0
Texas	288.8
Utah	159.8
Vermont	274.3
Virginia	290.8
Washington	284.9
West Virginia	354.1
Wisconsin	278.2
Wyoming	318.6

Source: Centers for Disease Control and Prevention. Smoking-Attributable Mortality, Morbidity, and Economic Costs (SAMMEC): Adult SAMMEC and Maternal and Child Health (MCH) SAMMEC software. Available at <http://www.cdc.gov/tobacco/sammecc>.