


**Number of Ohio youth projected to die prematurely from their smoking: 285,161**

## Adult Tobacco Use in Ohio

### Current Cigarette Smoking Among Adults Aged 18 and Older, 1999



## Youth Tobacco Use

GRADES 6-8		GRADES 9-12	
Current Cigarette Smoking	Current Any Tobacco Use	Current Cigarette Smoking	Current Any Tobacco Use
National*	9.2%	28.5%	34.8%
OH†	13.7%	33.4%	41.1%
Boys†	14.7%	33.7%	45.6%
Girls†	12.7%	33.0%	35.9%

Current Cigarette Smoking = smoked cigarettes on ≥1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥1 of the 30 days preceding the survey.

\*Source: National Youth Tobacco Survey, 1999.

†Source: Ohio Youth Tobacco Survey, 2000.


## Health Impact and Costs

AVERAGE ANNUAL DEATHS RELATED TO SMOKING, 1990-1994		AVERAGE ANNUAL YEARS OF POTENTIAL LIFE LOST,* 1990-1994	MEDICAL COSTS RELATED TO SMOKING, 1993	
Overall	19,527	270,475 years or an average of 13.9 years for each death due to smoking.	Ambulatory	\$484,900,000
Men	12,400		Hospital	\$765,890,000
Women	7,128		Nursing Home†	\$840,400,000
Death Rate	364/100,000		Drug	\$129,680,000
Rank	34		Other	\$273,180,000
(No. 1 is lowest death rate)		*Calculated to life expectancy	<b>Total</b>	<b>\$2,494,060,000</b>

**Smoking-attributable Medicaid expenditures, Ohio, fiscal year 1993: \$597,217,000**

†Preliminary estimates

## Lung Cancer Death Rate\*



\*1997 deaths per 100,000 population, adjusted to the 1970 total U.S. population.

Ohio

**Scheduled 2001 settlement payment to Ohio: \$348,780,049.22**

## Tobacco Control Funding, 2001



At 383% of the CDC Best Practices recommended lower estimate funding level, Ohio ranks 1st (No. 1 is the highest) for tobacco control funding. Ohio ranks 34th (No. 1 is the lowest death rate) for average annual deaths related to smoking.

<i>FUNDING SOURCE</i>	<i>FY01 AMOUNT</i>	<i>FUNDING CYCLE</i>
State Appropriation—Settlement (Tobacco Only)	\$234,861,000	7/00–6/01
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
<b>Subtotal: State Appropriation</b>	<b>\$234,861,000</b>	
Federal—CDC Office on Smoking and Health	\$1,525,232	6/00–5/01
Federal—SAMHSA	n/a	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	n/a	
<b>Subtotal: Federal/National Sources</b>	<b>\$1,525,232</b>	
<b>Total Per Capita Funding</b>	<b>\$236,386,232 \$20.82</b>	

The \$234,861,000 reflected above will be disbursed to an endowment fund where the funds will be used by the Foundation for tobacco use reduction programs. The Foundation may, but is not required to, treat these funds as an endowment. An additional \$435,120 is expected to be spent by the Department of Health for emergency assistance, including medication and oxygen for seniors whose health has been affected by tobacco use. An additional \$669,000 is expected to be spent by the Department of Public Safety to enforce Ohio's under-age tobacco use laws.

## Funding as a Percentage of CDC Best Practices Recommendations

	<i>RECOMMENDED FOR TOTAL PROGRAM ANNUAL COST</i>	<i>RECOMMENDED PER CAPITA FUNDING LEVEL</i>	<i>PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS</i>
Lower Estimate	\$61,735,000	\$5.52	383%
Upper Estimate	\$173,676,000	\$15.53	136%

## Excise Tax

Cigarette tax per pack . . . . .	24¢
Rank = 32 (No. 1 is highest tax)	
Number of packages of cigarettes sold and taxed, per capita, 1999 . . . . .	104.0
Rank = 39 (No. 1 is lowest cigarette sales)	
Federal and state taxes as a percentage of retail price . . . . .	18.1%
Annual gross tax revenue from cigarettes . . . . .	\$279,313,463
Smokeless tobacco tax . . . . .	.17% of wholesale sales price