
Health Impacts


| Smoking-Attributable Deaths, 1999 | | SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC | | Youth Projected to Die from Smoking, 1999–2000 | |
|---------------------------------------|--|---|---|--|--------------------------|
| Overall Men Women Death Rate | 7,412 4,933 2,479 326.6/100,000 | State Lung Cancer All States State CHD* All States | 101.3/100,000 90.2/100,000 60.6/100,000 59.7/100,000 | Overall Projected Death Rate Projected All | 111,273 9,905/100,000 |
| All States | 295.5/100,000 | State COPD† All States | 73.6/100,000 59.7/100,000 | States | 8,830/100,000 |

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99



Youth Tobacco Use


| GRADES 6–8 | | | GRADES 9–12 | | |
|-------------------|-------|-------------|-------------------|-------------|--|
| Current Cigarette | | Current Any | Current Cigarette | Current Any | |
| Smoking | | Tobacco Use | Smoking | Tobacco Use | |
| National* | 11.0% | 15.1% | 28.0% | 34.5% | |
| AL† | 19.1% | 26.5% | 30.2% | 37.6% | |
| Boys† | 22.1% | 32.0% | 32.5% | 44.3% | |
| Girls† | 15.7% | 20.4% | 27.8% | 30.7% | |

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Alabama Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*

*Calculations for racial/ethnic groups are based on combined 1999 and 2000 data.





Economic Impacts and Investments

Alabama is investing 8% of CDC's Best Practices lower estimated recommended funding and 3% of the upper estimated recommended funding. Alabama spent \$269 per capita on smoking-attributable direct medical expenditures. In 1998, about 9% (\$186,000,000 or \$352.36 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$433,000,000 Hospital \$300,000,000 Nursing Home \$241,000,000 Prescription Drugs \$110,000,000 Other \$86,000,000 Annual Total \$1,170,000,000 Annual Per Capita \$269

Smoking-Attributable **Productivity Costs, 1999**

| Annual Total Annual Per Capita | \$1,777,000,000 \$407 | | |
|-----------------------------------|--------------------------|--|--|
| | | | |
| | | | |

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$96,961,493 Gross cigarette tax revenue collected in 2000 \$70,665,000 Cigarette tax per pack was \$0.165 in 2001

Cigarette sales were 96.2 packs per capita in 2000

Investment in Tobacco Control

| FUNDING SOURCE | FY02 AMOUNT | FUNDING CYCLE |
|--|--------------|---------------|
| State Appropriation—Settlement (Tobacco Only)* | \$600,000 | 10/01-9/02 |
| State Appropriation—Excise Tax Revenue | \$0 | |
| State Appropriation—Other | \$0 | |
| Subtotal: State Appropriation | \$600,000 | |
| Federal—CDC Office on Smoking and Health | \$1,340,047 | 6/01-5/02 |
| Federal—SAMHSA | \$78,720 | 10/01-9/02 |
| Non-Government Source—American Legacy Foundation | \$208,156 | 10/00-9/01 |
| Non-Government Source—RWJF/AMA | \$0 | |
| Subtotal: Federal/National Sources | \$1,626,923 | |
| FY02 Total Investment in Tobacco Control | \$2,226,923 | |
| CDC Best Practices Recommended Annual Total (Lower F | \$26,740,000 | |
| CDC Best Practices Recommended Annual Total (Upper E | \$71,235,000 | |
| FY02 Per Capita Investment in Tobacco Control | \$0.50 | |

^{*}An additional \$2,000,000 was appropriated in FY02 to the Department of Education for drug, alcohol, or tobacco education and prevention programs.