

**Past Month Cigarette Smoking
Among Youth—Grades 9–12, 1997**

| State | Prevalence (Percentage) |
|------------------------|----------------------------|
| Weighted Data | |
| Alabama | 35.8 |
| Arkansas | 43.2 |
| Connecticut | 35.2 |
| District of Columbia | 22.7 |
| Hawaii | 29.2 |
| Iowa | 37.5 |
| Kentucky | 47.0 |
| Louisiana | 36.4 |
| Maine | 39.2 |
| Massachusetts | 34.4 |
| Michigan | 38.2 |
| Mississippi | 31.3 |
| Missouri | 40.3 |
| Montana | 38.1 |
| Nevada | 29.4 |
| New York | 32.9 |
| Ohio | 34.5 |
| Rhode Island | 35.4 |
| South Carolina | 38.6 |
| South Dakota | 44.0 |
| Utah | 16.4 |
| Vermont | 38.3 |
| West Virginia | 41.9 |
| Wisconsin | 36.0 |
| Wyoming | 37.4 |
| Unweighted Data | |
| California | 26.6 |
| Colorado | 36.6 |
| Delaware | 35.0 |
| Florida | 33.6 |
| New Hampshire | 39.6 |
| New Jersey | 37.9 |
| North Carolina | 35.8 |
| North Dakota | 45.0 |
| Tennessee | 38.6 |

**Past Month Smokeless Tobacco Use
Among Youth—Grades 9–12, 1997**

| State | Prevalence (Percentage) |
|------------------------|----------------------------|
| Weighted Data | |
| Alabama | 11.8 |
| Arkansas | 13.8 |
| Connecticut | 5.7 |
| District of Columbia | 3.2 |
| Hawaii | 3.4 |
| Iowa | 12.8 |
| Kentucky | 15.6 |
| Louisiana | 10.1 |
| Maine | 7.9 |
| Massachusetts | 6.0 |
| Michigan | 8.4 |
| Mississippi | 6.8 |
| Missouri | 9.1 |
| Montana | 21.0 |
| Nevada | 9.5 |
| New York | 5.1 |
| Ohio | 10.4 |
| Rhode Island | 6.1 |
| South Carolina | 6.7 |
| South Dakota | 17.7 |
| Utah | 6.5 |
| Vermont | 7.8 |
| West Virginia | 15.8 |
| Wisconsin | 11.4 |
| Wyoming | 22.5 |
| Unweighted Data | |
| California | 4.2 |
| Colorado | 12.7 |
| Delaware | 6.7 |
| Florida | 8.8 |
| New Hampshire | 7.2 |
| New Jersey | 5.4 |
| North Carolina | 7.4 |
| Tennessee | 14.2 |

Source: Youth Risk Behavior Surveillance System (YRBSS), 1997