




# U.S. Decennial Life Tables for 1989-91

Volume II, State Life Tables Number 32, New Mexico

From the CENTERS FOR DISEASE CONTROL AND PREVENTION/National Center for Health Statistics



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Centers for Disease Control and Prevention  
National Center for Health Statistics



**Copyright information**

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

---

**Suggested citation**

National Center for Health Statistics. U.S. decennial life tables for 1989–91, vol II, State life tables no. 32, New Mexico. Hyattsville, Maryland. 1998.

---

**Library of Congress Catalog Card Number 85-600190**

---


For sale by the U.S. Government Printing Office  
Superintendent of Documents  
Mail Stop: SSOP  
Washington, DC 20402-9328

---

# U.S. Decennial Life Tables for 1989-91

---

Volume II, State Life Tables Number 32, New Mexico



---

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Centers for Disease Control and Prevention  
National Center for Health Statistics

Hyattsville, Maryland  
May 1998

DHHS Publication No. PHS-98-1151-32

## **National Center for Health Statistics**

Edward J. Sondik, Ph.D., *Director*

Jack R. Anderson, *Deputy Director*

Jack R. Anderson, *Acting Associate Director for International Statistics*

Lester R. Curtin, Ph.D., *Acting Associate Director for Research and Methodology*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Analysis, Epidemiology, and Health Promotion*

P. Douglas Williams, *Acting Associate Director for Data Standards, Program Development, and Extramural Programs*

Edward L. Hunter, *Associate Director for Planning, Budget, and Legislation*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Vital and Health Statistics Systems*

Stephen E. Nieberding, *Associate Director for Management*

Charles J. Rothwell, *Associate Director for Data Processing and Services*

## **Division of Vital Statistics**

Mary Anne Freedman, *Director*

James A. Weed, Ph.D., *Deputy Director*

Robert J. Armstrong, *Actuarial Adviser*

Harry M. Rosenberg, Ph.D., *Chief, Mortality Statistics Branch*

Nicholas F. Pace, *Chief, Systems, Programming, and Statistical Resources Branch*

# Contents

Acknowledgments. . . . . iv  
Abstract . . . . . 1  
Introduction. . . . . 1  
Methodology. . . . . 1  
Results and discussion . . . . . 2  
Explanation of the columns of the life table. . . . . 2  
References. . . . . 3

**Detailed tables**

Average lifetime in years by race and sex: United States and each State in rank order, 1989–91. . . . . 4  
1. Life table for the total population: New Mexico, 1989–91 . . . . . 6  
2. Life table for males: New Mexico, 1989–91. . . . . 8  
3. Life table for females: New Mexico, 1989–91 . . . . . 10  
4. Life table for the white population: New Mexico, 1989–91. . . . . 12  
5. Life table for white males: New Mexico, 1989–91. . . . . 14  
6. Life table for white females: New Mexico, 1989–91 . . . . . 16  
7. Life table for the population other than white: New Mexico, 1989–91. . . . . 18  
8. Life table for males other than white: New Mexico, 1989–91 . . . . . 20  
9. Life table for females other than white: New Mexico, 1989–91 . . . . . 22  
10. Standard errors of the probability of dying: New Mexico, 1989–91 . . . . . 24  
11. Standard errors of the average remaining lifetime: New Mexico, 1989–91 . . . . . 26

## Acknowledgments

This report was prepared in the Division of Vital Statistics (DVS) under the guidance of an ad hoc committee chaired by Robert J. Armstrong and included Stephen C. Goss and Alice H. Wade of the Office of the Actuary, Social Security Administration; Gregory K. Spencer and Frederick W. Hollmann of the U.S. Bureau of the Census; and David P. Johnson, Lester R. Curtin, Nonie Atkinson, Kenneth D. Kochanek, Harry M. Rosenberg, Jeffrey D. Maurer, and Joseph D. Farrell from the National Center for Health Statistics.

Nonie Atkinson, formerly of the Office of Research and Methodology (ORM), was responsible for the overall computer systems analysis and design, and played a major role in writing the programs to produce the life tables and their variances. Lester R. Curtin, also of ORM, consulted on methodological issues including the preparation of standard errors for the life tables.

Joseph D. Farrell, Charles E. Royer, and David P. Johnson of the Systems, Programming, and Statistical Resources Branch,

DVS, coordinated data processing and developed computer processes that eased the workload of the actuarial statistician and the Publications Branch. They also provided major programming support in summarizing data basic to the calculation of the life tables.

Gregory K. Spencer and Frederick W. Hollmann of the U.S. Bureau of the Census furnished the modified-race populations that were used in the production of these tables.

Stephen C. Goss, Felicite C. Bell, and Bertram M. Kestenbaum of the Office of the Actuary, Social Security Administration, provided mortality data from the Medicare Program that were used at age 85 years and over. Vanetta A. Harrington of the Systems, Programming, and Statistical Resources Branch, DVS, provided content review, and Robert N. Anderson of the Mortality Statistics Branch, DVS, provided peer review. This report was edited by Demarius V. Miller and Patricia Keaton-Williams and typeset by Jacqueline M. Davis of the Publications Branch, Division of Data Services.

# New Mexico Life Tables: 1989–91

by Robert J. Armstrong, M.S.  
Division of Vital Statistics

## Abstract

The life tables in this report are current life tables for New Mexico based on age-specific death rates for the period 1989–91. The death rates were calculated using data from the 1990 census of population and deaths occurring in the United States to residents of New Mexico in the 3 years 1989–91. Presented are tables for the white population, the population other than white, and the black population, separately by sex and for both sexes combined, and also for the total population and for total males and total females. Standard errors of the probability of dying and of life expectancy are also provided.

## Introduction

The life tables in this report are current life tables for New Mexico based on age-specific death rates for the period 1989–91. With the exception of those aged 95 years and over (and to a lesser extent those aged 85–94 years), the death rates were calculated using data from the 1990 census of population and deaths occurring in the United States to residents of New Mexico in the 3 years 1989–91. Other publications in this decennial series present life tables for the United States and the other individual States. Generally, these reports show life tables calculated for the white population, the population other than white, and the black population separately by sex and for both sexes combined. Each of these reports also shows life tables for the total population, for total males, and for total females. Standard errors of the probability of dying and of life expectancy are also provided. However, life tables for the population other than white and for the black population in a State are not published when the total number of deaths for either males or females during the 3-year period is less than 700.

These life tables are the most recent in a series for the States that began with the 1939–41 period. Each of the tables in the series is based on a census of population and deaths in a 3-year period centered on the census year. Because State life tables are not currently produced on an annual basis, the decennial life tables are the only source of State life expectancy data available at the National Center for Health Statistics (NCHS).

---

**Keywords:** New Mexico • decennial life tables • 1989–91 • life expectancy

This report is 1 of 51 reports containing life tables for the individual States and the District of Columbia. A separate report describes the methods and formulas by which these life tables were prepared in *U.S. Decennial Life Tables for 1989–91, Volume I, Number 2, Methodology of the National and State Life Tables* (1).

## Methodology

The general methodology, with a few modifications, used in preparing these life tables was developed by Thomas N. E. Greville for the 1939–41 decennial life tables (2). The life tables are based on a complete count of deaths to residents of New Mexico that occurred anywhere in the United States during the 3 years of 1989, 1990, and 1991 and on the 1990 census of population for New Mexico. However, sometimes the observed death rates that these data produced did not meet certain well-established criteria, such as steadily increasing mortality with increasing age. For example, when the pattern of age-specific death rates at some ages was jagged rather than smooth or when the rates by race or sex were inconsistent, the observed death rates were adjusted slightly by moving deaths from one age group to another within the race-sex group. The total number of deaths in a race-sex group was never changed. Certain other adjustments were made. In accordance with standard practice, deaths for which age was not stated were allocated proportionately among the various age groups.

The population data used differ from the official data published by the U.S. Bureau of the Census because of age reporting problems in the 1990 census. Age was based on the respondents' direct reports of age at last birthday in the 1990 census. It was apparent that many respondents had reported their age at either the time of completion of the census form or at the time of the interview by an enumerator, which could have occurred several months after the April 1 reference date. As a result, reported age was biased upward and had to be modified.

Between the ages of 5 and 94 years, death rates were calculated using the total number of deaths in 1989–91 and 3 times the population shown in the 1990 census. However, since population counts at ages under 2 years are considered to be less reliable than those at other ages, life-table values at ages under 2 years were derived from the reported numbers of births for each of the years 1987 to 1991. At ages 2–4 years, the denominator of the death rates used the populations at ages

$x-1$ ,  $x$ , and  $x+1$  (instead of 3 times the population at age  $x$ ). Death rates at ages 95 years and over, where the data from the census and from registered deaths are scanty and the accuracy of the reporting of age is not as good as at younger ages, are based on data from the Medicare program. However, when the data from the Medicare program were judged to be unreliable (usually after age 97), an algorithm was used to produce the death rates. The new algorithm, which differed from the one used for the 1979–81 decennial life tables, incremented the death rates more rapidly resulting in lower life expectancies at the extreme ages than in the previous reports. The rates based on the Medicare program and on the algorithm are differentiated by race and sex but not by State, so the same rates are used for each State. As a consequence, the probabilities of dying and the life expectancies at ages 85 years and over may fail to adequately reflect variation in mortality among the States, but such variation is in general smaller than differences associated with race and sex. Death rates at ages 85–94 years were adjusted to provide a smooth transition between the death rates based on the census and registered deaths and those derived from the Medicare program.

The population and death statistics at ages under 85 years are known to be subject to reporting errors, but these were not considered to be serious enough to require adjustment prior to the calculation of the life tables. In some instances, fluctuations due to small numbers of deaths produced anomalous life-tables values, which were eliminated by minor redistribution of deaths by age. For a complete description of the methodology used in preparing these life tables, see *U.S. Decennial Life Tables for 1989–91, Volume I, Number 2, Methodology of the National and State Life Tables* (1).

## Results and discussion

The life tables in this report are current life tables and are based on age-specific death rates for the period 1989–91. They may also be characterized as “cross-sectional.” They assume that a hypothetical cohort is traced from birth until the death of the last survivor and that it is subject throughout its existence to the age-specific death rates observed for 1989–91. For example, [table 3](#) is a life table for females. This table shows the progression of a cohort starting with 100,000 live births who were subjected to the average annual death rates observed among females in New Mexico in the 3-year period 1989–91 during its passage through successive years of age.

Column 7 of [table 3](#) shows the average number of years of life remaining to those in the cohort who attain each birthday. This average remaining lifetime is commonly called the expectation of life, and the expectation of life at birth is frequently used as a measure of comparative longevity. According to the 1989–91 life tables for New Mexico, the expectation of life at birth is 72.20 years for total males and 79.33 years for total females. Among the 50 States and the District of Columbia in the expectation of life at birth for the total population, New Mexico ranks 25th.

The ranking table shows the average lifetime (or expectation of life at birth) by race and sex for the population of the

United States, each State, and the District of Columbia. The States are ranked using the life expectancy at birth for the total population of the State.

These life tables are based on a complete count of resident deaths in New Mexico during the 3 years 1989, 1990, and 1991. As such, they are not subject to sampling error. However, even complete counts may be considered as one of a large series of possible results that could have arisen under the same circumstances. This type of variation is known as random error. The standard errors shown in this report reflect random error only, not other errors such as misreporting of age on death certificates or in the census.

The probabilities of dying and the expectation of life presented in this report are “point estimates.” They do not give the reader an indication of how accurate they are. Therefore standard errors of these two measures are also presented. Standard errors can be used to develop confidence intervals within which the “point estimates” are believed to lie. Standard errors of the probability of dying and of life expectancy contain six and three decimal places, respectively, and are shown in [tables 10](#) and [11](#). In both cases, the standard errors contain one place more than the corresponding variable in the life tables. In computing confidence intervals, the limits are rounded to the same number of decimal places that the variable has in the life table.

Even though 68 percent confidence intervals are rarely used because of their high degree of uncertainty, they are shown here to demonstrate the method of construction of confidence intervals. To obtain a 68 percent confidence interval for the probability of dying at any age, take the point estimate from column 2 of the appropriate life table and add and subtract one standard error from the table that gives the standard errors of the probability of dying ([table 10](#)). The 95 percent confidence interval is obtained by adding and subtracting two standard errors. For example, the probability that a 50-year-old white female will die before her 51st birthday is 0.00316 with a standard error of 0.000401. Therefore, the 68 percent confidence interval is from 0.00276 to 0.00356 and the 95 percent confidence interval is from 0.00236 to 0.00396. The life expectancy of a 50-year-old white female is 32.16 years with a standard error of 0.089 years. The 68 percent confidence interval for the life expectancy is therefore from 32.07 to 32.25 years and the 95 percent confidence interval is from 31.98 to 32.34 years.

## Explanation of the columns of the life table

*Column 1—Age interval ( $x$  to  $x+1$ )*—The age interval shown in column 1 is the interval of 1 year between the two exact ages indicated. For instance, “21–22” indicates the interval between the 21st birthday and the 22d, in other words, the 22d year of life.

*Column 2—Proportion dying ( $q_x$ )*—This column shows the proportion of the members of the life-table cohort alive at the beginning of the indicated year of age who will die before reaching the next birthday on the basis of the mortality rates of



1989–91 in New Mexico. For example, for females who reach age 21, the proportion dying before reaching their 22d birthday is 0.00084—out of every 1,000 female babies surviving to age 21, 0.84 will die before reaching their 22d birthday.

*Column 3—Number surviving ( $l_x$ )*—This column shows the number of persons, starting with a cohort of 100,000 live births, who will survive to the birthday marking the beginning of the indicated year of age. Thus out of 100,000 female babies born alive in the cohort of [table 3](#), 99,271 will complete the first year of life and enter the second, 98,492 will reach age 21, and 71,130 will live to age 75.

*Column 4—Number dying ( $d_x$ )*—This column shows the number dying in each successive age interval out of 100,000 live births. Thus out of 100,000 females born alive, 729 will die in the first year of life, 82 in the 22d year, and 2,135 in the 76th year. Each figure in column 4 is the difference between two successive figures in column 3.

*Columns 5 and 6—Stationary population ( $L_x$  and  $T_x$ )*—Suppose that a group of 100,000 persons like that assumed in columns 3 and 4 is born every year, and that the proportion dying in each such group in each age interval throughout the lives of the members is exactly that shown in column 2. If there were no migration and if the births were evenly distributed over the year, the survivors of these births would constitute what is called a stationary population, because in such a population the number of persons living in any given age interval would never change. When an individual left an age interval, whether by death or growing older and entering the next higher age interval, his place would immediately be taken by someone entering from the next lower age interval. Thus a census taken at any time in such a stationary community would always show the same total population and the same numerical distribution of that population among the various age intervals. In such a stationary population supported by 100,000 annual births, column 3 shows the number of persons who, each year, will reach the exact age that marks the beginning of the age interval indicated in column 1, and column 4 shows the number of persons who will die each year in that year of age interval.

Column 5,  $L_x$ , shows the number of females in the stationary population in the indicated year of age. For example, the figure shown in [table 3](#) for the year of age 21–22 is 98,451. This means that in a stationary population supported by

100,000 annual births, and with proportions dying in each age interval always in accordance with column 2, a census taken on any date would show 98,451 persons at age 21 (that is, between exact ages 21 and 22 years).

Column 6,  $T_x$ , shows the total number of persons in the stationary population in the indicated year of age and all subsequent years of age. For example, in the stationary population of females described in the preceding paragraph, column 6 shows that there would be at any given moment a total of 5,854,616 persons who had reached their 21st birthday. The population at all ages 0 and above (in other words, the total female population of the stationary community) would be 7,932,733.

*Column 7—Average remaining lifetime ( ${}^o e_x$ )*—The average remaining lifetime (also called expectation of life) at any given age is the average number of years remaining to be lived by those surviving to that age, on the basis of a given set of age-specific rates of dying. In order to relate these figures to the preceding columns of the life table, it is necessary to observe that the figures in column 5 of the life tables can also be interpreted in terms of a single life-table cohort without introducing the concept of the stationary population. From this point of view, each figure in column 5 represents the total time in years lived between two indicated birthdays by all those reaching the younger age among the survivors of a cohort of 100,000 live births. Thus the figure of 98,451 for females in New Mexico in the year of age 21–22 is the total number of years of life lived between their 21st and 22d birthdays by the 98,492 (column 3) who reached their 21st birthday out of the original cohort of 100,000 females born alive. The corresponding figure (5,854,616) in column 6 is the total number of years lived after attaining age 21 by the 98,492 reaching that exact age. This number of years divided by the number of persons (5,854,616 divided by 98,492) gives 59.44 years as the average remaining lifetime at age 21 for females in New Mexico.

## References

1. U.S. decennial life tables for 1989–91, volume I, number 2, methodology of the national and State life tables. In progress.
2. Greville TNE. United States life tables and actuarial tables, 1939–41. Washington: U.S. Government Printing Office. 1947.

Average lifetime in years by race and sex: United States and each State in rank order, 1989-91

Rank	Area	Total			White			All other					
								Total			Black		
		Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
1	Hawaii	78.21	75.37	81.26	77.92	75.12	81.09	78.40	75.49	81.48	*	*	*
2	Minnesota	77.76	74.53	80.85	77.97	74.78	81.02	73.05	69.46	76.80	*	*	*
3	Utah	77.70	74.93	80.38	77.77	75.00	80.44	*	*	*	*	*	*
4	North Dakota	77.62	74.35	80.99	77.99	74.74	81.32	*	*	*	*	*	*
5	Iowa	77.29	73.89	80.54	77.38	73.98	80.62	*	*	*	*	*	*
6	Colorado	76.96	73.79	80.01	77.06	73.88	80.13	75.71	72.63	78.61	72.41	68.96	75.89
7	Nebraska	76.92	73.57	80.17	77.21	73.87	80.44	71.14	67.64	74.52	*	*	*
8	Connecticut	76.91	73.62	79.97	77.44	74.25	80.37	72.31	67.82	76.61	70.84	66.04	75.44
8	South Dakota	76.91	73.17	80.77	77.91	74.30	81.59	*	*	*	*	*	*
10	Idaho	76.88	73.88	79.93	76.89	73.90	79.93	*	*	*	*	*	*
11	Wisconsin	76.87	73.61	80.03	77.18	73.99	80.27	72.37	68.27	76.25	70.96	66.42	75.27
12	Washington	76.82	73.84	79.74	76.92	73.97	79.81	76.09	72.72	79.59	71.34	67.91	75.58
13	Kansas	76.76	73.40	79.99	77.06	73.72	80.25	72.77	69.25	76.26	71.22	67.48	75.04
14	Massachusetts	76.72	73.32	79.80	76.90	73.54	79.95	75.08	71.29	78.60	72.45	68.17	76.50
14	New Hampshire	76.72	73.52	79.77	76.68	73.48	79.74	*	*	*	*	*	*
16	Rhode Island	76.54	73.00	79.77	76.80	73.31	79.97	*	*	*	*	*	*
16	Vermont	76.54	73.29	79.68	76.50	73.25	79.65	*	*	*	*	*	*
18	Oregon	76.44	73.21	79.67	76.51	73.28	79.73	75.24	72.02	78.45	*	*	*
19	Maine	76.35	72.98	79.61	76.35	72.98	79.61	*	*	*	*	*	*
20	Montana	76.23	73.05	79.49	76.72	73.59	79.92	*	*	*	*	*	*
21	Wyoming	76.21	73.16	79.29	76.34	73.27	79.46	*	*	*	*	*	*
22	Arizona	76.10	72.66	79.58	76.42	73.04	79.84	72.76	68.89	76.81	70.84	67.20	74.90
23	California	75.86	72.53	79.19	75.92	72.61	79.26	75.79	72.34	79.18	69.65	65.43	74.07
24	Florida	75.84	72.10	79.60	76.82	73.19	80.46	69.82	65.40	74.19	68.77	64.26	73.28
25	New Mexico	75.74	72.20	79.33	76.08	72.66	79.53	73.41	68.97	77.93	*	*	*
26	New Jersey	75.42	72.16	78.49	76.46	73.37	79.34	70.73	66.59	74.66	68.47	63.87	72.88
27	Indiana	75.39	71.99	78.62	75.82	72.44	79.03	70.76	66.99	74.35	69.80	65.87	73.56
28	Pennsylvania	75.38	71.91	78.66	76.15	72.81	79.28	69.34	64.69	73.78	68.27	63.33	73.02
	United States	75.37	71.83	78.81	76.13	72.72	79.45	71.25	66.97	75.39	69.16	64.47	73.73
29	Ohio	75.32	71.99	78.45	75.93	72.70	78.95	70.86	66.70	74.82	70.15	65.80	74.29
30	Missouri	75.25	71.54	78.82	76.02	72.43	79.48	69.65	65.00	74.07	68.81	63.87	73.52
31	Virginia	75.22	71.77	78.56	76.34	73.04	79.48	71.17	67.03	75.27	70.05	65.75	74.37
32	Texas	75.14	71.41	78.87	75.75	72.08	79.42	71.25	67.08	75.38	69.79	65.36	74.23
33	Oklahoma	75.10	71.63	78.49	75.21	71.76	78.59	74.81	71.17	78.21	70.85	67.10	74.48
34	Michigan	75.04	71.71	78.24	76.18	73.06	79.14	69.22	64.68	73.65	68.49	63.68	73.18
35	Illinois	74.90	71.34	78.31	76.16	72.83	79.33	69.25	64.58	73.79	67.46	62.41	72.39
36	Alaska	74.83	71.60	78.60	75.83	72.82	79.40	71.67	67.65	76.17	*	*	*
37	Maryland	74.79	71.31	78.13	76.30	73.20	79.23	70.76	66.27	75.15	69.69	64.99	74.31
38	Delaware	74.76	71.63	77.74	75.76	72.75	78.62	70.06	66.39	73.63	69.26	65.51	72.91
39	New York	74.68	70.86	78.32	75.61	72.01	79.03	71.53	66.70	75.97	69.33	63.86	74.35
40	North Carolina	74.48	70.58	78.27	75.89	72.21	79.44	69.83	64.96	74.55	69.38	64.38	74.24
41	Kentucky	74.37	70.72	77.97	74.65	71.01	78.24	70.79	66.78	74.63	70.16	66.06	74.13
42	Arkansas	74.33	70.54	78.13	75.20	71.54	78.89	69.63	64.87	74.13	68.93	64.03	73.58
43	Tennessee	74.32	70.38	78.18	75.27	71.38	79.10	69.43	64.99	73.59	68.97	64.41	73.24
44	West Virginia	74.26	70.53	77.93	74.37	70.66	78.02	71.20	66.77	75.46	69.75	65.00	74.36
45	Nevada	74.18	70.96	77.76	74.44	71.26	77.99	72.74	69.15	76.42	*	*	*
46	Alabama	73.64	69.59	77.61	75.01	71.12	78.85	69.59	64.79	74.05	69.23	64.37	73.76
47	Georgia	73.61	69.65	77.46	75.24	71.46	78.94	69.21	64.49	73.65	68.79	63.98	73.34
48	South Carolina	73.51	69.59	77.34	75.33	71.62	78.97	69.09	64.37	73.57	68.82	64.07	73.35
49	Louisiana	73.05	69.10	76.93	74.87	71.15	78.54	68.99	64.33	73.43	68.62	63.84	73.16
50	Mississippi	73.03	68.90	77.10	74.78	70.74	78.82	69.54	64.84	73.91	69.41	64.66	73.82
51	District Of Columbia	67.99	61.97	74.23	76.09	71.36	81.06	64.97	58.14	72.03	64.44	57.53	71.61

\* Figure does not meet standards of reliability and precision.

## **Detailed tables**

**Table 1. Life table for the total population: New Mexico, 1989–91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0-1	.00855	100,000	855	99,332	7,574,197	75.74
1-2	.00078	99,145	78	99,106	7,474,865	75.39
2-3	.00057	99,067	56	99,039	7,375,759	74.45
3-4	.00045	99,011	45	98,988	7,276,720	73.49
4-5	.00036	98,966	35	98,949	7,177,732	72.53
5-6	.00031	98,931	31	98,915	7,078,783	71.55
6-7	.00029	98,900	29	98,886	6,979,868	70.58
7-8	.00027	98,871	26	98,858	6,880,982	69.60
8-9	.00023	98,845	23	98,834	6,782,124	68.61
9-10	.00020	98,822	19	98,812	6,683,290	67.63
10-11	.00017	98,803	17	98,795	6,584,478	66.64
11-12	.00017	98,786	16	98,778	6,485,683	65.65
12-13	.00023	98,770	23	98,758	6,386,905	64.66
13-14	.00038	98,747	38	98,728	6,288,147	63.68
14-15	.00060	98,709	59	98,680	6,189,419	62.70
15-16	.00084	98,650	83	98,608	6,090,739	61.74
16-17	.00108	98,567	106	98,514	5,992,131	60.79
17-18	.00127	98,461	125	98,398	5,893,617	59.86
18-19	.00140	98,336	139	98,266	5,795,219	58.93
19-20	.00149	98,197	146	98,125	5,696,953	58.02
20-21	.00158	98,051	154	97,973	5,598,828	57.10
21-22	.00167	97,897	164	97,815	5,500,855	56.19
22-23	.00172	97,733	168	97,650	5,403,040	55.28
23-24	.00173	97,565	168	97,481	5,305,390	54.38
24-25	.00171	97,397	167	97,313	5,207,909	53.47
25-26	.00167	97,230	162	97,149	5,110,596	52.56
26-27	.00164	97,068	159	96,989	5,013,447	51.65
27-28	.00163	96,909	158	96,830	4,916,458	50.73
28-29	.00166	96,751	160	96,671	4,819,628	49.81
29-30	.00171	96,591	166	96,508	4,722,957	48.90
30-31	.00177	96,425	170	96,340	4,626,449	47.98
31-32	.00182	96,255	176	96,167	4,530,109	47.06
32-33	.00187	96,079	180	95,990	4,433,942	46.15
33-34	.00192	95,899	183	95,807	4,337,952	45.23
34-35	.00196	95,716	188	95,622	4,242,145	44.32
35-36	.00201	95,528	192	95,432	4,146,523	43.41
36-37	.00207	95,336	198	95,237	4,051,091	42.49
37-38	.00214	95,138	203	95,037	3,955,854	41.58
38-39	.00221	94,935	210	94,830	3,860,817	40.67
39-40	.00229	94,725	217	94,617	3,765,987	39.76
40-41	.00238	94,508	224	94,396	3,671,370	38.85
41-42	.00248	94,284	234	94,167	3,576,974	37.94
42-43	.00260	94,050	245	93,927	3,482,807	37.03
43-44	.00276	93,805	259	93,675	3,388,880	36.13
44-45	.00295	93,546	275	93,409	3,295,205	35.23
45-46	.00319	93,271	298	93,122	3,201,796	34.33
46-47	.00346	92,973	321	92,812	3,108,674	33.44
47-48	.00374	92,652	347	92,479	3,015,862	32.55
48-49	.00398	92,305	367	92,121	2,923,383	31.67
49-50	.00419	91,938	385	91,746	2,831,262	30.80
50-51	.00442	91,553	405	91,351	2,739,516	29.92
51-52	.00472	91,148	430	90,933	2,648,165	29.05
52-53	.00509	90,718	461	90,487	2,557,232	28.19
53-54	.00556	90,257	502	90,006	2,466,745	27.33
54-55	.00612	89,755	550	89,480	2,376,739	26.48

**Table 1. Life table for the total population: New Mexico, 1989–91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55–56	.00673	89,205	600	88,906	2,287,259	25.64
56–57	.00737	88,605	653	88,278	2,198,353	24.81
57–58	.00809	87,952	712	87,596	2,110,075	23.99
58–59	.00889	87,240	775	86,853	2,022,479	23.18
59–60	.00975	86,465	843	86,043	1,935,626	22.39
60–61	.01066	85,622	913	85,165	1,849,583	21.60
61–62	.01161	84,709	984	84,218	1,764,418	20.83
62–63	.01262	83,725	1,056	83,197	1,680,200	20.07
63–64	.01370	82,669	1,133	82,102	1,597,003	19.32
64–65	.01485	81,536	1,211	80,931	1,514,901	18.58
65–66	.01605	80,325	1,289	79,680	1,433,970	17.85
66–67	.01731	79,036	1,368	78,352	1,354,290	17.14
67–68	.01870	77,668	1,453	76,942	1,275,938	16.43
68–69	.02032	76,215	1,548	75,441	1,198,996	15.73
69–70	.02222	74,667	1,659	73,837	1,123,555	15.05
70–71	.02439	73,008	1,780	72,118	1,049,718	14.38
71–72	.02680	71,228	1,909	70,273	977,600	13.73
72–73	.02945	69,319	2,042	68,298	907,327	13.09
73–74	.03223	67,277	2,168	66,193	839,029	12.47
74–75	.03506	65,109	2,282	63,968	772,836	11.87
75–76	.03801	62,827	2,389	61,633	708,868	11.28
76–77	.04121	60,438	2,490	59,193	647,235	10.71
77–78	.04471	57,948	2,591	56,652	588,042	10.15
78–79	.04868	55,357	2,695	54,010	531,390	9.60
79–80	.05327	52,662	2,805	51,259	477,380	9.06
80–81	.05854	49,857	2,918	48,398	426,121	8.55
81–82	.06441	46,939	3,024	45,427	377,723	8.05
82–83	.07092	43,915	3,114	42,358	332,296	7.57
83–84	.07796	40,801	3,181	39,210	289,938	7.11
84–85	.08561	37,620	3,221	36,009	250,728	6.66
85–86	.09488	34,399	3,264	32,768	214,719	6.24
86–87	.10549	31,135	3,284	29,493	181,951	5.84
87–88	.11655	27,851	3,246	26,228	152,458	5.47
88–89	.12762	24,605	3,140	23,034	126,230	5.13
89–90	.13891	21,465	2,982	19,974	103,196	4.81
90–91	.15145	18,483	2,799	17,084	83,222	4.50
91–92	.16564	15,684	2,598	14,384	66,138	4.22
92–93	.18041	13,086	2,361	11,906	51,754	3.95
93–94	.19511	10,725	2,093	9,679	39,848	3.72
94–95	.20977	8,632	1,810	7,727	30,169	3.49
95–96	.22502	6,822	1,535	6,054	22,442	3.29
96–97	.24126	5,287	1,276	4,649	16,388	3.10
97–98	.25689	4,011	1,030	3,496	11,739	2.93
98–99	.27175	2,981	810	2,575	8,243	2.77
99–100	.28751	2,171	624	1,859	5,668	2.61
100–101	.30418	1,547	471	1,311	3,809	2.46
101–102	.32182	1,076	346	903	2,498	2.32
102–103	.34049	730	249	606	1,595	2.19
103–104	.36024	481	173	395	989	2.05
104–105	.38113	308	117	249	594	1.93
105–106	.40324	191	77	152	345	1.81
106–107	.42663	114	49	89	193	1.70
107–108	.45137	65	29	51	104	1.59
108–109	.47755	36	17	27	53	1.49
109–110	.50525	19	10	14	26	1.39

**Table 2. Life table for males: New Mexico, 1989-91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
0-1	.00978	100,000	978	99,228	7,219,944	72.20
1-2	.00087	99,022	87	98,978	7,120,716	71.91
2-3	.00064	98,935	63	98,904	7,021,738	70.97
3-4	.00052	98,872	52	98,846	6,922,834	70.02
4-5	.00042	98,820	41	98,800	6,823,988	69.05
5-6	.00035	98,779	35	98,761	6,725,188	68.08
6-7	.00033	98,744	32	98,728	6,626,427	67.11
7-8	.00031	98,712	31	98,697	6,527,699	66.13
8-9	.00027	98,681	26	98,668	6,429,002	65.15
9-10	.00022	98,655	22	98,644	6,330,334	64.17
10-11	.00017	98,633	17	98,624	6,231,690	63.18
11-12	.00018	98,616	17	98,607	6,133,066	62.19
12-13	.00028	98,599	28	98,585	6,034,459	61.20
13-14	.00052	98,571	51	98,545	5,935,874	60.22
14-15	.00085	98,520	84	98,478	5,837,329	59.25
15-16	.00123	98,436	121	98,376	5,738,851	58.30
16-17	.00158	98,315	156	98,238	5,640,475	57.37
17-18	.00188	98,159	184	98,067	5,542,237	56.46
18-19	.00208	97,975	204	97,873	5,444,170	55.57
19-20	.00221	97,771	217	97,662	5,346,297	54.68
20-21	.00234	97,554	228	97,440	5,248,635	53.80
21-22	.00248	97,326	241	97,206	5,151,195	52.93
22-23	.00256	97,085	249	96,960	5,053,989	52.06
23-24	.00258	96,836	250	96,711	4,957,029	51.19
24-25	.00255	96,586	246	96,463	4,860,318	50.32
25-26	.00250	96,340	241	96,219	4,763,855	49.45
26-27	.00245	96,099	235	95,982	4,667,636	48.57
27-28	.00245	95,864	235	95,746	4,571,654	47.69
28-29	.00251	95,629	240	95,509	4,475,908	46.80
29-30	.00261	95,389	248	95,265	4,380,399	45.92
30-31	.00272	95,141	259	95,011	4,285,134	45.04
31-32	.00282	94,882	267	94,748	4,190,123	44.16
32-33	.00289	94,615	274	94,478	4,095,375	43.28
33-34	.00293	94,341	276	94,203	4,000,897	42.41
34-35	.00295	94,065	277	93,926	3,906,694	41.53
35-36	.00296	93,788	278	93,649	3,812,768	40.65
36-37	.00299	93,510	279	93,371	3,719,119	39.77
37-38	.00304	93,231	283	93,089	3,625,748	38.89
38-39	.00312	92,948	290	92,804	3,532,659	38.01
39-40	.00322	92,658	298	92,509	3,439,855	37.12
40-41	.00334	92,360	309	92,205	3,347,346	36.24
41-42	.00347	92,051	319	91,892	3,255,141	35.36
42-43	.00361	91,732	331	91,566	3,163,249	34.48
43-44	.00378	91,401	346	91,228	3,071,683	33.61
44-45	.00397	91,055	361	90,875	2,980,455	32.73
45-46	.00422	90,694	383	90,502	2,889,580	31.86
46-47	.00452	90,311	408	90,108	2,799,078	30.99
47-48	.00482	89,903	433	89,686	2,708,970	30.13
48-49	.00509	89,470	456	89,243	2,619,284	29.28
49-50	.00534	89,014	475	88,776	2,530,041	28.42
50-51	.00561	88,539	496	88,291	2,441,265	27.57
51-52	.00596	88,043	525	87,780	2,352,974	26.73
52-53	.00645	87,518	565	87,236	2,265,194	25.88
53-54	.00713	86,953	620	86,643	2,177,958	25.05
54-55	.00796	86,333	687	85,989	2,091,315	24.22
55-56	.00886	85,646	759	85,266	2,005,326	23.41
56-57	.00978	84,887	831	84,471	1,920,060	22.62
57-58	.01074	84,056	903	83,605	1,835,589	21.84

**Table 2. Life table for males: New Mexico, 1989–91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
58–59	.01173	83,153	976	82,665	1,751,984	21.07
59–60	.01276	82,177	1,049	81,653	1,669,319	20.31
60–61	.01384	81,128	1,123	80,566	1,587,666	19.57
61–62	.01498	80,005	1,198	79,406	1,507,100	18.84
62–63	.01617	78,807	1,275	78,170	1,427,694	18.12
63–64	.01743	77,532	1,351	76,857	1,349,524	17.41
64–65	.01876	76,181	1,429	75,466	1,272,667	16.71
65–66	.02010	74,752	1,503	74,001	1,197,201	16.02
66–67	.02153	73,249	1,577	72,461	1,123,200	15.33
67–68	.02323	71,672	1,665	70,840	1,050,739	14.66
68–69	.02536	70,007	1,775	69,119	979,899	14.00
69–70	.02795	68,232	1,906	67,279	910,780	13.35
70–71	.03095	66,326	2,053	65,299	843,501	12.72
71–72	.03423	64,273	2,201	63,173	778,202	12.11
72–73	.03771	62,072	2,341	60,902	715,029	11.52
73–74	.04119	59,731	2,460	58,501	654,127	10.95
74–75	.04463	57,271	2,556	55,994	595,626	10.40
75–76	.04823	54,715	2,639	53,395	539,632	9.86
76–77	.05224	52,076	2,720	50,716	486,237	9.34
77–78	.05675	49,356	2,801	47,955	435,521	8.82
78–79	.06202	46,555	2,888	45,111	387,566	8.32
79–80	.06820	43,667	2,978	42,179	342,455	7.84
80–81	.07555	40,689	3,074	39,152	300,276	7.38
81–82	.08379	37,615	3,151	36,039	261,124	6.94
82–83	.09238	34,464	3,184	32,872	225,085	6.53
83–84	.10059	31,280	3,147	29,706	192,213	6.14
84–85	.10842	28,133	3,050	26,609	162,507	5.78
85–86	.11799	25,083	2,959	23,603	135,898	5.42
86–87	.12941	22,124	2,863	20,692	112,295	5.08
87–88	.14164	19,261	2,728	17,897	91,603	4.76
88–89	.15432	16,533	2,552	15,256	73,706	4.46
89–90	.16743	13,981	2,341	12,811	58,450	4.18
90–91	.18142	11,640	2,111	10,585	45,639	3.92
91–92	.19682	9,529	1,876	8,590	35,054	3.68
92–93	.21310	7,653	1,631	6,838	26,464	3.46
93–94	.22954	6,022	1,382	5,331	19,626	3.26
94–95	.24522	4,640	1,138	4,071	14,295	3.08
95–96	.26004	3,502	911	3,047	10,224	2.92
96–97	.27536	2,591	713	2,234	7,177	2.77
97–98	.28943	1,878	544	1,606	4,943	2.63
98–99	.30390	1,334	405	1,132	3,337	2.50
99–100	.31910	929	297	781	2,205	2.37
100–101	.33505	632	211	526	1,424	2.25
101–102	.35181	421	148	347	898	2.13
102–103	.36940	273	101	222	551	2.02
103–104	.38787	172	67	139	329	1.91
104–105	.40726	105	43	83	190	1.81
105–106	.42762	62	26	49	107	1.71
106–107	.44900	36	16	28	58	1.61
107–108	.47145	20	10	15	30	1.52
108–109	.49503	10	5	8	15	1.43
109–110	.51978	5	2	4	7	1.35

**Table 3. Life table for females: New Mexico, 1989–91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0-1	.00729	100,000	729	99,441	7,932,733	79.33
1-2	.00069	99,271	68	99,237	7,833,292	78.91
2-3	.00049	99,203	49	99,179	7,734,055	77.96
3-4	.00037	99,154	37	99,136	7,634,876	77.00
4-5	.00030	99,117	29	99,102	7,535,740	76.03
5-6	.00027	99,088	27	99,075	7,436,638	75.05
6-7	.00024	99,061	24	99,049	7,337,563	74.07
7-8	.00022	99,037	22	99,025	7,238,514	73.09
8-9	.00020	99,015	20	99,006	7,139,489	72.11
9-10	.00017	98,995	17	98,986	7,040,483	71.12
10-11	.00016	98,978	15	98,971	6,941,497	70.13
11-12	.00015	98,963	16	98,954	6,842,526	69.14
12-13	.00018	98,947	18	98,939	6,743,572	68.15
13-14	.00024	98,929	24	98,917	6,644,633	67.17
14-15	.00034	98,905	33	98,889	6,545,716	66.18
15-16	.00044	98,872	44	98,850	6,446,827	65.20
16-17	.00054	98,828	53	98,801	6,347,977	64.23
17-18	.00063	98,775	63	98,743	6,249,176	63.27
18-19	.00070	98,712	68	98,678	6,150,433	62.31
19-20	.00074	98,644	74	98,607	6,051,755	61.35
20-21	.00079	98,570	78	98,532	5,953,148	60.39
21-22	.00084	98,492	82	98,451	5,854,616	59.44
22-23	.00087	98,410	85	98,367	5,756,165	58.49
23-24	.00087	98,325	86	98,282	5,657,798	57.54
24-25	.00085	98,239	84	98,197	5,559,516	56.59
25-26	.00083	98,155	81	98,114	5,461,319	55.64
26-27	.00082	98,074	81	98,034	5,363,205	54.69
27-28	.00081	97,993	79	97,954	5,265,171	53.73
28-29	.00081	97,914	79	97,874	5,167,217	52.77
29-30	.00082	97,835	81	97,795	5,069,343	51.82
30-31	.00083	97,754	81	97,713	4,971,548	50.86
31-32	.00085	97,673	83	97,632	4,873,835	49.90
32-33	.00088	97,590	85	97,547	4,776,203	48.94
33-34	.00093	97,505	91	97,459	4,678,656	47.98
34-35	.00100	97,414	98	97,365	4,581,197	47.03
35-36	.00109	97,316	106	97,264	4,483,832	46.07
36-37	.00118	97,210	114	97,152	4,386,568	45.12
37-38	.00126	97,096	123	97,035	4,289,416	44.18
38-39	.00132	96,973	128	96,909	4,192,381	43.23
39-40	.00138	96,845	133	96,778	4,095,472	42.29
40-41	.00143	96,712	139	96,643	3,998,694	41.35
41-42	.00150	96,573	145	96,500	3,902,051	40.41
42-43	.00160	96,428	155	96,351	3,805,551	39.47
43-44	.00175	96,273	168	96,189	3,709,200	38.53
44-45	.00194	96,105	187	96,011	3,613,011	37.59
45-46	.00217	95,918	208	95,815	3,517,000	36.67
46-47	.00243	95,710	233	95,593	3,421,185	35.75
47-48	.00269	95,477	256	95,349	3,325,592	34.83
48-49	.00291	95,221	277	95,082	3,230,243	33.92
49-50	.00310	94,944	294	94,797	3,135,161	33.02
50-51	.00330	94,650	313	94,494	3,040,364	32.12
51-52	.00355	94,337	335	94,169	2,945,870	31.23
52-53	.00382	94,002	359	93,823	2,851,701	30.34
53-54	.00410	93,643	384	93,451	2,757,878	29.45
54-55	.00442	93,259	412	93,053	2,664,427	28.57



Table 3. Life table for females: New Mexico, 1989–91—Con.

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55–56	.00475	92,847	441	92,626	2,571,374	27.69
56–57	.00513	92,406	474	92,169	2,478,748	26.82
57–58	.00562	91,932	517	91,673	2,386,579	25.96
58–59	.00626	91,415	572	91,129	2,294,906	25.10
59–60	.00700	90,843	636	90,525	2,203,777	24.26
60–61	.00778	90,207	702	89,856	2,113,252	23.43
61–62	.00858	89,505	768	89,121	2,023,396	22.61
62–63	.00945	88,737	839	88,317	1,934,275	21.80
63–64	.01040	87,898	914	87,441	1,845,958	21.00
64–65	.01141	86,984	992	86,488	1,758,517	20.22
65–66	.01249	85,992	1,075	85,455	1,672,029	19.44
66–67	.01361	84,917	1,156	84,339	1,586,574	18.68
67–68	.01476	83,761	1,236	83,144	1,502,235	17.93
68–69	.01595	82,525	1,316	81,867	1,419,091	17.20
69–70	.01727	81,209	1,403	80,507	1,337,224	16.47
70–71	.01874	79,806	1,495	79,059	1,256,717	15.75
71–72	.02044	78,311	1,601	77,511	1,177,658	15.04
72–73	.02246	76,710	1,723	75,848	1,100,147	14.34
73–74	.02479	74,987	1,859	74,058	1,024,299	13.66
74–75	.02732	73,128	1,998	72,129	950,241	12.99
75–76	.03001	71,130	2,135	70,062	878,112	12.35
76–77	.03285	68,995	2,266	67,863	808,050	11.71
77–78	.03584	66,729	2,391	65,533	740,187	11.09
78–79	.03911	64,338	2,517	63,079	674,654	10.49
79–80	.04282	61,821	2,647	60,498	611,575	9.89
80–81	.04694	59,174	2,777	57,786	551,077	9.31
81–82	.05160	56,397	2,910	54,941	493,291	8.75
82–83	.05719	53,487	3,059	51,958	438,350	8.20
83–84	.06397	50,428	3,226	48,815	386,392	7.66
84–85	.07201	47,202	3,399	45,503	337,577	7.15
85–86	.08195	43,803	3,589	42,009	292,074	6.67
86–87	.09301	40,214	3,740	38,343	250,065	6.22
87–88	.10437	36,474	3,807	34,570	211,722	5.80
88–89	.11550	32,667	3,773	30,781	177,152	5.42
89–90	.12676	28,894	3,663	27,062	146,371	5.07
90–91	.13955	25,231	3,521	23,471	119,309	4.73
91–92	.15420	21,710	3,348	20,036	95,838	4.41
92–93	.16931	18,362	3,108	16,808	75,802	4.13
93–94	.18414	15,254	2,809	13,849	58,994	3.87
94–95	.19901	12,445	2,477	11,207	45,145	3.63
95–96	.21475	9,968	2,141	8,898	33,938	3.40
96–97	.23143	7,827	1,811	6,921	25,040	3.20
97–98	.24775	6,016	1,491	5,271	18,119	3.01
98–99	.26375	4,525	1,193	3,929	12,848	2.84
99–100	.27957	3,332	932	2,866	8,919	2.68
100–101	.29635	2,400	711	2,045	6,053	2.52
101–102	.31413	1,689	531	1,423	4,008	2.37
102–103	.33298	1,158	385	966	2,585	2.23
103–104	.35296	773	273	636	1,619	2.10
104–105	.37413	500	187	407	983	1.97
105–106	.39658	313	124	251	576	1.84
106–107	.42038	189	80	149	325	1.72
107–108	.44560	109	48	85	176	1.61
108–109	.47233	61	29	46	91	1.50
109–110	.50068	32	16	24	45	1.40

**Table 4. Life table for the white population: New Mexico, 1989–91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0–1	.00813	100,000	813	99,357	7,607,906	76.08
1–2	.00070	99,187	70	99,152	7,508,549	75.70
2–3	.00052	99,117	51	99,092	7,409,397	74.75
3–4	.00042	99,066	42	99,045	7,310,305	73.79
4–5	.00034	99,024	33	99,007	7,211,260	72.82
5–6	.00030	98,991	30	98,976	7,112,253	71.85
6–7	.00027	98,961	27	98,948	7,013,277	70.87
7–8	.00025	98,934	25	98,922	6,914,329	69.89
8–9	.00023	98,909	22	98,898	6,815,407	68.91
9–10	.00019	98,887	19	98,877	6,716,509	67.92
10–11	.00016	98,868	16	98,860	6,617,632	66.93
11–12	.00016	98,852	16	98,844	6,518,772	65.94
12–13	.00022	98,836	22	98,826	6,419,928	64.96
13–14	.00036	98,814	35	98,797	6,321,102	63.97
14–15	.00055	98,779	54	98,751	6,222,305	62.99
15–16	.00078	98,725	77	98,687	6,123,554	62.03
16–17	.00099	98,648	97	98,599	6,024,867	61.07
17–18	.00116	98,551	114	98,494	5,926,268	60.13
18–19	.00128	98,437	126	98,374	5,827,774	59.20
19–20	.00135	98,311	133	98,244	5,729,400	58.28
20–21	.00142	98,178	139	98,109	5,631,156	57.36
21–22	.00150	98,039	147	97,965	5,533,047	56.44
22–23	.00154	97,892	151	97,817	5,435,082	55.52
23–24	.00155	97,741	152	97,665	5,337,265	54.61
24–25	.00154	97,589	150	97,514	5,239,600	53.69
25–26	.00151	97,439	147	97,366	5,142,086	52.77
26–27	.00148	97,292	144	97,220	5,044,720	51.85
27–28	.00148	97,148	144	97,076	4,947,500	50.93
28–29	.00151	97,004	146	96,931	4,850,424	50.00
29–30	.00156	96,858	152	96,782	4,753,493	49.08
30–31	.00162	96,706	156	96,628	4,656,711	48.15
31–32	.00167	96,550	162	96,469	4,560,083	47.23
32–33	.00172	96,388	165	96,306	4,463,614	46.31
33–34	.00176	96,223	170	96,138	4,367,308	45.39
34–35	.00181	96,053	174	95,966	4,271,170	44.47
35–36	.00186	95,879	178	95,790	4,175,204	43.55
36–37	.00191	95,701	183	95,610	4,079,414	42.63
37–38	.00198	95,518	188	95,424	3,983,804	41.71
38–39	.00204	95,330	195	95,233	3,888,380	40.79
39–40	.00212	95,135	202	95,034	3,793,147	39.87
40–41	.00220	94,933	209	94,829	3,698,113	38.95
41–42	.00230	94,724	217	94,616	3,603,284	38.04
42–43	.00241	94,507	228	94,393	3,508,668	37.13
43–44	.00256	94,279	241	94,158	3,414,275	36.21
44–45	.00275	94,038	258	93,909	3,320,117	35.31
45–46	.00298	93,780	280	93,639	3,226,208	34.40
46–47	.00326	93,500	304	93,348	3,132,569	33.50
47–48	.00352	93,196	329	93,032	3,039,221	32.61
48–49	.00376	92,867	349	92,692	2,946,189	31.72
49–50	.00396	92,518	366	92,335	2,853,497	30.84
50–51	.00418	92,152	385	91,960	2,761,162	29.96
51–52	.00447	91,767	410	91,561	2,669,202	29.09
52–53	.00485	91,357	443	91,136	2,577,641	28.22
53–54	.00534	90,914	485	90,671	2,486,505	27.35
54–55	.00592	90,429	536	90,161	2,395,834	26.49

**Table 4. Life table for the white population: New Mexico, 1989–91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55–56	.00656	89,893	590	89,598	2,305,673	25.65
56–57	.00723	89,303	646	88,980	2,216,075	24.82
57–58	.00797	88,657	707	88,304	2,127,095	23.99
58–59	.00878	87,950	772	87,564	2,038,791	23.18
59–60	.00965	87,178	841	86,758	1,951,227	22.38
60–61	.01056	86,337	912	85,880	1,864,469	21.60
61–62	.01150	85,425	983	84,934	1,778,589	20.82
62–63	.01250	84,442	1,056	83,914	1,693,655	20.06
63–64	.01358	83,386	1,132	82,821	1,609,741	19.30
64–65	.01472	82,254	1,211	81,648	1,526,920	18.56
65–66	.01590	81,043	1,289	80,399	1,445,272	17.83
66–67	.01714	79,754	1,367	79,071	1,364,873	17.11
67–68	.01852	78,387	1,452	77,661	1,285,802	16.40
68–69	.02015	76,935	1,550	76,161	1,208,141	15.70
69–70	.02207	75,385	1,663	74,553	1,131,980	15.02
70–71	.02427	73,722	1,789	72,828	1,057,427	14.34
71–72	.02673	71,933	1,923	70,971	984,599	13.69
72–73	.02943	70,010	2,061	68,979	913,628	13.05
73–74	.03227	67,949	2,192	66,853	844,649	12.43
74–75	.03516	65,757	2,312	64,601	777,796	11.83
75–76	.03818	63,445	2,422	62,234	713,195	11.24
76–77	.04146	61,023	2,530	59,758	650,961	10.67
77–78	.04501	58,493	2,633	57,176	591,203	10.11
78–79	.04900	55,860	2,738	54,491	534,027	9.56
79–80	.05357	53,122	2,846	51,699	479,536	9.03
80–81	.05879	50,276	2,955	48,799	427,837	8.51
81–82	.06460	47,321	3,057	45,792	379,038	8.01
82–83	.07109	44,264	3,147	42,691	333,246	7.53
83–84	.07820	41,117	3,215	39,509	290,555	7.07
84–85	.08604	37,902	3,261	36,271	251,046	6.62
85–86	.09556	34,641	3,311	32,986	214,775	6.20
86–87	.10649	31,330	3,336	29,662	181,789	5.80
87–88	.11784	27,994	3,299	26,345	152,127	5.43
88–89	.12900	24,695	3,185	23,102	125,782	5.09
89–90	.14015	21,510	3,015	20,002	102,680	4.77
90–91	.15247	18,495	2,820	17,085	82,678	4.47
91–92	.16654	15,675	2,610	14,370	65,593	4.18
92–93	.18140	13,065	2,370	11,880	51,223	3.92
93–94	.19653	10,695	2,102	9,644	39,343	3.68
94–95	.21184	8,593	1,821	7,682	29,699	3.46
95–96	.22760	6,772	1,541	6,002	22,017	3.25
96–97	.24414	5,231	1,277	4,592	16,015	3.06
97–98	.26009	3,954	1,028	3,440	11,423	2.89
98–99	.27538	2,926	806	2,523	7,983	2.73
99–100	.29135	2,120	618	1,811	5,460	2.58
100–101	.30824	1,502	463	1,271	3,649	2.43
101–102	.32612	1,039	339	870	2,378	2.29
102–103	.34504	700	241	579	1,508	2.15
103–104	.36505	459	168	375	929	2.03
104–105	.38622	291	112	235	554	1.90
105–106	.40862	179	73	142	319	1.78
106–107	.43232	106	46	83	177	1.67
107–108	.45740	60	27	46	94	1.56
108–109	.48393	33	16	25	48	1.46
109–110	.51200	17	9	13	23	1.36

**Table 5. Life table for white males: New Mexico, 1989–91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0-1	.00919	100,000	919	99,262	7,265,823	72.66
1-2	.00079	99,081	79	99,041	7,166,561	72.33
2-3	.00059	99,002	58	98,973	7,067,520	71.39
3-4	.00048	98,944	48	98,921	6,968,547	70.43
4-5	.00039	98,896	38	98,877	6,869,626	69.46
5-6	.00033	98,858	33	98,841	6,770,749	68.49
6-7	.00031	98,825	31	98,809	6,671,908	67.51
7-8	.00029	98,794	28	98,780	6,573,099	66.53
8-9	.00026	98,766	26	98,753	6,474,319	65.55
9-10	.00021	98,740	20	98,730	6,375,566	64.57
10-11	.00017	98,720	17	98,712	6,276,836	63.58
11-12	.00017	98,703	16	98,694	6,178,124	62.59
12-13	.00026	98,687	26	98,674	6,079,430	61.60
13-14	.00047	98,661	47	98,638	5,980,756	60.62
14-15	.00077	98,614	76	98,576	5,882,118	59.65
15-16	.00111	98,538	109	98,484	5,783,542	58.69
16-17	.00143	98,429	140	98,359	5,685,058	57.76
17-18	.00169	98,289	166	98,205	5,586,699	56.84
18-19	.00186	98,123	183	98,032	5,488,494	55.93
19-20	.00197	97,940	194	97,843	5,390,462	55.04
20-21	.00208	97,746	203	97,645	5,292,619	54.15
21-22	.00220	97,543	214	97,436	5,194,974	53.26
22-23	.00228	97,329	222	97,217	5,097,538	52.37
23-24	.00230	97,107	223	96,996	5,000,321	51.49
24-25	.00228	96,884	221	96,773	4,903,325	50.61
25-26	.00225	96,663	217	96,554	4,806,552	49.73
26-27	.00222	96,446	214	96,339	4,709,998	48.84
27-28	.00222	96,232	214	96,125	4,613,659	47.94
28-29	.00229	96,018	220	95,908	4,517,534	47.05
29-30	.00239	95,798	229	95,684	4,421,626	46.16
30-31	.00250	95,569	239	95,450	4,325,942	45.26
31-32	.00260	95,330	247	95,206	4,230,492	44.38
32-33	.00267	95,083	254	94,956	4,135,286	43.49
33-34	.00271	94,829	257	94,700	4,040,330	42.61
34-35	.00272	94,572	257	94,443	3,945,630	41.72
35-36	.00273	94,315	258	94,186	3,851,187	40.83
36-37	.00276	94,057	259	93,927	3,757,001	39.94
37-38	.00280	93,798	263	93,666	3,663,074	39.05
38-39	.00288	93,535	270	93,400	3,569,408	38.16
39-40	.00298	93,265	277	93,127	3,476,008	37.27
40-41	.00309	92,988	288	92,844	3,382,881	36.38
41-42	.00321	92,700	297	92,552	3,290,037	35.49
42-43	.00334	92,403	309	92,248	3,197,485	34.60
43-44	.00350	92,094	322	91,933	3,105,237	33.72
44-45	.00369	91,772	338	91,603	3,013,304	32.83
45-46	.00393	91,434	360	91,254	2,921,701	31.95
46-47	.00423	91,074	385	90,881	2,830,447	31.08
47-48	.00452	90,689	410	90,484	2,739,566	30.21
48-49	.00478	90,279	432	90,063	2,649,082	29.34
49-50	.00501	89,847	449	89,622	2,559,019	28.48
50-51	.00525	89,398	469	89,163	2,469,397	27.62
51-52	.00558	88,929	497	88,681	2,380,234	26.77
52-53	.00608	88,432	537	88,164	2,291,553	25.91
53-54	.00679	87,895	596	87,597	2,203,389	25.07
54-55	.00767	87,299	670	86,963	2,115,792	24.24

**Table 5. Life table for white males: New Mexico, 1989–91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55–56	.00863	86,629	748	86,255	2,028,829	23.42
56–57	.00961	85,881	826	85,469	1,942,574	22.62
57–58	.01062	85,055	903	84,603	1,857,105	21.83
58–59	.01162	84,152	978	83,664	1,772,502	21.06
59–60	.01265	83,174	1,052	82,648	1,688,838	20.30
60–61	.01372	82,122	1,127	81,558	1,606,190	19.56
61–62	.01485	80,995	1,202	80,394	1,524,632	18.82
62–63	.01603	79,793	1,279	79,153	1,444,238	18.10
63–64	.01729	78,514	1,358	77,835	1,365,085	17.39
64–65	.01862	77,156	1,436	76,438	1,287,250	16.68
65–66	.01995	75,720	1,511	74,965	1,210,812	15.99
66–67	.02137	74,209	1,586	73,416	1,135,847	15.31
67–68	.02308	72,623	1,676	71,785	1,062,431	14.63
68–69	.02522	70,947	1,789	70,052	990,646	13.96
69–70	.02785	69,158	1,927	68,194	920,594	13.31
70–71	.03091	67,231	2,078	66,193	852,400	12.68
71–72	.03425	65,153	2,232	64,037	786,207	12.07
72–73	.03781	62,921	2,379	61,732	722,170	11.48
73–74	.04136	60,542	2,503	59,291	660,438	10.91
74–75	.04487	58,039	2,605	56,736	601,147	10.36
75–76	.04856	55,434	2,692	54,088	544,411	9.82
76–77	.05267	52,742	2,778	51,353	490,323	9.30
77–78	.05723	49,964	2,859	48,535	438,970	8.79
78–79	.06248	47,105	2,943	45,633	390,435	8.29
79–80	.06855	44,162	3,028	42,648	344,802	7.81
80–81	.07572	41,134	3,114	39,577	302,154	7.35
81–82	.08377	38,020	3,185	36,428	262,577	6.91
82–83	.09225	34,835	3,213	33,228	226,149	6.49
83–84	.10057	31,622	3,181	30,032	192,921	6.10
84–85	.10878	28,441	3,094	26,894	162,889	5.73
85–86	.11897	25,347	3,015	23,839	135,995	5.37
86–87	.13113	22,332	2,929	20,868	112,156	5.02
87–88	.14403	19,403	2,794	18,006	91,288	4.70
88–89	.15696	16,609	2,607	15,305	73,282	4.41
89–90	.16985	14,002	2,378	12,813	57,977	4.14
90–91	.18329	11,624	2,131	10,558	45,164	3.89
91–92	.19822	9,493	1,882	8,552	34,606	3.65
92–93	.21437	7,611	1,631	6,796	26,054	3.42
93–94	.23140	5,980	1,384	5,288	19,258	3.22
94–95	.24804	4,596	1,140	4,026	13,970	3.04
95–96	.26329	3,456	910	3,001	9,944	2.88
96–97	.27914	2,546	711	2,191	6,943	2.73
97–98	.29399	1,835	539	1,565	4,752	2.59
98–99	.30869	1,296	400	1,096	3,187	2.46
99–100	.32413	896	291	751	2,091	2.33
100–101	.34033	605	206	502	1,340	2.21
101–102	.35735	399	142	328	838	2.10
102–103	.37522	257	97	208	510	1.99
103–104	.39398	160	63	129	302	1.88
104–105	.41368	97	40	77	173	1.78
105–106	.43436	57	25	45	96	1.68
106–107	.45608	32	14	25	51	1.58
107–108	.47888	18	9	13	26	1.49
108–109	.50282	9	4	7	13	1.41
109–110	.52797	5	3	3	6	1.32

**Table 6. Life table for white females: New Mexico, 1989–91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0-1	.00702	100,000	702	99,455	7,953,314	79.53
1-2	.00061	99,298	61	99,268	7,853,859	79.09
2-3	.00045	99,237	45	99,214	7,754,591	78.14
3-4	.00035	99,192	35	99,175	7,655,377	77.18
4-5	.00028	99,157	28	99,143	7,556,202	76.20
5-6	.00026	99,129	26	99,116	7,457,059	75.23
6-7	.00024	99,103	23	99,092	7,357,943	74.25
7-8	.00022	99,080	22	99,069	7,258,851	73.26
8-9	.00019	99,058	19	99,049	7,159,782	72.28
9-10	.00017	99,039	17	99,031	7,060,733	71.29
10-11	.00015	99,022	15	99,014	6,961,702	70.30
11-12	.00015	99,007	15	98,999	6,862,688	69.32
12-13	.00018	98,992	18	98,984	6,763,689	68.33
13-14	.00024	98,974	23	98,962	6,664,705	67.34
14-15	.00032	98,951	32	98,935	6,565,743	66.35
15-16	.00042	98,919	42	98,898	6,466,808	65.37
16-17	.00052	98,877	51	98,851	6,367,910	64.40
17-18	.00060	98,826	60	98,796	6,269,059	63.44
18-19	.00066	98,766	65	98,734	6,170,263	62.47
19-20	.00070	98,701	69	98,666	6,071,529	61.51
20-21	.00074	98,632	73	98,596	5,972,863	60.56
21-22	.00078	98,559	77	98,520	5,874,267	59.60
22-23	.00080	98,482	79	98,443	5,775,747	58.65
23-24	.00080	98,403	79	98,363	5,677,304	57.69
24-25	.00078	98,324	77	98,285	5,578,941	56.74
25-26	.00076	98,247	75	98,209	5,480,656	55.78
26-27	.00074	98,172	73	98,136	5,382,447	54.83
27-28	.00073	98,099	72	98,062	5,284,311	53.87
28-29	.00073	98,027	72	97,991	5,186,249	52.91
29-30	.00074	97,955	73	97,919	5,088,258	51.94
30-31	.00075	97,882	72	97,846	4,990,339	50.98
31-32	.00076	97,810	74	97,772	4,892,493	50.02
32-33	.00078	97,736	77	97,698	4,794,721	49.06
33-34	.00084	97,659	82	97,618	4,697,023	48.10
34-35	.00091	97,577	89	97,533	4,599,405	47.14
35-36	.00100	97,488	97	97,439	4,501,872	46.18
36-37	.00109	97,391	106	97,339	4,404,433	45.22
37-38	.00116	97,285	113	97,228	4,307,094	44.27
38-39	.00122	97,172	119	97,113	4,209,866	43.32
39-40	.00127	97,053	123	96,992	4,112,753	42.38
40-41	.00132	96,930	128	96,866	4,015,761	41.43
41-42	.00138	96,802	134	96,735	3,918,895	40.48
42-43	.00148	96,668	142	96,598	3,822,160	39.54
43-44	.00162	96,526	157	96,447	3,725,562	38.60
44-45	.00181	96,369	174	96,283	3,629,115	37.66
45-46	.00204	96,195	196	96,097	3,532,832	36.73
46-47	.00230	95,999	221	95,888	3,436,735	35.80
47-48	.00255	95,778	244	95,656	3,340,847	34.88
48-49	.00277	95,534	265	95,401	3,245,191	33.97
49-50	.00296	95,269	282	95,128	3,149,790	33.06
50-51	.00316	94,987	300	94,837	3,054,662	32.16
51-52	.00342	94,687	323	94,525	2,959,825	31.26
52-53	.00368	94,364	348	94,190	2,865,300	30.36
53-54	.00397	94,016	373	93,830	2,771,110	29.47
54-55	.00428	93,643	401	93,442	2,677,280	28.5

**Table 6. Life table for white females: New Mexico, 1989–91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55–56	.00460	93,242	429	93,028	2,583,838	27.71
56–57	.00498	92,813	463	92,582	2,490,810	26.84
57–58	.00548	92,350	506	92,097	2,398,228	25.97
58–59	.00612	91,844	562	91,563	2,306,131	25.11
59–60	.00687	91,282	627	90,969	2,214,568	24.26
60–61	.00766	90,655	694	90,308	2,123,599	23.43
61–62	.00846	89,961	761	89,580	2,033,291	22.60
62–63	.00932	89,200	832	88,783	1,943,711	21.79
63–64	.01026	88,368	907	87,915	1,854,928	20.99
64–65	.01126	87,461	984	86,969	1,767,013	20.20
65–66	.01232	86,477	1,066	85,944	1,680,044	19.43
66–67	.01342	85,411	1,145	84,838	1,594,100	18.66
67–68	.01454	84,266	1,226	83,653	1,509,262	17.91
68–69	.01573	83,040	1,306	82,388	1,425,609	17.17
69–70	.01706	81,734	1,394	81,037	1,343,221	16.43
70–71	.01855	80,340	1,490	79,595	1,262,184	15.71
71–72	.02027	78,850	1,599	78,050	1,182,589	15.00
72–73	.02233	77,251	1,725	76,389	1,104,539	14.30
73–74	.02471	75,526	1,867	74,593	1,028,150	13.61
74–75	.02730	73,659	2,011	72,653	953,557	12.95
75–76	.03005	71,648	2,153	70,572	880,904	12.29
76–77	.03296	69,495	2,290	68,350	810,332	11.66
77–78	.03603	67,205	2,422	65,994	741,982	11.04
78–79	.03936	64,783	2,550	63,508	675,988	10.43
79–80	.04312	62,233	2,683	60,892	612,480	9.84
80–81	.04730	59,550	2,817	58,141	551,588	9.26
81–82	.05201	56,733	2,950	55,258	493,447	8.70
82–83	.05765	53,783	3,101	52,233	438,189	8.15
83–84	.06451	50,682	3,269	49,047	385,956	7.62
84–85	.07266	47,413	3,445	45,690	336,909	7.11
85–86	.08268	43,968	3,635	42,150	291,219	6.62
86–87	.09389	40,333	3,787	38,440	249,069	6.18
87–88	.10542	36,546	3,853	34,619	210,629	5.76
88–89	.11661	32,693	3,812	30,787	176,010	5.38
89–90	.12784	28,881	3,692	27,035	145,223	5.03
90–91	.14058	25,189	3,541	23,418	118,188	4.69
91–92	.15525	21,648	3,361	19,967	94,770	4.38
92–93	.17049	18,287	3,118	16,728	74,803	4.09
93–94	.18567	15,169	2,816	13,761	58,075	3.83
94–95	.20107	12,353	2,484	11,111	44,314	3.59
95–96	.21737	9,869	2,145	8,796	33,203	3.36
96–97	.23434	7,724	1,810	6,819	24,407	3.16
97–98	.25091	5,914	1,484	5,172	17,588	2.97
98–99	.26715	4,430	1,184	3,838	12,416	2.80
99–100	.28318	3,246	919	2,787	8,578	2.64
100–101	.30017	2,327	698	1,978	5,791	2.49
101–102	.31818	1,629	519	1,369	3,813	2.34
102–103	.33727	1,110	374	923	2,444	2.20
103–104	.35750	736	263	605	1,521	2.07
104–105	.37895	473	179	383	916	1.94
105–106	.40169	294	118	235	533	1.81
106–107	.42579	176	75	138	298	1.70
107–108	.45134	101	46	78	160	1.59
108–109	.47842	55	26	42	82	1.48
109–110	.50712	29	15	22	40	1.38

**Table 7. Life table for the population other than white: New Mexico, 1989-91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0-1	.01056	100,000	1,056	99,218	7,341,338	73.41
1-2	.00117	98,944	115	98,886	7,242,120	73.19
2-3	.00079	98,829	79	98,789	7,143,234	72.28
3-4	.00060	98,750	59	98,721	7,044,445	71.34
4-5	.00046	98,691	45	98,669	6,945,724	70.38
5-6	.00040	98,646	40	98,626	6,847,055	69.41
6-7	.00036	98,606	35	98,589	6,748,429	68.44
7-8	.00033	98,571	33	98,554	6,649,840	67.46
8-9	.00028	98,538	27	98,525	6,551,286	66.48
9-10	.00023	98,511	23	98,499	6,452,761	65.50
10-11	.00019	98,488	18	98,479	6,354,262	64.52
11-12	.00019	98,470	20	98,460	6,255,783	63.53
12-13	.00030	98,450	29	98,436	6,157,323	62.54
13-14	.00053	98,421	52	98,395	6,058,887	61.56
14-15	.00085	98,369	84	98,327	5,960,492	60.59
15-16	.00123	98,285	120	98,226	5,862,165	59.64
16-17	.00159	98,165	156	98,086	5,763,939	58.72
17-18	.00190	98,009	186	97,917	5,665,853	57.81
18-19	.00213	97,823	208	97,719	5,567,936	56.92
19-20	.00229	97,615	224	97,503	5,470,217	56.04
20-21	.00245	97,391	238	97,272	5,372,714	55.17
21-22	.00262	97,153	255	97,025	5,275,442	54.30
22-23	.00273	96,898	264	96,766	5,178,417	53.44
23-24	.00275	96,634	266	96,502	5,081,651	52.59
24-25	.00272	96,368	262	96,237	4,985,149	51.73
25-26	.00266	96,106	255	95,979	4,888,912	50.87
26-27	.00260	95,851	250	95,726	4,792,933	50.00
27-28	.00259	95,601	247	95,478	4,697,207	49.13
28-29	.00263	95,354	250	95,229	4,601,729	48.26
29-30	.00271	95,104	258	94,975	4,506,500	47.39
30-31	.00280	94,846	266	94,713	4,411,525	46.51
31-32	.00289	94,580	273	94,443	4,316,812	45.64
32-33	.00297	94,307	280	94,167	4,222,369	44.77
33-34	.00305	94,027	287	93,883	4,128,202	43.90
34-35	.00312	93,740	293	93,594	4,034,319	43.04
35-36	.00320	93,447	299	93,298	3,940,725	42.17
36-37	.00330	93,148	307	92,994	3,847,427	41.30
37-38	.00342	92,841	317	92,682	3,754,433	40.44
38-39	.00356	92,524	330	92,359	3,661,751	39.58
39-40	.00373	92,194	344	92,022	3,569,392	38.72
40-41	.00393	91,850	361	91,670	3,477,370	37.86
41-42	.00414	91,489	378	91,299	3,385,700	37.01
42-43	.00436	91,111	398	90,912	3,294,401	36.16
43-44	.00459	90,713	416	90,505	3,203,489	35.31
44-45	.00484	90,297	438	90,078	3,112,984	34.48
45-46	.00513	89,859	460	89,629	3,022,906	33.64
46-47	.00545	89,399	488	89,155	2,933,277	32.81
47-48	.00580	88,911	515	88,654	2,844,122	31.99
48-49	.00613	88,396	543	88,124	2,755,468	31.17
49-50	.00645	87,853	566	87,570	2,667,344	30.36
50-51	.00678	87,287	592	86,991	2,579,774	29.56
51-52	.00712	86,695	617	86,387	2,492,783	28.75
52-53	.00746	86,078	643	85,756	2,406,396	27.96
53-54	.00779	85,435	665	85,103	2,320,640	27.16
54-55	.00812	84,770	689	84,426	2,235,537	26.37



**Table 7. Life table for the population other than white: New Mexico, 1989–91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55–56	.00845	84,081	710	83,726	2,151,111	25.58
56–57	.00882	83,371	735	83,003	2,067,385	24.80
57–58	.00932	82,636	771	82,250	1,984,382	24.01
58–59	.01002	81,865	821	81,455	1,902,132	23.23
59–60	.01091	81,044	884	80,603	1,820,677	22.47
60–61	.01190	80,160	953	79,683	1,740,074	21.71
61–62	.01294	79,207	1,025	78,694	1,660,391	20.96
62–63	.01408	78,182	1,101	77,632	1,581,697	20.23
63–64	.01534	77,081	1,182	76,490	1,504,065	19.51
64–65	.01670	75,899	1,268	75,265	1,427,575	18.81
65–66	.01822	74,631	1,360	73,951	1,352,310	18.12
66–67	.01986	73,271	1,455	72,543	1,278,359	17.45
67–68	.02150	71,816	1,544	71,044	1,205,816	16.79
68–69	.02307	70,272	1,622	69,461	1,134,772	16.15
69–70	.02459	68,650	1,688	67,806	1,065,311	15.52
70–71	.02618	66,962	1,753	66,086	997,505	14.90
71–72	.02792	65,209	1,821	64,298	931,419	14.28
72–73	.02973	63,388	1,885	62,446	867,121	13.68
73–74	.03160	61,503	1,943	60,532	804,675	13.08
74–75	.03353	59,560	1,997	58,561	744,143	12.49
75–76	.03539	57,563	2,037	56,544	685,582	11.91
76–77	.03738	55,526	2,076	54,488	629,038	11.33
77–78	.03999	53,450	2,138	52,381	574,550	10.75
78–79	.04368	51,312	2,241	50,192	522,169	10.18
79–80	.04854	49,071	2,382	47,880	471,977	9.62
80–81	.05460	46,689	2,549	45,415	424,097	9.08
81–82	.06137	44,140	2,709	42,785	378,682	8.58
82–83	.06829	41,431	2,829	40,017	335,897	8.11
83–84	.07426	38,602	2,867	37,168	295,880	7.66
84–85	.07907	35,735	2,825	34,323	258,712	7.24
85–86	.08496	32,910	2,796	31,512	224,389	6.82
86–87	.09191	30,114	2,768	28,730	192,877	6.40
87–88	.10003	27,346	2,735	25,978	164,147	6.00
88–89	.11018	24,611	2,712	23,255	138,169	5.61
89–90	.12259	21,899	2,685	20,557	114,914	5.25
90–91	.13773	19,214	2,646	17,891	94,357	4.91
91–92	.15471	16,568	2,563	15,286	76,466	4.62
92–93	.17098	14,005	2,395	12,808	61,180	4.37
93–94	.18159	11,610	2,108	10,556	48,372	4.17
94–95	.18738	9,502	1,780	8,612	37,816	3.98
95–96	.19586	7,722	1,513	6,965	29,204	3.78
96–97	.20830	6,209	1,293	5,563	22,239	3.58
97–98	.22089	4,916	1,086	4,373	16,676	3.39
98–99	.23370	3,830	895	3,382	12,303	3.21
99–100	.24726	2,935	726	2,572	8,921	3.04
100–101	.26160	2,209	578	1,920	6,349	2.87
101–102	.27677	1,631	451	1,406	4,429	2.71
102–103	.29282	1,180	346	1,007	3,023	2.56
103–104	.30981	834	258	705	2,016	2.42
104–105	.32778	576	189	482	1,311	2.28
105–106	.34679	387	134	320	829	2.14
106–107	.36690	253	93	206	509	2.01
107–108	.38818	160	62	129	303	1.89
108–109	.41070	98	40	78	174	1.78
109–110	.43452	58	25	45	96	1.66

**Table 8. Life table for males other than white: New Mexico, 1989–91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0-1	.01259	100,000	1,259	99,063	6,896,501	68.97
1-2	.00127	98,741	126	98,677	6,797,438	68.84
2-3	.00093	98,615	91	98,570	6,698,761	67.93
3-4	.00070	98,524	70	98,489	6,600,191	66.99
4-5	.00055	98,454	54	98,427	6,501,702	66.04
5-6	.00047	98,400	46	98,377	6,403,275	65.07
6-7	.00044	98,354	43	98,333	6,304,898	64.10
7-8	.00040	98,311	40	98,290	6,206,565	63.13
8-9	.00035	98,271	34	98,254	6,108,275	62.16
9-10	.00027	98,237	27	98,224	6,010,021	61.18
10-11	.00021	98,210	21	98,199	5,911,797	60.20
11-12	.00022	98,189	21	98,179	5,813,598	59.21
12-13	.00039	98,168	39	98,149	5,715,419	58.22
13-14	.00077	98,129	75	98,092	5,617,270	57.24
14-15	.00130	98,054	127	97,990	5,519,178	56.29
15-16	.00191	97,927	187	97,833	5,421,188	55.36
16-17	.00250	97,740	244	97,618	5,323,355	54.46
17-18	.00299	97,496	292	97,350	5,225,737	53.60
18-19	.00335	97,204	325	97,042	5,128,387	52.76
19-20	.00358	96,879	347	96,705	5,031,345	51.93
20-21	.00381	96,532	368	96,348	4,934,640	51.12
21-22	.00404	96,164	388	95,970	4,838,292	50.31
22-23	.00418	95,776	400	95,576	4,742,322	49.51
23-24	.00420	95,376	401	95,175	4,646,746	48.72
24-25	.00414	94,975	393	94,779	4,551,571	47.92
25-26	.00402	94,582	380	94,392	4,456,792	47.12
26-27	.00393	94,202	370	94,016	4,362,400	46.31
27-28	.00389	93,832	365	93,650	4,268,384	45.49
28-29	.00395	93,467	370	93,282	4,174,734	44.67
29-30	.00409	93,097	380	92,907	4,081,452	43.84
30-31	.00425	92,717	394	92,520	3,988,545	43.02
31-32	.00439	92,323	405	92,120	3,896,025	42.20
32-33	.00451	91,918	415	91,711	3,803,905	41.38
33-34	.00460	91,503	421	91,292	3,712,194	40.57
34-35	.00467	91,082	425	90,870	3,620,902	39.75
35-36	.00473	90,657	428	90,443	3,530,032	38.94
36-37	.00482	90,229	435	90,011	3,439,589	38.12
37-38	.00494	89,794	444	89,572	3,349,578	37.30
38-39	.00513	89,350	458	89,121	3,260,006	36.49
39-40	.00536	88,892	477	88,653	3,170,885	35.67
40-41	.00564	88,415	499	88,166	3,082,232	34.86
41-42	.00594	87,916	522	87,655	2,994,066	34.06
42-43	.00623	87,394	544	87,122	2,906,411	33.26
43-44	.00651	86,850	566	86,567	2,819,289	32.46
44-45	.00678	86,284	585	85,992	2,732,722	31.67
45-46	.00707	85,699	606	85,396	2,646,730	30.88
46-47	.00742	85,093	631	84,777	2,561,334	30.10
47-48	.00782	84,462	661	84,132	2,476,557	29.32
48-49	.00829	83,801	694	83,454	2,392,425	28.55
49-50	.00880	83,107	732	82,741	2,308,971	27.78
50-51	.00939	82,375	773	81,989	2,226,230	27.03
51-52	.00998	81,602	815	81,195	2,144,241	26.28
52-53	.01051	80,787	849	80,362	2,063,046	25.54
53-54	.01090	79,938	871	79,503	1,982,684	24.80
54-55	.01120	79,067	886	78,624	1,903,181	24.07

**Table 8. Life table for males other than white: New Mexico, 1989-91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55-56	.01144	78,181	894	77,734	1,824,557	23.34
56-57	.01174	77,287	907	76,834	1,746,823	22.60
57-58	.01224	76,380	935	75,913	1,669,989	21.86
58-59	.01307	75,445	986	74,952	1,594,076	21.13
59-60	.01418	74,459	1,055	73,931	1,519,124	20.40
60-61	.01546	73,404	1,135	72,837	1,445,193	19.69
61-62	.01677	72,269	1,212	71,663	1,372,356	18.99
62-63	.01814	71,057	1,289	70,412	1,300,693	18.30
63-64	.01952	69,768	1,361	69,088	1,230,281	17.63
64-65	.02091	68,407	1,431	67,691	1,161,193	16.97
65-66	.02240	66,976	1,500	66,226	1,093,502	16.33
66-67	.02402	65,476	1,572	64,690	1,027,276	15.69
67-68	.02573	63,904	1,645	63,082	962,586	15.06
68-69	.02755	62,259	1,715	61,402	899,504	14.45
69-70	.02948	60,544	1,785	59,651	838,102	13.84
70-71	.03160	58,759	1,856	57,832	778,451	13.25
71-72	.03388	56,903	1,928	55,938	720,619	12.66
72-73	.03621	54,975	1,991	53,979	664,681	12.09
73-74	.03853	52,984	2,042	51,963	610,702	11.53
74-75	.04090	50,942	2,083	49,900	558,739	10.97
75-76	.04313	48,859	2,108	47,805	508,839	10.41
76-77	.04566	46,751	2,135	45,684	461,034	9.86
77-78	.04940	44,616	2,204	43,514	415,350	9.31
78-79	.05514	42,412	2,338	41,243	371,836	8.77
79-80	.06293	40,074	2,522	38,813	330,593	8.25
80-81	.07298	37,552	2,741	36,182	291,780	7.77
81-82	.08410	34,811	2,927	33,347	255,598	7.34
82-83	.09430	31,884	3,007	30,381	222,251	6.97
83-84	.10088	28,877	2,913	27,420	191,870	6.64
84-85	.10357	25,964	2,689	24,620	164,450	6.33
85-86	.10599	23,275	2,467	22,041	139,830	6.01
86-87	.11028	20,808	2,295	19,661	117,789	5.66
87-88	.11722	18,513	2,170	17,428	98,128	5.30
88-89	.12842	16,343	2,099	15,294	80,700	4.94
89-90	.14373	14,244	2,047	13,220	65,406	4.59
90-91	.16278	12,197	1,985	11,205	52,186	4.28
91-92	.18354	10,212	1,875	9,274	40,981	4.01
92-93	.20275	8,337	1,690	7,492	31,707	3.80
93-94	.21368	6,647	1,420	5,937	24,215	3.64
94-95	.21885	5,227	1,144	4,655	18,278	3.50
95-96	.22903	4,083	935	3,615	13,623	3.34
96-97	.24048	3,148	757	2,770	10,008	3.18
97-98	.25250	2,391	604	2,089	7,238	3.03
98-99	.26513	1,787	474	1,550	5,149	2.88
99-100	.27838	1,313	365	1,130	3,599	2.74
100-101	.29230	948	277	809	2,469	2.61
101-102	.30692	671	206	568	1,660	2.47
102-103	.32226	465	150	390	1,092	2.35
103-104	.33837	315	107	262	702	2.23
104-105	.35529	208	74	171	440	2.11
105-106	.37306	134	50	110	269	2.00
106-107	.39171	84	33	67	159	1.89
107-108	.41130	51	21	41	92	1.79
108-109	.43186	30	13	24	51	1.69
109-110	.45345	17	8	13	27	1.59

**Table 9. Life table for females other than white: New Mexico, 1989-91**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
0-1	.00851	100,000	851	99,373	7,792,696	77.93
1-2	.00106	99,149	105	99,096	7,693,323	77.59
2-3	.00066	99,044	65	99,012	7,594,227	76.68
3-4	.00049	98,979	48	98,954	7,495,215	75.73
4-5	.00036	98,931	36	98,913	7,396,261	74.76
5-6	.00034	98,895	33	98,878	7,297,348	73.79
6-7	.00029	98,862	29	98,848	7,198,470	72.81
7-8	.00025	98,833	24	98,821	7,099,622	71.83
8-9	.00021	98,809	22	98,798	7,000,801	70.85
9-10	.00019	98,787	18	98,778	6,902,003	69.87
10-11	.00017	98,769	16	98,761	6,803,225	68.88
11-12	.00017	98,753	17	98,744	6,704,464	67.89
12-13	.00020	98,736	20	98,727	6,605,720	66.90
13-14	.00029	98,716	28	98,702	6,506,993	65.92
14-15	.00040	98,688	40	98,668	6,408,291	64.93
15-16	.00054	98,648	53	98,621	6,309,623	63.96
16-17	.00068	98,595	67	98,562	6,211,002	63.00
17-18	.00080	98,528	79	98,488	6,112,440	62.04
18-19	.00090	98,449	88	98,405	6,013,952	61.09
19-20	.00098	98,361	96	98,313	5,915,547	60.14
20-21	.00106	98,265	105	98,212	5,817,234	59.20
21-22	.00115	98,160	113	98,104	5,719,022	58.26
22-23	.00122	98,047	120	97,986	5,620,918	57.33
23-24	.00126	97,927	124	97,865	5,522,932	56.40
24-25	.00127	97,803	124	97,742	5,425,067	55.47
25-26	.00127	97,679	124	97,617	5,327,325	54.54
26-27	.00127	97,555	124	97,492	5,229,708	53.61
27-28	.00129	97,431	126	97,368	5,132,216	52.68
28-29	.00132	97,305	128	97,241	5,034,848	51.74
29-30	.00137	97,177	133	97,110	4,937,607	50.81
30-31	.00142	97,044	138	96,975	4,840,497	49.88
31-32	.00147	96,906	143	96,834	4,743,522	48.95
32-33	.00153	96,763	148	96,690	4,646,688	48.02
33-34	.00160	96,615	154	96,538	4,549,998	47.09
34-35	.00169	96,461	163	96,379	4,453,460	46.17
35-36	.00178	96,298	171	96,213	4,357,081	45.25
36-37	.00189	96,127	182	96,036	4,260,868	44.33
37-38	.00200	95,945	191	95,849	4,164,832	43.41
38-39	.00212	95,754	203	95,653	4,068,983	42.49
39-40	.00224	95,551	214	95,443	3,973,330	41.58
40-41	.00238	95,337	227	95,224	3,877,887	40.68
41-42	.00253	95,110	241	94,990	3,782,663	39.77
42-43	.00270	94,869	256	94,741	3,687,673	38.87
43-44	.00289	94,613	273	94,476	3,592,932	37.97
44-45	.00311	94,340	294	94,193	3,498,456	37.08
45-46	.00337	94,046	316	93,888	3,404,263	36.20
46-47	.00367	93,730	344	93,558	3,310,375	35.32
47-48	.00396	93,386	370	93,201	3,216,817	34.45
48-49	.00421	93,016	391	92,821	3,123,616	33.58
49-50	.00440	92,625	408	92,421	3,030,795	32.72
50-51	.00459	92,217	423	92,005	2,938,374	31.86
51-52	.00480	91,794	440	91,574	2,846,369	31.01
52-53	.00504	91,354	461	91,123	2,754,795	30.16
53-54	.00535	90,893	486	90,650	2,663,672	29.31
54-55	.00571	90,407	517	90,149	2,573,022	28.46

**Table 9. Life table for females other than white: New Mexico, 1989–91—Con.**

Age in years	Proportion dying	Of 100,000 born alive		Stationary population		Average remaining lifetime
		Number living at beginning of year of age (3)	Number dying during year of age (4)	In year of age (5)	In this year of age and all subsequent years (6)	Average number of years of life remaining at beginning of year of age (7)
Period of life between two exact ages stated (1)	Proportion of persons alive at beginning of year of age dying during year (2)	$l_x$	$d_x$	$L_x$	$T_x$	${}^o e_x$
x to x+1	$q_x$					
55–56	.00610	89,890	548	89,616	2,482,873	27.62
56–57	.00652	89,342	582	89,051	2,393,257	26.79
57–58	.00702	88,760	624	88,448	2,304,206	25.96
58–59	.00764	88,136	673	87,799	2,215,758	25.14
59–60	.00836	87,463	732	87,098	2,127,959	24.33
60–61	.00915	86,731	793	86,334	2,040,861	23.53
61–62	.01000	85,938	859	85,508	1,954,527	22.74
62–63	.01097	85,079	934	84,612	1,869,019	21.97
63–64	.01211	84,145	1,019	83,636	1,784,407	21.21
64–65	.01342	83,126	1,116	82,568	1,700,771	20.46
65–66	.01492	82,010	1,223	81,398	1,618,203	19.73
66–67	.01652	80,787	1,334	80,120	1,536,805	19.02
67–68	.01806	79,453	1,435	78,735	1,456,685	18.33
68–69	.01940	78,018	1,513	77,262	1,377,950	17.66
69–70	.02057	76,505	1,574	75,718	1,300,688	17.00
70–71	.02172	74,931	1,627	74,117	1,224,970	16.35
71–72	.02301	73,304	1,687	72,460	1,150,853	15.70
72–73	.02441	71,617	1,748	70,743	1,078,393	15.06
73–74	.02597	69,869	1,815	68,962	1,007,650	14.42
74–75	.02765	68,054	1,881	67,114	938,688	13.79
75–76	.02935	66,173	1,942	65,202	871,574	13.17
76–77	.03106	64,231	1,995	63,233	806,372	12.55
77–78	.03295	62,236	2,051	61,211	743,139	11.94
78–79	.03520	60,185	2,118	59,126	681,928	11.33
79–80	.03794	58,067	2,203	56,966	622,802	10.73
80–81	.04111	55,864	2,297	54,715	565,836	10.13
81–82	.04478	53,567	2,398	52,368	511,121	9.54
82–83	.04938	51,169	2,527	49,906	458,753	8.97
83–84	.05498	48,642	2,674	47,305	408,847	8.41
84–85	.06135	45,968	2,820	44,557	361,542	7.87
85–86	.07018	43,148	3,028	41,634	316,985	7.35
86–87	.07960	40,120	3,194	38,523	275,351	6.86
87–88	.08923	36,926	3,295	35,278	236,828	6.41
88–89	.09937	33,631	3,342	31,960	201,550	5.99
89–90	.11063	30,289	3,351	28,614	169,590	5.60
90–91	.12419	26,938	3,345	25,266	140,976	5.23
91–92	.14001	23,593	3,303	21,941	115,710	4.90
92–93	.15591	20,290	3,164	18,708	93,769	4.62
93–94	.16780	17,126	2,873	15,690	75,061	4.38
94–95	.17521	14,253	2,497	13,004	59,371	4.17
95–96	.18338	11,756	2,156	10,678	46,367	3.94
96–97	.19682	9,600	1,890	8,655	35,689	3.72
97–98	.21089	7,710	1,626	6,897	27,034	3.51
98–99	.22557	6,084	1,372	5,398	20,137	3.31
99–100	.23911	4,712	1,127	4,149	14,739	3.13
100–101	.25346	3,585	909	3,131	10,590	2.95
101–102	.26866	2,676	719	2,317	7,459	2.79
102–103	.28478	1,957	557	1,678	5,142	2.63
103–104	.30187	1,400	423	1,189	3,464	2.47
104–105	.31998	977	312	821	2,275	2.33
105–106	.33918	665	226	552	1,454	2.19
106–107	.35953	439	158	360	902	2.05
107–108	.38110	281	107	228	542	1.93
108–109	.40397	174	70	139	314	1.80
109–110	.42821	104	45	81	175	1.69

**Table 10. Standard errors of the probability of dying: New Mexico, 1989–91**

Exact age in years							All other					
	Total			White			Total			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
0	.000321	.000481	.000423	.000344	.000513	.000457	.000852	.001311	.001085	*	*	*
1	.000098	.000145	.000132	.000102	.000152	.000136	.000287	.000422	.000390	*	*	*
2	.000086	.000129	.000114	.000091	.000135	.000121	.000248	.000379	.000320	*	*	*
3	.000076	.000115	.000099	.000081	.000121	.000106	.000216	.000331	.000277	*	*	*
4	.000068	.000102	.000088	.000072	.000108	.000095	.000189	.000294	.000238	*	*	*
5	.000063	.000094	.000084	.000067	.000099	.000090	.000179	.000273	.000232	*	*	*
6	.000060	.000091	.000079	.000064	.000096	.000085	.000170	.000264	.000216	*	*	*
7	.000058	.000088	.000075	.000062	.000093	.000081	.000163	.000255	.000202	*	*	*
8	.000055	.000083	.000072	.000059	.000087	.000077	.000152	.000239	.000189	*	*	*
9	.000051	.000075	.000068	.000054	.000079	.000074	.000139	.000213	.000177	*	*	*
10	.000047	.000067	.000065	.000050	.000072	.000071	.000127	.000189	.000165	*	*	*
11	.000047	.000069	.000065	.000051	.000073	.000071	.000130	.000196	.000167	*	*	*
12	.000057	.000087	.000072	.000060	.000091	.000077	.000163	.000263	.000191	*	*	*
13	.000073	.000119	.000084	.000077	.000124	.000090	.000219	.000372	.000228	*	*	*
14	.000092	.000153	.000099	.000096	.000158	.000106	.000280	.000488	.000273	*	*	*
15	.000110	.000185	.000113	.000114	.000190	.000121	.000338	.000596	.000318	*	*	*
16	.000124	.000210	.000126	.000129	.000216	.000134	.000387	.000686	.000358	*	*	*
17	.000136	.000231	.000137	.000141	.000237	.000145	.000427	.000757	.000392	*	*	*
18	.000144	.000245	.000145	.000149	.000252	.000153	.000457	.000808	.000421	*	*	*
19	.000150	.000257	.000151	.000155	.000263	.000160	.000480	.000845	.000446	*	*	*
20	.000157	.000268	.000158	.000162	.000275	.000166	.000503	.000881	.000472	*	*	*
21	.000163	.000280	.000165	.000168	.000286	.000172	.000526	.000917	.000499	*	*	*
22	.000166	.000285	.000167	.000170	.000292	.000174	.000539	.000936	.000516	*	*	*
23	.000164	.000282	.000165	.000168	.000289	.000171	.000539	.000934	.000520	*	*	*
24	.000159	.000274	.000159	.000163	.000281	.000165	.000529	.000918	.000515	*	*	*
25	.000153	.000264	.000153	.000157	.000271	.000158	.000515	.000894	.000506	*	*	*
26	.000149	.000256	.000149	.000152	.000263	.000153	.000504	.000874	.000500	*	*	*
27	.000146	.000252	.000146	.000149	.000259	.000149	.000499	.000867	.000498	*	*	*
28	.000146	.000253	.000144	.000149	.000260	.000147	.000504	.000877	.000504	*	*	*
29	.000148	.000258	.000144	.000151	.000264	.000147	.000515	.000900	.000515	*	*	*
30	.000150	.000263	.000145	.000153	.000270	.000147	.000529	.000929	.000527	*	*	*
31	.000152	.000268	.000145	.000155	.000274	.000147	.000541	.000955	.000539	*	*	*
32	.000154	.000272	.000148	.000157	.000278	.000150	.000555	.000982	.000554	*	*	*
33	.000156	.000275	.000153	.000160	.000281	.000155	.000569	.001006	.000574	*	*	*
34	.000159	.000277	.000160	.000163	.000283	.000162	.000585	.001029	.000598	*	*	*
35	.000163	.000280	.000168	.000166	.000286	.000171	.000603	.001054	.000625	*	*	*
36	.000167	.000285	.000177	.000170	.000290	.000180	.000624	.001084	.000656	*	*	*
37	.000171	.000290	.000185	.000175	.000295	.000189	.000648	.001122	.000688	*	*	*
38	.000176	.000297	.000191	.000179	.000302	.000195	.000676	.001170	.000723	*	*	*
39	.000181	.000305	.000198	.000184	.000309	.000201	.000708	.001227	.000760	*	*	*
40	.000186	.000313	.000204	.000189	.000318	.000207	.000744	.001293	.000800	*	*	*
41	.000193	.000323	.000212	.000196	.000327	.000215	.000784	.001365	.000844	*	*	*
42	.000201	.000335	.000223	.000204	.000339	.000226	.000828	.001442	.000896	*	*	*
43	.000212	.000352	.000238	.000215	.000355	.000242	.000878	.001523	.000958	*	*	*
44	.000226	.000372	.000258	.000229	.000376	.000263	.000934	.001608	.001031	*	*	*
45	.000243	.000397	.000283	.000247	.000403	.000289	.001000	.001703	.001119	*	*	*
46	.000262	.000426	.000309	.000267	.000432	.000316	.001072	.001809	.001216	*	*	*
47	.000281	.000455	.000334	.000286	.000462	.000342	.001144	.001923	.001308	*	*	*
48	.000297	.000479	.000355	.000303	.000487	.000365	.001207	.002040	.001378	*	*	*
49	.000310	.000500	.000372	.000316	.000507	.000383	.001260	.002157	.001425	*	*	*
50	.000323	.000522	.000390	.000330	.000528	.000401	.001311	.002281	.001465	*	*	*
51	.000339	.000548	.000409	.000346	.000554	.000423	.001365	.002410	.001510	*	*	*
52	.000357	.000578	.000429	.000365	.000587	.000445	.001417	.002525	.001562	*	*	*
53	.000377	.000615	.000450	.000387	.000627	.000466	.001470	.002619	.001629	*	*	*
54	.000398	.000654	.000470	.000411	.000671	.000487	.001525	.002697	.001710	*	*	*
55	.000420	.000694	.000491	.000435	.000715	.000508	.001577	.002761	.001794	*	*	*
56	.000442	.000733	.000513	.000458	.000757	.000531	.001633	.002833	.001879	*	*	*
57	.000465	.000771	.000539	.000482	.000798	.000558	.001703	.002935	.001979	*	*	*
58	.000488	.000809	.000569	.000507	.000838	.000590	.001797	.003089	.002098	*	*	*
59	.000513	.000848	.000602	.000532	.000877	.000624	.001911	.003288	.002235	*	*	*

**Table 10. Standard errors of the probability of dying: New Mexico, 1989–91—Con.**

Exact age in years							All other					
	Total			White			Total			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
60	.000538	.000887	.000635	.000557	.000917	.000658	.002036	.003511	.002381	*	*	*
61	.000562	.000927	.000667	.000582	.000957	.000691	.002167	.003737	.002536	*	*	*
62	.000588	.000968	.000701	.000608	.000998	.000725	.002310	.003972	.002714	*	*	*
63	.000615	.001010	.000737	.000635	.001040	.000761	.002466	.004207	.002921	*	*	*
64	.000643	.001053	.000775	.000663	.001084	.000798	.002636	.004446	.003158	*	*	*
65	.000671	.001096	.000813	.000691	.001127	.000836	.002824	.004698	.003427	*	*	*
66	.000701	.001142	.000853	.000720	.001173	.000875	.003026	.004972	.003713	*	*	*
67	.000737	.001202	.000898	.000757	.001234	.000920	.003231	.005264	.003996	*	*	*
68	.000784	.001281	.000951	.000805	.001316	.000974	.003434	.005576	.004257	*	*	*
69	.000841	.001382	.001015	.000865	.001421	.001041	.003636	.005912	.004499	*	*	*
70	.000909	.001501	.001090	.000935	.001546	.001119	.003852	.006280	.004749	*	*	*
71	.000984	.001632	.001174	.001014	.001683	.001207	.004088	.006678	.005025	*	*	*
72	.001063	.001769	.001266	.001097	.001827	.001303	.004330	.007091	.005309	*	*	*
73	.001140	.001905	.001358	.001177	.001969	.001399	.004572	.007514	.005596	*	*	*
74	.001215	.002040	.001447	.001255	.002110	.001493	.004819	.007958	.005886	*	*	*
75	.001291	.002183	.001538	.001335	.002260	.001589	.005060	.008403	.006167	*	*	*
76	.001378	.002348	.001638	.001426	.002433	.001694	.005324	.008905	.006465	*	*	*
77	.001478	.002540	.001751	.001530	.002633	.001811	.005669	.009581	.006830	*	*	*
78	.001599	.002773	.001887	.001655	.002873	.001953	.006154	.010537	.007320	*	*	*
79	.001746	.003055	.002053	.001807	.003163	.002123	.006795	.011790	.007961	*	*	*
80	.001921	.003396	.002244	.001985	.003511	.002320	.007598	.013370	.008743	*	*	*
81	.002119	.003790	.002461	.002187	.003915	.002543	.008513	.015155	.009648	*	*	*
82	.002346	.004236	.002718	.002421	.004375	.002808	.009482	.016934	.010702	*	*	*
83	.002601	.004721	.003020	.002687	.004885	.003121	.010384	.018402	.011854	*	*	*
84	.002891	.005257	.003375	.002992	.005458	.003491	.011202	.019524	.013076	*	*	*
85	.003244	.005920	.003807	.003366	.006177	.003941	.012103	.020614	.014560	*	*	*
86	.003671	.006750	.004315	.003819	.007084	.004473	.013188	.022027	.016239	*	*	*
87	.004164	.007738	.004891	.004342	.008160	.005074	.014576	.024042	.018215	*	*	*
88	.004728	.008893	.005538	.004931	.009396	.005742	.016543	.027188	.020788	*	*	*
89	.005385	.010254	.006286	.005604	.010819	.006505	.019339	.031859	.024298	*	*	*
90	.006206	.011962	.007224	.006435	.012572	.007453	.023420	.038754	.029371	*	*	*
91	.007259	.014203	.008417	.007498	.014859	.008655	.029152	.048408	.036535	*	*	*
92	.008531	.017004	.009837	.008780	.017718	.010082	.036594	.060879	.045896	*	*	*
93	.009969	.020296	.011420	.010249	.021158	.011688	.043757	.072306	.055276	*	*	*
94	.011535	.023930	.013138	.011889	.025120	.013463	.048343	.078559	.061929	*	*	*
95	.012586	.026686	.014424	.013052	.027920	.014909	.048038	.091561	.058132	*	*	*
96	.014954	.031855	.017127	.015529	.033472	.017713	.055980	.104539	.068552	*	*	*
97	.017959	.038535	.020546	.018676	.040655	.021266	.066095	.123117	.081473	*	*	*
98	.021913	.047752	.025039	.022868	.050418	.026011	.077951	.151324	.095267	*	*	*
99	.026609	.059197	.030225	.027863	.062995	.031472	.091171	.174633	.111866	*	*	*
100	.032985	.074159	.037363	.034744	.079527	.039122	.106602	.206008	.130310	*	*	*
101	.041682	.094195	.047155	.044180	.101702	.049683	.127611	.249765	.155249	*	*	*
102	.053775	.122752	.060693	.057413	.134265	.064337	.155842	.301567	.190204	*	*	*
103	.071063	.162131	.080228	.076630	.180389	.085768	.192951	.366857	.236743	*	*	*
104	.092727	.220061	.103806	.102186	.254601	.113106	.224644	.432365	.274407	*	*	*
105	.120362	.287568	.134609	.135425	.342977	.149533	.268044	.521354	.326251	*	*	*
106	.165474	.378693	.186822	.194023	.512626	.212853	.324801	.554624	.413980	*	*	*
107	.213434	.494228	.240437	.251611	.608355	.280516	.414632	.841267	.498607	*	*	*
108	.303382	.660665	.346648	.381086	.953058	.422461	.518941	.911537	.652870	*	*	*
109	.417039	.855690	.483985	.538357	.999999	.592951	.686815	.999999	.907046	*	*	*

\* Figure does not meet standards of reliability and precision.

Table 11. Standard errors of the average remaining lifetime: New Mexico, 1989–91

Exact age in years	Total			White			All other					
							Total			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
0	.078	.112	.105	.082	.116	.109	.281	.391	.393	*	*	*
1	.075	.107	.100	.078	.111	.104	.277	.385	.387	*	*	*
2	.075	.107	.100	.078	.111	.103	.276	.385	.386	*	*	*
3	.075	.107	.099	.077	.111	.103	.276	.384	.385	*	*	*
4	.075	.106	.099	.077	.110	.103	.276	.384	.385	*	*	*
5	.074	.106	.099	.077	.110	.102	.276	.384	.385	*	*	*
6	.074	.106	.099	.077	.110	.102	.275	.383	.384	*	*	*
7	.074	.106	.099	.077	.110	.102	.275	.383	.384	*	*	*
8	.074	.106	.099	.077	.110	.102	.275	.383	.384	*	*	*
9	.074	.106	.098	.077	.109	.102	.275	.383	.384	*	*	*
10	.074	.106	.098	.077	.109	.102	.275	.383	.384	*	*	*
11	.074	.106	.098	.076	.109	.102	.275	.383	.384	*	*	*
12	.074	.106	.098	.076	.109	.101	.275	.383	.384	*	*	*
13	.074	.105	.098	.076	.109	.101	.275	.382	.383	*	*	*
14	.074	.105	.098	.076	.109	.101	.275	.382	.383	*	*	*
15	.073	.105	.098	.076	.109	.101	.274	.382	.383	*	*	*
16	.073	.105	.097	.076	.108	.101	.274	.381	.383	*	*	*
17	.073	.104	.097	.075	.108	.100	.273	.380	.382	*	*	*
18	.073	.103	.097	.075	.107	.100	.273	.379	.382	*	*	*
19	.072	.103	.097	.075	.106	.100	.272	.378	.381	*	*	*
20	.072	.102	.096	.074	.105	.099	.271	.377	.381	*	*	*
21	.071	.101	.096	.074	.105	.099	.271	.376	.380	*	*	*
22	.071	.100	.095	.073	.104	.098	.270	.374	.379	*	*	*
23	.070	.100	.095	.073	.103	.098	.269	.373	.379	*	*	*
24	.070	.099	.095	.072	.102	.098	.268	.372	.378	*	*	*
25	.070	.098	.094	.072	.101	.097	.268	.371	.378	*	*	*
26	.069	.097	.094	.071	.101	.097	.267	.370	.377	*	*	*
27	.069	.097	.094	.071	.100	.097	.267	.369	.377	*	*	*
28	.069	.096	.093	.071	.100	.096	.266	.369	.376	*	*	*
29	.068	.096	.093	.071	.099	.096	.266	.368	.376	*	*	*
30	.068	.095	.093	.070	.099	.096	.266	.367	.375	*	*	*
31	.068	.095	.093	.070	.098	.096	.265	.367	.375	*	*	*
32	.068	.095	.093	.070	.098	.095	.265	.366	.375	*	*	*
33	.067	.094	.092	.070	.097	.095	.264	.366	.374	*	*	*
34	.067	.094	.092	.069	.097	.095	.264	.365	.374	*	*	*
35	.067	.093	.092	.069	.096	.095	.264	.364	.374	*	*	*
36	.067	.093	.092	.069	.096	.095	.263	.364	.373	*	*	*
37	.066	.092	.091	.069	.095	.094	.263	.363	.373	*	*	*
38	.066	.092	.091	.068	.095	.094	.263	.363	.372	*	*	*
39	.066	.092	.091	.068	.094	.094	.262	.362	.372	*	*	*
40	.066	.091	.091	.068	.094	.094	.262	.361	.371	*	*	*
41	.066	.091	.090	.068	.094	.093	.261	.361	.371	*	*	*
42	.065	.090	.090	.067	.093	.093	.261	.360	.370	*	*	*
43	.065	.090	.090	.067	.093	.093	.260	.359	.370	*	*	*
44	.065	.090	.090	.067	.092	.092	.260	.358	.369	*	*	*
45	.064	.089	.089	.066	.092	.092	.259	.357	.368	*	*	*
46	.064	.089	.089	.066	.091	.092	.258	.356	.367	*	*	*
47	.064	.088	.088	.066	.091	.091	.257	.354	.366	*	*	*
48	.063	.087	.088	.065	.090	.091	.256	.352	.365	*	*	*
49	.063	.087	.087	.065	.089	.090	.255	.350	.364	*	*	*
50	.062	.086	.087	.064	.089	.089	.254	.348	.362	*	*	*
51	.062	.085	.086	.064	.088	.089	.252	.346	.361	*	*	*
52	.062	.085	.086	.063	.087	.088	.251	.344	.360	*	*	*
53	.061	.084	.085	.063	.087	.087	.250	.342	.358	*	*	*
54	.061	.083	.084	.062	.086	.087	.249	.339	.357	*	*	*
55	.060	.082	.084	.062	.085	.086	.248	.337	.356	*	*	*
56	.059	.081	.083	.061	.084	.085	.246	.335	.355	*	*	*
57	.059	.081	.082	.061	.083	.085	.245	.332	.354	*	*	*
58	.058	.080	.081	.060	.082	.084	.244	.330	.352	*	*	*
59	.058	.079	.081	.059	.081	.083	.243	.328	.351	*	*	*



Table 11. Standard errors of the average remaining lifetime: New Mexico, 1989–91—Con.

Exact age in years	Total			White			All other					
	Both sexes	Male	Female	Both sexes	Male	Female	Total			Black		
							Both sexes	Male	Female	Both sexes	Male	Female
60	.057	.078	.080	.059	.080	.082	.242	.326	.350	*	*	*
61	.057	.077	.079	.058	.079	.081	.241	.324	.349	*	*	*
62	.056	.076	.079	.058	.078	.081	.240	.322	.348	*	*	*
63	.056	.076	.078	.057	.078	.080	.239	.319	.346	*	*	*
64	.055	.075	.077	.057	.077	.079	.238	.317	.345	*	*	*
65	.055	.074	.076	.056	.076	.078	.236	.314	.344	*	*	*
66	.054	.074	.076	.056	.076	.078	.235	.312	.342	*	*	*
67	.054	.073	.075	.055	.075	.077	.234	.310	.341	*	*	*
68	.053	.073	.075	.055	.075	.076	.233	.307	.339	*	*	*
69	.053	.072	.074	.055	.074	.076	.231	.305	.337	*	*	*
70	.053	.072	.073	.054	.074	.075	.230	.303	.335	*	*	*
71	.052	.072	.073	.054	.074	.075	.229	.301	.334	*	*	*
72	.052	.071	.072	.054	.073	.074	.228	.299	.332	*	*	*
73	.052	.071	.071	.053	.073	.073	.227	.298	.331	*	*	*
74	.052	.071	.071	.053	.073	.072	.226	.297	.329	*	*	*
75	.051	.071	.070	.053	.073	.072	.226	.296	.328	*	*	*
76	.051	.071	.070	.052	.073	.071	.226	.295	.328	*	*	*
77	.051	.071	.069	.052	.073	.071	.226	.295	.328	*	*	*
78	.051	.072	.069	.052	.074	.071	.227	.296	.329	*	*	*
79	.051	.072	.069	.053	.074	.071	.228	.298	.330	*	*	*
80	.052	.073	.069	.053	.075	.070	.230	.301	.332	*	*	*
81	.052	.074	.069	.053	.076	.071	.232	.304	.334	*	*	*
82	.052	.076	.069	.054	.078	.071	.234	.308	.336	*	*	*
83	.053	.077	.070	.054	.080	.071	.237	.313	.339	*	*	*
84	.054	.079	.070	.055	.082	.072	.241	.319	.343	*	*	*
85	.055	.082	.071	.056	.085	.072	.246	.326	.349	*	*	*
86	.056	.085	.072	.058	.088	.073	.253	.336	.357	*	*	*
87	.058	.089	.074	.059	.092	.075	.262	.349	.368	*	*	*
88	.060	.093	.075	.061	.096	.077	.274	.367	.382	*	*	*
89	.062	.099	.078	.064	.102	.079	.289	.390	.401	*	*	*
90	.065	.105	.081	.067	.109	.082	.307	.419	.422	*	*	*
91	.069	.113	.084	.070	.117	.086	.328	.454	.447	*	*	*
92	.073	.123	.089	.074	.126	.090	.350	.495	.472	*	*	*
93	.077	.133	.093	.078	.137	.095	.368	.534	.492	*	*	*
94	.082	.144	.099	.084	.149	.100	.381	.571	.503	*	*	*
95	.088	.158	.105	.090	.163	.107	.394	.620	.511	*	*	*
96	.097	.178	.116	.099	.184	.118	.425	.675	.549	*	*	*
97	.108	.203	.128	.111	.211	.132	.460	.745	.592	*	*	*
98	.122	.234	.144	.126	.245	.149	.500	.827	.639	*	*	*
99	.139	.273	.164	.145	.289	.169	.544	.905	.694	*	*	*
100	.161	.321	.188	.169	.344	.196	.596	1.005	.758	*	*	*
101	.189	.384	.220	.200	.417	.231	.660	1.125	.838	*	*	*
102	.224	.465	.260	.240	.516	.276	.736	1.257	.934	*	*	*
103	.269	.568	.311	.293	.648	.334	.819	1.403	1.039	*	*	*
104	.322	.698	.371	.358	.827	.406	.897	1.549	1.137	*	*	*
105	.389	.844	.447	.443	1.046	.500	1.001	1.721	1.273	*	*	*
106	.478	1.023	.550	.560	1.350	.630	1.132	1.892	1.457	*	*	*
107	.575	1.231	.663	.690	1.622	.778	1.301	2.306	1.647	*	*	*
108	.707	1.468	.821	.887	2.176	.993	1.464	2.356	1.906	*	*	*
109	.796	1.609	.931	1.031	2.641	1.145	1.593	2.433	2.116	*	*	*

\* Figure does not meet standards of reliability and precision.

For a list of reports published by the National Center for Health Statistics contact:

Data Dissemination Branch  
National Center for Health Statistics  
Centers for Disease Control and Prevention  
6525 Belcrest Road, Room 1064  
Hyattsville, MD 20782-2003  
(301) 436-8500  
Internet: [www.cdc.gov/nchswww/](http://www.cdc.gov/nchswww/)

# U.S. Decennial Life Tables, 1989–91

These 55 reports are published once each 10-year period by the National Center for Health Statistics.

## VOLUME I

- Number 1** *United States Life Tables.* This first report contains life tables by single years of age from birth to age 110 for the United States. Tables are included for the total population, the white population, the population other than white, and the black population. Within these large populations are tables showing the race-sex categories of male, female, and both sexes combined. Standard error tables for the probability of dying and of the average remaining lifetime are included.
- Number 2** *Methodology of the National and State Life Tables.* This report describes in detail the methods of construction of the national and State life tables.
- Number 3** *Some Trends and Comparisons of United States Life Table Data: 1900–1991.* This report deals with trends and interpretations related to life expectancy and survivorship.
- Number 4** *United States Life Tables Eliminating Certain Causes of Death.* This report provides life tables analyzed by major groups of causes of death.

## VOLUME II

### Numbers

- 1 through 51** *Alaska through Wyoming, State Life Tables.* Each of these 51 reports contains life tables for a particular State and a table that ranks each State in the order of life expectancy. All States have tables for the total population and the white population by sex. In addition, 40 States have tables for the other than white population and 33 have tables for the black population. Standard error tables for the probability of dying and of the average remaining lifetime are included.

**DEPARTMENT OF  
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention  
National Center for Health Statistics  
6525 Belcrest Road  
Hyattsville, Maryland 20782-2003

---

OFFICIAL BUSINESS  
PENALTY FOR PRIVATE USE, \$300

STANDARD MAIL (A)  
POSTAGE & FEES PAID  
PHS/NCHS  
PERMIT NO. G-281