Table 11. Mothers who smoked cigarettes during pregnancy, according to mother's detailed race, Hispanic origin, age, and education: Selected States, 1989–2001

[Data are based on birth certificates]

Characteristic of mother	1989	1990	1995	1996	1997	1998	1999	2000	2001
Race of mother <sup>1</sup>	Percent of mothers who smoked <sup>2</sup>								
All races	19.5	18.4	13.9	13.6	13.2	12.9	12.6	12.2	12.0
White Black or African American American Indian or Alaska Native Asian or Pacific Islander <sup>3</sup> Chinese Japanese Fillpino Hawaiian Other Asian or Pacific Islander	20.4 17.1 23.0 5.7 2.7 8.2 5.1 19.3 4.2	19.4 15.9 22.4 5.5 2.0 8.0 5.3 21.0 3.8	15.0 10.6 20.9 3.4 0.8 5.2 3.4 15.9 2.7	14.7 10.2 21.3 3.3 0.7 4.8 3.5 15.3 2.7	14.3 9.7 20.8 3.2 1.0 4.7 3.4 15.8 2.5	14.0 9.5 20.2 3.1 0.8 4.8 3.3 16.8 2.4	13.6 9.3 20.2 2.9 0.5 4.5 3.3 14.7 2.3	13.2 9.1 20.0 2.8 0.6 4.2 3.2 14.4 2.3	13.0 9.0 19.9 2.8 0.7 3.8 3.2 14.8 2.3
Hispanic origin and race of mother <sup>4</sup>									
Hispanic or Latino Mexican Puerto Rican Cuban Central and South American Other and unknown Hispanic or Latino Not Hispanic or Latino:	8.0 6.3 14.5 6.9 3.6 12.1	6.7 5.3 13.6 6.4 3.0 10.8	4.3 3.1 10.4 4.1 1.8 8.2	4.3 3.1 11.0 4.7 1.8 9.1	4.1 2.9 11.0 4.2 1.8 8.5	4.0 2.8 10.7 3.7 1.5 8.0	3.7 2.6 10.5 3.3 1.4 7.7	3.5 2.4 10.3 3.3 1.5 7.4	3.2 2.4 9.7 3.0 1.3 6.8
WhiteBlack or African American	21.7 17.2	21.0 15.9	17.1 10.6	16.9 10.3	16.5 9.8	16.2 9.6	15.9 9.4	15.6 9.2	15.5 9.1
Age of mother <sup>1</sup>									
Under 15 years 15–19 years 15–17 years 18–19 years 20–24 years 25–29 years 30–34 years 35–39 years 40–54 years <sup>5</sup>	7.7 22.2 19.0 23.9 23.5 19.0 15.7 13.6 13.2	7.5 20.8 17.6 22.5 22.1 18.0 15.3 13.3 12.3	7.3 16.8 14.6 18.1 17.1 12.8 11.4 12.0 10.1	7.7 17.2 15.4 18.3 16.8 12.3 10.9 11.7 10.1	8.1 17.6 15.5 18.8 16.6 11.8 10.0 11.1	7.7 17.8 15.5 19.2 16.5 11.4 9.3 10.6 10.0	7.8 18.1 15.5 19.5 16.7 11.0 8.6 9.9 9.5	7.1 17.8 15.0 19.2 16.8 10.5 8.0 9.1 9.5	6.0 17.5 14.4 19.0 17.0 10.3 7.6 8.6 9.3
Education of mother <sup>6</sup>	Percent of mothers 20 years of age and over who smoked <sup>2</sup>								
0–8 years	18.9 42.2 22.8 13.7 5.0	17.5 40.5 21.9 12.8 4.5	11.0 32.0 18.3 10.6 2.7	10.3 31.1 18.0 10.4 2.6	9.9 30.2 17.5 9.9 2.4	9.5 29.3 17.1 9.6 2.2	8.9 29.0 16.9 9.4 2.1	7.9 28.2 16.6 9.1 2.0	7.2 27.6 16.5 9.2 1.9

<sup>&</sup>lt;sup>1</sup>Data from States that did not require the reporting of mother's tobacco use during pregnancy on the birth certificate are not included. Reporting area for tobacco use increased from 43 States and the District of Columbia (DC) in 1989 to 49 States and DC in 2000–01 (see Appendix II, Tobacco use).

NOTES: The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Interpretation of trend data should take into consideration expansion of reporting areas and immigration. Data for additional years are available (see Appendix III).

SOURCES: Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System, Birth File. Martin JA, Hamilton BE, Ventura SJ, Menacker F, Park MM, Sutton PD. Births: Final Data for 2001. National vital statistics reports; vol 51, no 2. Hyattsville, Maryland: National Center for Health Statistics, 2002; Births: Final data for each data year 1997–2000. National vital statistics reports. Hyattsville, Maryland; Final natality statistics for each data year 1989–96. Monthly vital statistics report. Hyattsville, Maryland.

Health, United States, 2003

<sup>&</sup>lt;sup>2</sup>Excludes live births for whom smoking status of mother is unknown.

<sup>&</sup>lt;sup>3</sup>Maternal tobacco use during pregnancy was not reported on the birth certificates of California, which in 2000 accounted for 32 percent of the births to Asian or Pacific Islander mothers.

<sup>&</sup>lt;sup>4</sup>Data from States that did not require the reporting of either Hispanic origin of mother or tobacco use during pregnancy on the birth certificate are not included. Reporting area for tobacco use and Hispanic origin of mother increased from 42 States and DC in 1989 to 49 States and DC in 2000–01. See Appendix II, Hispanic origin; Tobacco use.

<sup>&</sup>lt;sup>5</sup>Prior to 1997 data are for live births to mothers 45–49 years of age.

<sup>&</sup>lt;sup>6</sup>Data from States that did not require the reporting of either mother's education or tobacco use during pregnancy on the birth certificate are not included. Reporting area for tobacco use and education of mother increased from 42 States and DC in 1989 to 49 States and DC in 2000–01. See Appendix II, Education; Hispanic origin).