Table 45 (page 1 of 3). Death rates for homicide, according to sex, race, Hispanic origin, and age: United States, selected years 1950–2001

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950¹	1960¹	1970	1980	1990	1995	2000 ²	2001	
All persons	Deaths per 100,000 resident population								
All ages, age adjusted All ages, crude	5.1	5.0	8.8	10.4	9.4	8.3	5.9	7.1	
	5.0	4.6	8.1	10.6	9.9	8.5	6.0	7.1	
Under 1 year.	4.4	4.8	4.3	5.9	8.4	8.2	9.2	8.2	
1–14 years	0.6	0.6	1.1	1.5	1.8	1.9	1.3	1.3	
1–4 years	0.6	0.7	1.9	2.5	2.5	2.9	2.3	2.7	
5–14 years	0.5	0.5	0.9	1.2	1.5	1.5	0.9	0.8	
15–24 years	5.8	5.6	11.3	15.4	19.7	19.6	12.6	13.3	
15–19 years	3.9	3.9	7.7	10.5	16.9	17.8	9.5	9.4	
20–24 years. 25–44 years 25–34 years. 35–44 years. 45–64 years. 55–64 years.	8.5 8.9 9.3 8.4 5.0 5.9 3.9	7.7 8.5 9.2 7.8 5.3 6.1 4.1	15.6 14.9 16.2 13.5 8.7 10.0 7.1	20.2 17.5 19.3 14.9 9.0 11.0 7.0	22.2 14.7 17.4 11.6 6.3 7.5 5.0	21.5 11.9 14.4 9.4 5.4 6.0 4.4	16.0 8.7 10.4 7.1 4.0 4.7 3.0	17.3 11.2 13.1 9.5 5.4 6.3 4.0	
65 years and over	3.0	2.7	4.6	5.5	4.0	3.1	2.4	2.7	
65-74 years.	3.2	2.8	4.9	5.7	3.8	3.2	2.4	2.9	
75-84 years.	2.5	2.3	4.0	5.2	4.3	3.0	2.4	2.5	
85 years and over	2.3	2.4	4.2	5.3	4.6	3.2	2.4	2.4	
Male All ages, age adjusted ³ All ages, crude	7.9 7.7	7.5 6.8	14.3 13.1	16.6 17.1	14.8 15.9	12.8 13.4	9.0 9.3	10.8 11.1	
Under 1 year	4.5	4.7	4.5	6.3	8.8	9.0	10.4	9.5	
	0.6	0.6	1.2	1.6	2.0	2.2	1.5	1.5	
	0.5	0.7	1.9	2.7	2.7	3.1	2.5	3.0	
5-14 years	0.6	0.5	1.0	1.2	1.7	1.9	1.1	0.9	
15-24 years	8.6	8.4	18.2	24.0	32.5	32.8	20.9	22.2	
15-19 years	5.5	5.7	12.1	15.9	27.8	29.1	15.5	15.7	
20-24 years	13.5	11.8	25.6	32.2	36.9	36.5	26.7	28.9	
25-44 years	13.8	12.8	24.4	28.9	23.5	18.2	13.3	17.2	
25–34 years. 35–44 years. 45–64 years	14.4 13.2 8.1 9.5 6.3	13.9 11.7 8.1 9.4 6.4	26.8 21.7 14.8 16.8 12.1	31.9 24.5 15.2 18.4 11.8	27.7 18.6 10.2 11.9 8.0	22.5 14.0 8.3 9.2 7.0	16.7 10.3 6.0 6.9 4.6	20.8 13.9 8.1 9.5 5.9	
65 years and over	4.8	4.3	7.7	8.8	5.8	4.2	3.3	3.6	
	5.2	4.6	8.5	9.2	5.8	4.5	3.4	4.0	
	3.9	3.7	5.9	8.1	5.7	3.7	3.2	3.1	
	2.5	3.6	7.4	7.5	6.7	4.1	3.3	3.2	
Female	0.4	0.0	0.7		4.0	0.7		0.0	
All ages, age adjusted ³	2.4	2.6	3.7	4.4	4.0	3.7	2.8	3.3	
All ages, crude	2.4	2.4	3.4	4.5	4.2	3.8	2.8	3.3	
Under 1 year	4.2	4.9	4.1	5.6	8.0	7.4	7.9	6.9	
	0.6	0.5	1.0	1.4	1.6	1.5	1.1	1.1	
	0.7	0.7	1.9	2.2	2.3	2.6	2.1	2.4	
	0.5	0.4	0.7	1.1	1.2	1.0	0.7	0.7	
	3.0	2.8	4.6	6.6	6.2	5.9	3.9	3.9	
15–19 years.	2.4	1.9	3.2	4.9	5.4	5.8	3.1	2.7	
20–24 years.	3.7	3.8	6.2	8.2	7.0	6.0	4.7	5.1	
25–44 years .	4.2	4.3	5.8	6.4	6.0	5.6	4.0	5.2	
25–34 years.	4.5	4.6	6.0	6.9	7.1	6.3	4.1	5.3	
35–44 years.	3.8	4.0	5.7	5.7	4.8	4.9	4.0	5.1	
45–64 years	1.9	2.5	3.1	3.4	2.8	2.6	2.1	2.8	
45–54 years.	2.3	2.9	3.7	4.1	3.2	2.9	2.5	3.2	
55–64 years.	1.4	2.0	2.5	2.8	2.3	2.1	1.6	2.2	
65 years and over	1.4	1.3	2.3	3.3	2.8	2.4	1.8	2.0	
65–74 years.	1.3	1.3	2.2	3.0	2.2	2.1	1.6	2.0	
75–84 years	1.4	1.3	2.7	3.5	3.4	2.6	2.0	2.1	
	2.1	1.6	2.5	4.3	3.8	2.9	2.0	2.0	
White male ⁴ All ages, age adjusted ³	3.8	2.0	7.0	10.4	ρo	7.9	. 50	71	
All ages, crude	3.6	3.9 3.6	7.2 6.6	10.4 10.7	8.3 8.8	7.3 7.5	5.2 5.2	7.1 7.2	
Under 1 year. 1–14 years	4.3	3.8	2.9	4.3	6.4	7.1	8.2	7.3	
	0.4	0.5	0.7	1.2	1.3	1.5	1.2	1.1	
	3.2	5.0	7.6	15.1	15.2	15.9	9.9	11.2	
	5.4	5.5	11.6	17.2	13.0	10.4	7.4	11.5	
	4.9	5.7	12.5	18.5	14.7	12.1	8.4	12.3	
	6.1	5.2	10.8	15.2	11.1	8.7	6.5	10.7	
45–64 years	4.8	4.6	8.3	9.8	6.9	5.5	4.1	6.4	
	3.8	3.1	5.4	6.7	4.1	2.9	2.5	3.0	

See footnotes at end of table.

184 Health, United States, 2003

Table 45 (page 2 of 3). Death rates for homicide, according to sex, race, Hispanic origin, and age: United States, selected years 1950-2001

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950¹	1960¹	1970	1980	1990	1995	2000²	2001	
Black or African American male ⁴	Deaths per 100,000 resident population								
All ages, age adjusted ³ All ages, crude	47.0 44.7	42.3 35.0	78.2 66.0	69.4 65.7	63.1 68.5	51.1 54.5	35.4 37.2	36.2 38.3	
Under 1 year. 1–14 years ⁵ 15–24 years 25–44 years 25–34 years. 35–44 years. 45–64 years 65 years and over	1.8 53.8 92.8 104.3 80.0 46.0 16.5	10.3 1.5 43.2 80.5 86.4 74.4 44.6 17.3	14.3 4.4 98.3 140.2 154.5 124.0 82.3 33.3	18.6 4.1 82.6 130.0 142.9 109.3 70.6 30.9	21.4 5.8 137.1 105.4 123.7 81.2 41.4 25.7	20.3 5.8 129.4 75.8 95.1 54.9 33.4 20.3	23.3 3.1 85.3 55.8 73.9 38.5 21.9 12.8	21.0 3.6 85.7 58.1 78.8 38.5 22.7 11.5	
American Indian or Alaska Native male ⁴							'		
All ages, age adjusted ³ All ages, crude				23.3 23.1	16.7 16.6	14.4 15.7	10.7 10.7	9.3 9.6	
15–24 years				35.4 39.2 22.1	25.1 25.7 14.8	28.0 24.6 12.0	17.0 17.0 *	16.1 13.6 8.9	
Asian or Pacific Islander male ⁴							•		
All ages, age adjusted ³ All ages, crude				9.1 8.3	7.3 7.9	7.2 7.5	4.3 4.4	6.0 6.3	
15–24 years				9.3 11.3 10.4	14.9 9.6 7.0	17.2 7.4 7.7	7.8 4.6 6.1	9.1 8.5 7.7	
Hispanic or Latino male ^{4,6}									
All ages, age adjusted ³ All ages, crude					27.4 31.0	20.4 23.5	11.8 13.4	12.9 14.5	
Under 1 year					8.7 3.1 55.4	5.9 3.2 54.7	6.6 1.7 28.5	7.9 1.5 30.5	
25–44 years		 			46.4 50.9 39.3	28.8 33.0 22.8	17.2 19.9 13.5	19.2 21.6 15.9	
45–64 years					20.5 9.4	14.7 5.8	9.1 4.4	9.8 5.5	
White, not Hispanic or Latino male ⁶							!		
All ages, age adjusted ³ All ages, crude					5.6 5.8	4.8 4.9	3.6 3.6	5.6 5.6	
Under 1 year					5.4 0.9	6.8 1.1	8.3 1.0	6.8 1.0	
15–24 years					7.5 8.7	7.2 7.2	4.7 5.2	5.7 9.4	
25–34 years					9.3 8.0	7.7 6.7	5.2 5.2	9.3 9.5	
45–64 years					5.7 3.7	4.6 2.6	3.6 2.3	5.9 2.8	
White female ⁴ All ages, age adjusted ³	1.4	1.5	2.3	3.2	2.7	2.7	2.1	2.6	
All ages, crude	1.4	1.4	2.1	3.2	2.8	2.7	2.1	2.6	
Under 1 year	3.9 0.4	3.5 0.4	2.9 0.7	4.3 1.1	5.1 1.0	5.1 1.1	5.0 0.8	5.1 0.9	
15–24 years	1.3 2.0	1.5 2.1	2.7 3.3	4.7 4.2	4.0 3.8	4.0 3.7	2.7 2.9	3.0 4.0	
45–64 years	1.5 1.2	1.7 1.2	2.1 1.9	2.6 2.9	2.3 2.2	2.2 2.0	1.8 1.6	2.4 1.8	
Black or African American female ⁴									
All ages, age adjusted ³ All ages, crude	11.1 11.5	11.4 10.4	14.7 13.2	13.2 13.5	12.5 13.4	10.4 10.9	7.1 7.2	7.4 7.4	
Under 1 year	1.8	13.8 1.2	10.7 3.1	12.8 3.3	22.8 4.7	20.1 3.4	22.2 2.7	16.7 2.1	
15–24 years	16.5 22.5	11.9 22.7	17.7 25.3	18.4 22.6	18.9 21.0	16.4 17.1	10.7 11.0	8.9 12.5	
45–64 years	6.8 3.6	10.3 3.0	13.4 7.4	10.8 8.0	6.5 9.4	5.9 6.8	4.5 3.5	5.7 3.7	

See footnotes at end of table.

Health, United States, 2003

185

Table 45 (page 3 of 3). Death rates for homicide, according to sex, race, Hispanic origin, and age: United States, selected years 1950-2001

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950¹	1960¹	1970	1980	1990	1995	2000 ²	2001
American Indian or Alaska Native female ⁴	Deaths per 100,000 resident population							
All ages, age adjusted ³ All ages, crude				8.1 7.7	4.6 4.8	5.3 5.1	3.0 2.9	4.2 4.2
15–24 years				* 13.7	* 6.9	* 8.3	* 5.9	* 5.9
45–64 years				*	*	*	*	*
Asian or Pacific Islander female ⁴								
All ages, age adjusted ³ All ages, crude				3.1 3.1	2.8 2.8	2.4 2.6	1.7 1.7	2.5 2.7
15–24 years				* 4.6	* 3.8	3.4 3.6	* 2.2	4.0
45–64 years				*	*	2.2	2.0	2.7
Hispanic or Latino female ^{4,6}								
All ages, age adjusted ³ All ages, crude					4.3 4.7	4.0 4.2	2.8 2.8	3.1 3.2
Jnder 1 year					* 1.9	* 1.8	7.4 1.0	5.5 1.0
15–24 years					8.1 6.1	6.4 5.6	3.7 3.7	4.0 4.8
45-64 years					3.3	3.4 2.4	2.9 2.4	3.0 2.0
White, not Hispanic or Latino female ⁶								
All ages, age adjusted ³ All ages, crude					2.5 2.5	2.3 2.3	1.9 1.9	2.5 2.4
Jnder 1 year					4.4	4.4	4.1	4.7
I–14 years					0.8 3.3	0.9 3.4	0.8 2.3	0.9 2.7
25–44 years					3.5 2.2	3.3 1.9	2.7 1.6	3.7 2.3
65 years and over					2.2	1.9	1.6	1.8

^{- - -} Data not available

NOTES: Population estimates used to compute rates for 1991-2000 differ from those used previously. Starting with Health, United States, 2003, rates for 1991-99 were revised using intercensal population estimates based on Census 2000. Rates for 2000 were revised based on Census 2000 counts. Rates for 2001 were computed using 2000-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Figures for 2001 include September 11 related deaths for which death certificates were filed as of October 24, 2002. Underlying cause of death code numbers are based on the applicable revision of the International Classification of Diseases (ICD) for data years shown. For the period 1980-98, causes were coded using ICD-9 codes that are most nearly comparable with the 113 cause list for ICD-10. See Appendix II, tables IV and updated V for new terrorism-related ICD-10 codes. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see www.cdc.gov/ncipc/wisqars, a Web-based interactive database for injury data. Data for additional years are available. See Appendix III.

SOURCES: Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington: U.S. Government Printing Office. 1968; numerator data from National Vital Statistics System, annual mortality files; denominator data from national population estimates for race groups from table 1 and unpublished Hispanic population estimates for 1985-96 prepared by the Housing and Household Economic Statistics Division, U.S. Bureau of the Census; additional mortality tables are available at

www.cdc.gov/nchs/datawh/statab/unpubd/mortabs.htm; Arias E, Anderson RN, Kung HC, Murphy SL, Kochanek KD. Deaths: Final data for 2001. National vital statistics reports. Vol 52 no 3. Hyattsville, Maryland: National Center for Health Statistics. 2003.

186 Health, United States, 2003

^{*} Rates based on fewer than 20 deaths are considered unreliable and are not shown.

¹Includes deaths of persons who were not residents of the 50 States and the District of Columbia.

²Starting with 1999 data, cause of death is coded according to ICD-10. To estimate change between 1998 and 1999, compare the 1999 rate with the comparability-modified rate for 1998. See Appendix II, Comparability ratio and tables V and VI.

³Age-adjusted rates are calculated using the year 2000 standard population starting with *Health, United States, 2001*. See Appendix II, Age adjustment. ⁴The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for the American Indian or Alaska Native and Asian or Pacific Islander populations are known to be underestimated. See Appendix II, Race, for a discussion of sources of bias in death rates by race and Hispanic origin.

⁵In 1950 rate is for the age group under 15 years.

⁶Prior to 1997, excludes data from States lacking an Hispanic-origin item on the death certificate. See Appendix II, Hispanic origin.