Table 63 (page 1 of 3). Use of selected substances by high school seniors, eighth-, and tenth-graders, according to sex and race: United States, selected years 1980–2002

[Data are based on a survey of high school seniors and eighth-graders in the coterminous United States]

Substance, sex, race, and grade in school	1980	1990	1991	1995	1998	1999	2000	2001	2002
Cigarettes			Pe	rcent using s	substance in	the past mo	nth		
All seniors	30.5	29.4	28.3	33.5	35.1	34.6	31.4	29.5	26.7
Male	26.8 33.4	29.1 29.2	29.0 27.5	34.5 32.0	36.3 33.3	35.4 33.5	32.8 29.7	29.7 28.7	27.4 25.5
White	31.0 25.2	32.5 12.0	31.8 9.4	37.3 15.0	41.0 14.9	39.1 14.9	36.6 13.6	34.1 12.9	30.9 11.3
All tenth-graders			20.8	27.9	27.6	25.7	23.9	21.3	17.7
Male Female			20.8 20.7	27.7 27.9	26.2 29.1	25.2 25.8	23.8 23.6	20.9 21.5	16.7 18.6
White			23.9 6.4	31.2 12.2	32.4 13.8	29.1 11.0	27.3 11.3	24.0 10.9	20.8 9.1
All eighth-graders			14.3	19.1	19.1	17.5	14.6	12.2	10.7
Male			15.5 13.1	18.8 19.0	18.0 19.8	16.7 17.7	14.3 14.7	12.2 12.0	11.0 10.4
White			15.0 5.3	21.7 8.2	21.1 10.8	19.0 10.7	16.4 8.4	12.8 8.0	11.1 7.3
Marijuana									
All seniors	33.7	14.0	13.8	21.2	22.8	23.1	21.6	22.4	21.5
Male	37.8 29.1	16.1 11.5	16.1 11.2	24.6 17.2	26.5 18.8	26.3 19.7	24.7 18.3	25.6 19.1	25.3 17.4
White	34.2 26.5	15.6 5.2	15.0 6.5	21.5 17.8	24.2 18.3	23.4 20.4	22.0 17.5	23.9 16.5	22.8 16.4
All tenth-graders			8.7	17.2	18.7	19.4	19.7	19.8	17.8
Male			10.1 7.3	19.2 15.0	20.3 17.2	21.8 17.0	23.3 16.2	22.7 16.8	19.3 16.4
White			9.4 3.8	17.7 15.1	19.5 14.5	20.2 14.7	20.1 17.0	20.4 16.5	19.1 14.4
All eighth-graders			3.2	9.1	9.7	9.7	9.1	9.2	8.3
Male			3.8 2.6	9.8 8.2	10.3 8.8	10.5 8.8	10.2 7.8	11.0 7.3	9.5 7.1
White			3.0 2.1	9.0 7.0	8.9 9.4	8.5 10.0	8.3 8.5	8.6 7.7	7.9 7.1
Cocaine									
All seniors	5.2	1.9	1.4	1.8	2.4	2.6	2.1	2.1	2.3
Male	6.0 4.3	2.3 1.3	1.7 0.9	2.2 1.3	3.0 1.7	3.3 1.8	2.7 1.6	2.5 1.6	2.7 1.8
White	5.4 2.0	1.8 0.5	1.3 0.8	1.7 0.4	2.7 0.4	2.8 0.5	2.2 1.0	2.3 0.6	2.8 0.2
All tenth-graders			0.7	1.7	2.1	1.9	1.8	1.3	1.6
Male			0.7 0.6	1.8 1.5	2.4 1.8	2.2 1.6	2.1 1.4	1.5 1.2	1.8 1.4
White			0.6 0.2	1.7 0.4	2.0 0.8	1.9 0.3	1.7 0.4	1.2 0.3	1.7 0.4
All eighth-graders			0.5	1.2	1.4	1.3	1.2	1.2	1.1
Male			0.7 0.4	1.1 1.2	1.5 1.2	1.4 1.2	1.3 1.1	1.1 1.2	1.1 1.1
White			0.4 0.4	1.0 0.4	1.0 0.6	1.1 0.3	1.1 0.5	1.1 0.4	1.0 0.5

See footnotes at end of table.

Health, United States, 2003 219

Table 63 (page 2 of 3). Use of selected substances by high school seniors, eighth-, and tenth-graders, according to sex and race: United States, selected years 1980–2002

[Data are based on a survey of high school seniors and eighth-graders in the coterminous United States]

Substance, sex, race, and grade in school	1980	1990	1991	1995	1998	1999	2000	2001	2002	
Inhalants	Percent using substance in the past month									
All seniors	1.4	2.7	2.4	3.2	2.3	2.0	2.2	1.7	1.5	
Male	1.8 1.0	3.5 2.0	3.3 1.6	3.9 2.5	2.9 1.7	2.5 1.5	2.9 1.7	2.3 1.1	2.2 0.8	
White	1.4 1.0	3.0 1.5	2.4 1.5	3.7 1.1	2.6 1.0	2.1 0.4	2.1 2.1	1.8 1.3	1.3 1.2	
All tenth-graders			2.7	3.5	2.9	2.6	2.6	2.5	2.4	
Male			2.9 2.6	3.8 3.2	3.2 2.6	2.9 2.2	3.0 2.2	2.5 2.4	2.3 2.4	
White			2.9 2.0	3.9 1.2	3.3 1.1	2.9 0.8	2.8 1.5	2.5 0.9	2.6 1.5	
All eighth-graders			4.4	6.1	4.8	5.0	4.5	4.0	3.8	
Male			4.1 4.7	5.6 6.6	4.8 4.7	4.6 5.3	4.1 4.8	3.6 4.3	3.5 3.9	
White			4.5 2.3	7.0 2.3	5.3 2.2	5.6 2.3	4.5 2.3	4.1 2.6	3.9 2.7	
MDMA (Ecstasy)										
All seniors					1.5	2.5	3.6	2.8	2.4	
Male					2.3 0.8	2.6 2.5	4.1 3.1	3.7 2.0	2.6 2.1	
White					1.8 0.2	2.7 0.0	3.9 1.9	2.8 0.9	2.5 0.5	
All tenth-graders					1.3	1.8	2.6	2.6	1.8	
Male					1.4 1.1	1.7 1.9	2.5 2.5	3.5 1.6	1.6 1.8	
White					1.3 0.7	2.1 0.3	2.5 1.8	2.6 1.0	2.3 0.5	
All eighth-graders					0.9	0.8	1.4	1.8	1.4	
Male					1.0 0.7	0.9 0.7	1.6 1.2	1.9 1.8	1.5 1.3	
White					0.9 0.4	0.9 0.4	1.4 0.8	2.0 1.1	1.0 0.6	
Alcohol ¹										
All seniors	72.0	57.1	54.0	51.3	52.0	51.0	50.0	49.8	48.6	
Male Female	77.4 66.8	61.3 52.3	58.4 49.0	55.7 47.0	57.6 46.9	55.3 46.8	54.0 46.1	54.7 45.1	52.3 45.1	
White	75.8 47.7	62.2 32.9	57.7 34.4	54.8 37.4	57.6 33.6	54.9 30.8	55.3 29.3	55.3 29.6	52.7 30.7	
All tenth-graders			42.8	38.8	38.8	40.0	41.0	39.0	35.4	
Male			45.5 40.3	39.7 37.8	40.0 37.7	42.3 38.1	43.3 38.6	41.1 36.8	35.3 35.7	
White			45.7 30.2	41.3 24.9	42.5 24.2	43.4 24.6	44.3 24.7	41.0 26.0	39.0 23.2	
All eighth-graders			25.1	24.6	23.0	24.0	22.4	21.5	19.6	
Male Female			26.3 23.8	25.0 24.0	24.0 21.9	24.8 23.3	22.5 22.0	22.3 20.6	19.1 20.0	
White			26.0 17.8	25.4 17.3	24.0 15.4	25.6 16.8	23.9 15.1	22.5 14.9	20.4 14.7	

See footnotes at end of table.

220 Health, United States, 2003

Table 63 (page 3 of 3). Use of selected substances by high school seniors, eighth-, and tenth-graders, according to sex and race: United States, selected years 1980–2002

[Data are based on a survey of high school seniors and eighth-graders in the coterminous United States]

Substance, sex, race, and grade in school	1980	1990	1991	1995	1998	1999	2000	2001	2002	
Binge drinking ²	Percent in last 2 weeks									
All seniors	41.2	32.2	29.8	29.8	31.5	30.8	30.0	29.7	28.6	
Male	52.1 30.5	39.1 24.4	37.8 21.2	36.9 23.0	39.2 24.0	38.1 23.6	36.7 23.5	36.0 23.7	34.2 23.0	
White	44.6 17.0	36.2 11.6	32.9 11.8	32.9 15.5	36.6 12.7	34.8 11.9	34.4 11.0	34.5 12.6	32.9 10.4	
All tenth-graders			22.9	24.0	24.4	25.6	26.2	24.9	22.4	
Male			26.4 19.5	26.4 21.5	26.7 22.2	29.7 21.8	29.8 22.5	28.6 21.4	23.8 21.0	
WhiteBlack or African American			24.4 14.4	25.7 12.3	26.8 12.5	27.7 12.9	28.5 12.9	26.4 12.3	24.6 12.4	
All eighth-graders			12.9	14.5	13.7	15.2	14.1	13.2	12.4	
Male			14.3 11.4	15.1 13.9	14.4 12.7	16.4 13.9	14.4 13.6	13.7 12.4	12.5 12.1	
White			12.6 9.9	14.5 10.0	13.5 9.1	15.2 10.8	14.6 9.3	13.1 8.8	12.3 9.9	

^{- - -} Data not available.

NOTES: Because of methodological differences among the National Household Survey on Drug Abuse (NHSDA), Monitoring the Future Study (MTF), and Youth Risk Behavior Survey (YRBS), rates of substance use measured by these surveys are not directly comparable. See Appendix I, NHSDA, MTF, and YRBS. Data for additional years are available (see Appendix III).

SOURCE: National Institutes of Health, National Institute on Drug Abuse (NIDA), Monitoring the Future Study, Annual surveys.

Health, United States, 2003 221

^{0.0} Quantity more than zero but less than 0.05.

¹ In 1993 the alcohol question was changed to indicate that a "drink" meant "more than a few sips." 1993 data, available electronically, are based on a half sample.

²Five or more alcoholic drinks in a row at least once in the prior 2-week period.