Table 128 (page 1 of 2). Medicaid coverage among persons under 65 years of age, according to selected characteristics: United States, selected years 1984–2001

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1984	1989	1995	1996	1997 ¹	1998	1999	2000	2001
	Number in millions								
Total ²	14.0	15.4	26.6	25.8	22.9	21.1	21.9	22.9	25.2
		Percent of population							
Total, age adjusted ^{2,3}	6.7 6.8	7.1 7.2	11.3 11.5	10.9 11.1	9.6 9.7	8.8 8.9	9.0 9.1	9.4 9.5	10.3 10.4
Age									
Jnder 18 years Under 6 years 6–17 years 18–44 years 18–24 years 25–34 years 35–44 years 45–64 years 55–64 years 55–64 years	11.9 15.5 10.1 5.1 6.4 5.3 3.5 3.4 3.2 3.6	12.6 15.7 10.9 5.2 6.8 5.2 4.0 4.3 3.8 4.9	21.5 29.3 17.4 7.8 10.4 8.2 5.9 5.6 5.1 6.4	20.7 28.2 16.9 7.6 9.7 7.8 6.2 5.3 4.9 5.9	$18.4 \\ 24.7 \\ 15.2 \\ 6.6 \\ 8.8 \\ 6.8 \\ 5.2 \\ 4.6 \\ 4.0 \\ 5.6 \\$	17.1 22.4 14.5 5.8 8.0 5.7 4.6 4.5 4.1 5.0	18.1 23.5 15.5 5.7 8.1 5.7 4.3 4.4 3.9 5.3	$19.4 \\ 24.3 \\ 17.0 \\ 5.6 \\ 8.1 \\ 5.5 \\ 4.3 \\ 4.5 \\ 4.2 \\ 4.9 \\$	21.2 25.8 19.0 6.3 8.4 6.2 5.1 4.7 4.4 5.2
Sex ³									
lale emale	5.2 8.0	5.6 8.6	9.2 13.3	8.9 12.8	8.1 11.0	7.5 10.1	7.7 10.4	8.0 10.8	8.9 11.6
Race ^{3,4}									
White only	4.6 18.9 # 9.1	5.1 17.8 # 11.3	8.8 26.0 # 10.7	8.7 23.0 # *11.5	7.5 20.5 # 9.4	6.7 19.6 # 6.7	6.9 18.7 41.3 8.4	7.2 19.4 44.2 7.8	8.1 20.4 15.5 8.8
2 or more races							15.8	15.6	14.6
Hispanic origin and race ^{3,4}									
Hispanic or Latino Mexican Puerto Rican Cuban Other Hispanic or Latino Not Hispanic or Latino White only Black or African American only	12.2 11.1 28.6 4.8 7.4 6.2 3.7 19.1	12.7 11.5 26.9 7.8 10.4 6.6 4.2 17.8	19.8 18.8 31.1 13.8 16.9 10.2 7.1 25.6	18.5 17.6 31.3 *13.1 15.0 9.7 7.0 22.7	16.0 15.3 28.9 8.2 13.9 8.7 6.2 20.3	14.1 12.6 24.5 *9.1 13.9 8.0 5.7 19.4	14.1 12.4 27.0 8.3 13.8 8.2 6.0 18.7	14.2 12.5 27.6 9.7 14.1 8.6 6.3 19.3	16.0 14.6 28.5 12.2 15.0 9.3 7.0 20.3
Age and percent of poverty level ⁵									
All ages: ³ Below 100 percent 100–149 percent 150–199 percent 200 percent or more	30.5 7.5 3.1 0.6	35.3 11.0 5.0 1.1	44.7 18.0 7.9 1.8	42.9 17.4 8.0 1.7	38.8 17.5 7.4 1.7	37.9 16.0 7.2 1.8	36.8 18.6 9.8 2.0	37.2 20.3 10.8 2.3	39.0 23.5 13.3 2.6
Under 18 years: Below 100 percent 100–149 percent 150–199 percent 200 percent or more	43.1 9.0 4.4 0.8	47.8 12.3 6.1 1.6	66.0 27.2 13.1 3.3	65.2 26.6 12.2 2.8	59.7 30.2 12.2 2.9	58.7 25.9 12.8 3.2	59.9 33.5 18.0 3.7	60.9 37.1 21.5 4.7	64.3 41.4 26.5 5.3

See footnotes at end of table.

Table 128 (page 2 of 2). Medicaid coverage among persons under 65 years of age, according to selected characteristics: United States, selected years 1984–2001

1997¹ Characteristic 1984 1989 1995 1996 1998 1999 2000 2001 Geographic region³ Percent of population 11.2 10.8 6.8 11.7 9.8 10.1 10.5 8.5 11.5 7.2 7.5 10.3 8.7 8.2 7.5 7.3 7.9 9.0 5.0 6.4 11.1 11.1 8.6 8.6 8.9 9.4 10.7 6.9 8.2 12.4 12.4 11.4 9.7 10.3 10.2 10.6 Location of residence³ Within MSA⁶ 10.4 7.1 7.0 9.5 9.9 8.8 9.8 11.1 8.5 8.4 Outside MSA⁶ 7.8 12.0 9.8 11.5 11.9 5.9 12.7 12.4

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Estimates calculated upon request.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error of 20-30 percent. Data not shown have a relative standard error of greater than 30 percent.

- - Data not available.

¹In 1997 the National Health Interview Survey (NHIS) was redesigned, including changes to the questions on health insurance coverage. See Appendix I, National Health Interview Survey and Appendix II, Health insurance coverage.

²Includes all other races not shown separately and unknown poverty level. ³Estimates are for persons under 65 years of age and are age adjusted to the year 2000 standard using three age groups: under 18 years, 18–44 years, and 45–64 years. See Annendix II. Age adjustment

years. See Appendix II, Age adjustment. ⁴The race groups, white, black, American Indian and Alaska Native (Al/AN), Asian, Native Hawaiian and Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with data year 1999 race-specific estimates are tabulated according to 1997 Standards for Federal data on Race and Ethnicity and are not strictly comparable with estimates for earlier years. The five single race categories plus multiple race categories shown in the table conform to 1997 Standards. The 1999 and later race-specific estimates are for persons who reported only one racial group; the category "2 or more races" includes persons who reported more than one racial group. Prior to data year 1999, data were tabulated according to 1977 Standards with four racial groups and the category "Asian only" included Native Hawaiian and Other Pacific Islander. Estimates for single race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. The effect of the 1997 Standard on the 1999 estimates can be seen by comparing 1999 data tabulated according to the two Standards: Age-adjusted estimates based on the 1977 Standards of the percent with Medicaid are: 0.1 percentage points higher for the Al/AN group than estimates based on the 1997 Standards. See Appendix II, Race.

⁵Missing family income data were imputed for 15–17 percent of the sample under 65 years of age in 1994–96. Percent of poverty level was unknown for 19 percent of sample persons under 65 in 1997, 24 percent in 1998, 27 percent in 1999, and 26 percent in 2000 and 2001. See Appendix II, Family income; Poverty level. ⁶MSA is metropolitan statistical area.

NOTES: Medicaid includes other public assistance through 1996. Starting in 1997 includes state-sponsored health plans. Starting in 1999 includes State Children's Health Insurance Program (SCHIP). In 2001, 7.9 percent were covered by Medicaid, 1.2 percent by state-sponsored health plans, and 1.2 percent by SCHIP.

SOURCES: Centers for Disease Control and Prevention, National Center for Health Statistics, National Health Interview Survey, health insurance supplements (1984, 1989, 1994–1996). Starting in 1997 data are from the family core questionnaires.