

WARNING!!! WARNING!!! WARNING!!! WARNING!!!

THIS TEXT DOCUMENT IS OVER 200 PAGES LONG!!!!!!

DO NOT PRINT THE DOCUMENT UNLESS YOU ARE ABLE
TO PRINT THIS MANY PAGES ON YOUR PRINTER!

YOU ARE ADVISED TO DOWNLOAD THE DOCUMENT TO
YOUR PERSONAL COMPUTER AND THEN VIEW OR
PRINT IT FROM WITHIN A WORD PROCESSOR!!!!
SET YOUR MARGINS TO ZERO AND USE A FONT
THAT MIMICS A TEXT PRINTER.

WARNING!!! WARNING!!! WARNING!!! WARNING!!!

Third National Health and Nutrition Examination Survey
(NHANES III), 1988-94

Catalog Number 76300

NHANES III LABORATORY DATA FILE DOCUMENTATION

Ages one year and older

December 1996

Table of Contents

| | |
|---|--|
| Introduction | |
| Guidelines for Data Users. | |
| Survey Description | |
| Sample Design and Analysis Guidelines. | |
| Data Preparation and Processing Procedures | |
| General References | |
| NHANES III Laboratory Data | |
| General Information | |
| Data File Index | |
| Data File Item Descriptions, Codes, Counts, and Notes | |
| References | |

Introduction

The National Center for Health Statistics (NCHS) of the Centers for Disease Control and Prevention (CDC) collects, analyzes, and disseminates data on the health status of U.S. residents. The results of surveys, analyses, and studies are made known through a number of data release mechanisms including publications, mainframe computer data files, CD-ROMs (Search and Retrieval Software, Statistical Export and Tabulation System (SETS)), and the Internet (<http://www.cdc.gov/nchswww/nchshome.htm>).

The National Health and Nutrition Examination Survey (NHANES) is a periodic survey conducted by NCHS. The third National Health and Nutrition Examination Survey (NHANES III), conducted from 1988 through 1994, was the seventh in a series of these surveys based on a complex, multi-stage sample plan. It was designed to provide national estimates of the health and nutritional status of the United States' civilian, noninstitutionalized population aged two months and older.

Data from NHANES III are being released in five public release data files:

NHANES III Household Adult Data File (Catalog Number 77560)

NHANES III Household Youth Data File (Catalog Number 77550)

NHANES III Examination Data File (Catalog Number 76200)

NHANES III Laboratory Data File (Catalog Number 76300)

NHANES III Dietary Recall Data Files (Catalog Number 76700)

A table showing the location of the interview and examination components in the five NHANES III public release data files follows.

Location of the interview and examination components in the five NHANES III public release data files

Data File

| Topic | HA | HY | EXAM | LAB | DIET |
|---|----|----|------|-----|------|
| Sample weights | X | X | X | X | . |
| Age/race/sex | X | X | X | X | . |
| Ethnic background | X | X | . | . | . |
| Household composition | X | X | . | . | . |
| Individual characteristics | X | X | . | . | . |
| Health insurance | X | X | . | . | . |
| Family background | X | X | . | . | . |
| Occupation of family head | X | X | . | . | . |
| Housing characteristics | X | X | . | . | . |
| Family characteristics | X | X | . | . | . |
| Orientation | X | X | . | . | . |
| Health services | X | X | . | . | . |
| Selected health conditions | X | X | X | . | . |
| Diabetes questions | X | . | . | . | . |
| High blood pressure and cholesterol questions | X | . | . | . | . |
| Cardiovascular disease questions | X | . | . | . | . |
| Musculoskeletal conditions | X | . | . | . | . |
| Physical functioning questions | X | . | . | . | . |
| Gallbladder disease questions | X | . | . | . | . |

Location of the interview and examination components in the five NHANES III public release data files (continued)

Data File

| Topic | HA | HY | EXAM | LAB | DIET |
|-----------------------------------|----|----|------|-----|------|
| Kidney conditions | X | . | . | . | . |
| Respiratory and allergy questions | X | X | . | . | . |
| Diet questions | X | . | . | . | . |
| Food frequency | X | . | X | . | . |
| Vision questions | X | X | . | . | . |
| Hearing questions | X | X | . | . | . |
| Dental care and status | X | X | . | . | . |
| Tobacco | X | . | X | . | . |
| Occupation | X | . | . | . | . |
| Language usage | X | X | . | . | . |
| Exercise | X | . | . | . | . |
| Social support/residence | X | . | . | . | . |
| Vitamin/mineral/medicine usage | X | X | X | . | . |
| Blood pressure measurement | X | . | X | . | . |
| Birth | . | X | X | . | . |
| Infant feeding practices/diet | . | X | . | . | . |
| Motor and social development | . | X | . | . | . |
| Functional impairment | X | X | . | . | . |
| School attendance | . | X | . | . | . |
| Cognitive function | . | X | X | . | . |

Location of the interview and examination components in the five NHANES III public release data files (continued)

Data File

| Topic | HA | HY | EXAM | LAB | DIET |
|---|----|----|------|-----|------|
| Alcohol and drug use | . | . | X | . | . |
| Reproductive health | . | . | X | . | . |
| Diagnostic interview schedule | . | . | X | . | . |
| Activity | . | . | X | . | . |
| Physician's examination | . | . | X | . | . |
| Height and weight | . | . | X | . | . |
| Body measurements | . | . | X | . | . |
| Dental examination | . | . | X | . | . |
| Allergy skin test | . | . | X | . | . |
| Audiometry | . | . | X | . | . |
| Tympanometry | . | . | X | . | . |
| WISC and WRAT | . | . | X | . | . |
| Spirometry | . | . | X | . | . |
| Bone densitometry | . | . | X | . | . |
| Gallbladder ultrasonography | . | . | X | . | . |
| Central nervous system function evaluation | . | . | X | . | . |
| Fundus photography | . | . | X | . | . |
| Physical function evaluation | . | . | X | . | . |
| Fasting questions | . | . | . | X | . |

Location of the interview and examination components in the five NHANES III public release data files (continued)

Data File

| Topic | HA | HY | EXAM | LAB | DIET |
|-------------------------------------|----|----|------|-----|------|
| Laboratory tests on blood and urine | . | . | . | X | . |
| Total nutrient intakes | . | . | X | . | . |
| Individual foods | . | . | . | . | X |
| Combination foods | . | . | . | . | X |
| Ingredients | . | . | . | . | X |

Data File Definitions

- HA - Household Adult Data File
- HY - Household Youth Data File
- EXAM - Examination Data File
- LAB - Laboratory Data File
- DIET - Dietary Recall Data Files

This document includes the documentation for the NHANES III Laboratory Data File and also contains a general overview of the survey and the use of the data files. The general overview includes five sections. The first section, entitled "Guidelines for Data Users," contains important information about the use of the data files. The second section, "Survey Description," is a brief overview of the survey plan and operation. The third section, "Sample Design and Analysis Guidelines," describes some technical aspects of the sampling plan and discusses some analytic issues particularly related to the use of data from complex sample surveys. The "Data Preparation and Processing Procedures" section describes the editing conventions and the codes used to represent the data. The last and fifth section, "General References," includes a reference list for the survey overview sections of the document.

Public Use Data Files for the third National Health and Nutrition Examination Survey will also be available from the National Technical Information Service (NTIS). A list of NCHS public use data tapes available for purchase from NTIS may be obtained from the Data Dissemination Branch at NCHS. Information regarding a bibliography (on disk) of journal articles citing data from all the NHANES and the availability of NHANES III data in CD-ROM/SETS software format can be obtained from the Data Dissemination Branch(301-436-8500) or by writing to:

Data Dissemination Branch
National Center for Health Statistics
Room 1018
6525 Belcrest Road
Hyattsville, Maryland 20782

NTIS can be contacted at:

NTIS - Computer Products Office
5285 Port Royal Road
Springfield, Virginia 22161
(703) 487-4807

Copies of all NHANES III questionnaires and data collection forms are included in the Plan and Operation of the Third National Health and Nutrition Examination Survey, 1988-94 (NCHS, 1994; U.S. DHHS, 1996). This publication, along with detailed information on NHANES procedures, interviewing, data collection, quality control techniques, survey design, nonresponse, and sample weighting can be found on the NHANES III Reference Manuals and Reports CD-ROM (U.S. DHHS, 1996). Information on how to order this CD-ROM is available from the Data Dissemination Branch at NCHS at the address and telephone number given above.

GUIDELINES FOR DATA USERS

Please refer to the following important information before analyzing data.

NHANES III Background Documents

- o The Plan and Operation of the Third National Health and Nutrition Examination Survey, 1988-94, (NCHS, 1994; U.S. DHHS, 1996) provides an overview of the survey and includes copies of the survey forms.
- o The sample design, nonresponse, and analytic guidelines documents on the NHANES III Reference Manuals and Reports CD-ROM (U.S. DHHS, 1996) discuss the reasons that sample weights and the complex survey design should be taken into account when conducting any analysis.
- o Instruction manuals, laboratory procedures, and other NHANES III reference manuals on the NHANES III Reference Manuals and Reports CD-ROM (U.S. DHHS, 1996) are also available for further information on the details of the survey.

Analytic Data Set Preparation

- o Most NHANES III survey design and demographic variables are found only on the Adult and Youth Household Data Files. In preparing a data set for analysis, other data files must be merged with either or both of these files to obtain many important analytic variables.
- o All of the NHANES III public use data files are linked with the common survey participant identification number (SEQN). Merging information from multiple NHANES III data files using this variable ensures that the appropriate information for each survey participant is linked correctly.
- o NHANES III public use data files do not have the same number of records on each file. The Household Questionnaire Files (divided into two files, Adult and Youth) contain more records than the Examination Data File because not everyone who was interviewed completed the examination. The Laboratory Data File contains data only for persons aged one year and older. The Individual Foods Data File based on the dietary recall has multiple records for each person rather than the one record per sample person contained in the other data files.
- o For each data file, SAS program code with standard variable names and labels is provided as separate text files on the CD-ROM that contains the data files. This SAS program code can be used to create a SAS data set from the data file.
- o Modifications were made to items in the questionnaires, laboratory, and examination components over the course of the survey; as a result, data may not be available for certain variables for the full six years. In addition, variables may differ by phase since some changes were implemented between phases. Users are encouraged to read the Notes sections of this document carefully for information about changes.

- o Extremely high and low values have been verified whenever possible, and numerous consistency checks have been performed. Nonetheless, users should examine the range and frequency of values before analyzing data.
- o Some data were not ready for release at the time of this publication due to continued processing of the data or analysis of laboratory specimens. A listing of those data are available in the general information section of each data file.
- o Confidential and administrative data are not being released to the public. Additionally, some variables have been recoded to help protect the confidentiality of the survey participants. For example, all age-related variables were recoded to 90+ years for persons who were 90 years of age and older.
- o Some variable names may differ from those used in the Phase 1 NHANES III Provisional Data Release and some variables included in the Phase 1 provisional release may not appear on these files.
- o Although the data files have been edited carefully, errors may be detected. Please notify NCHS staff (301-436-8500) of any errors in the data file or the documentation.

Analytic Considerations

- o NHANES III (1988-94) was designed so that the survey's first three years, 1988-91, its last three years, 1991-94, and the entire six years were national probability samples. Analysts are encouraged to use all six years of survey results.
- o Sample weights are available for analyzing NHANES III data. One of the following three sample weights will be appropriate for nearly all analyses: interviewed sample final weight (WTPFQX6), examined sample final weight (WTPFEX6), and mobile examination center (MEC)- and home-examined sample final weight (WTPFHX6). Choosing which of these sample weights to use in any analysis depends on the variables being used. A good rule of thumb is to use "the least common denominator" approach. In this approach, the user checks the variables of interest. The variable that was collected on the smallest number of persons is the "least common denominator," and the sample weight that applies to that variable is the appropriate one to use for that analysis. For more detailed information, see the Analytic and Reporting Guidelines for NHANES III (U.S. DHHS, 1996).

Referencing or Citing NHANES III Data

- o In publications, please acknowledge NCHS as the original data source. For instance, the reference for the NHANES III Laboratory Data File is:

U.S. Department of Health and Human Services (DHHS). National Center

for Health Statistics. Third National Health and Nutrition Examination Survey, 1988-1994, NHANES III Laboratory Data File (CD-ROM). Public Use Data File Documentation Number 76200. Hyattsville, MD.: Centers for Disease Control and Prevention, 1996. Available from National Technical Information Service (NTIS), Springfield, VA. Acrobat. PDF format; includes access software: Adobe Systems, Inc. Acrobat Reader 2.1.

- o Please place the acronym "NHANES III" in the titles or abstracts of journal articles and other publications in order to facilitate the retrieval of such materials in bibliographic searches.

SURVEY DESCRIPTION

The third National Health and Nutrition Examination Survey (NHANES III) was the seventh in a series of large health examination surveys conducted in the United States beginning in 1960. Three of these surveys, the National Health Examination Surveys (NHES), were conducted in the 1960's (NCHS, 1965; NCHS, 1967; NCHS, 1969). In 1970, an expanded nutrition component was added to provide data with which to assess nutritional status and dietary practices, and the name was changed to the National Health and Nutrition Examination Survey (Miller, 1973; Engel, 1978; McDowell, 1981). A special survey of Hispanic populations in the United States was conducted during 1982-1984 (NCHS, 1985).

The general structure of the NHANES III sample design was similar to that of the previous NHANES. All of the surveys used complex, multi-stage, stratified, clustered samples of civilian, noninstitutionalized populations. NHANES III was the first NHANES without an upper age limit; in fact, the age range for the survey was two months and older. A home examination option was employed for the first time in order to obtain examination data for very young children and for elderly persons who were unable to visit the mobile examination center (MEC). The home examination included only a subset of the components used in the full MEC examination since it would have been difficult to collect some types of data in a home setting. A detailed description of design specifications and copies of the data collection forms can be found in the Plan and Operation of the Third National Health and Nutrition Examination Survey, 1988-1994 (NCHS, 1994; U.S. DHHS, 1996).

NHANES III was conducted from October 1988 through October 1994 in two phases, each of which comprised a national probability sample. The first phase was conducted from October 18, 1988, through October 24, 1991, at 44 locations. The second phase was conducted from September 20, 1991, through October 15, 1994, at 45 different locations. In NHANES III, 39,695 persons were selected over the six years; of those, 33,994 (86%) were interviewed in their homes. All interviewed persons were invited to the MEC for a medical examination. Seventy-eight percent (30,818) of the selected persons were examined in the MEC, and an additional 493 persons were given a special, limited examination in their homes.

Data collection began with a household interview. Several questionnaires were administered in the household: Household Screener Questionnaire, Family Questionnaire, Household Adult Questionnaire, and Household Youth Questionnaire.

At the MEC, an examination was performed, and five automated questionnaires or interviews were administered: MEC Adult Questionnaire, MEC Youth Questionnaire, MEC Proxy Questionnaire, 24-Hour Dietary Recall, and Dietary Food Frequency (ages 12-16 years). The health examination component included a variety of tests and procedures. The examinee's age at the time of the interview and other factors determined which procedures were administered. Blood and urine specimens were obtained, and a number of tests and measurements were performed including body measurements, spirometry, fundus photography, x-rays, electrocardiography, allergy and glucose tolerance tests, and ultrasonography. Measurements were taken of bone density, hearing, and physical, cognitive, and central nervous system functions. A physician performed a limited standardized medical examination

and a dentist performed a standardized dental examination. While some of the blood and urine analyses were performed in the MEC laboratory, most analyses were conducted elsewhere by contract laboratories.

A home examination was conducted for those sample persons aged 2-11 months and aged 20 years or older who were unable to visit the mobile examination center. The home examination consisted of an abbreviated version of the tests and interviews performed in the MEC. Depending on age of the sample person, the components included body measurements, blood pressure, spirometry, venipuncture, physical function evaluation, and a questionnaire to inquire about infant feeding, selected health conditions, cognitive function, tobacco use, and reproductive history.

SAMPLE DESIGN AND ANALYSIS GUIDELINES

Sample Design

The general structure of the NHANES III sample design is the same as that of the previous NHANES. Each of these surveys used a stratified, multi-stage probability design. The major design parameters of the two previous NHANES and the special Hispanic HANES, as well as NHANES III, have been previously summarized (Miller, 1973; McDowell, 1981; NCHS, 1985; NCHS, 1994). The NHANES III sample was designed to be self-weighting within a primary sampling unit (PSU) for subdomains (age, sex, and race-ethnic groups). While the sample was fairly close to self-weighting nationally for each of these subdomain groups, it was not representative of the total population, which includes institutionalized, non-civilian persons that were outside the scope of the survey.

The NHANES III sample represented the total civilian, noninstitutionalized population, two months of age or over, in the 50 states and the District of Columbia of the United States. The first stage of the design consisted of selecting a sample of 81 PSU's that were mostly individual counties. In a few cases, adjacent counties were combined to keep PSU's above a minimum population size. The PSU's were stratified and selected with probability proportional to size (PPS). Thirteen large counties (strata) were chosen with certainty (probability of one). For operational reasons, these 13 certainty PSU's were divided into 21 survey locations. After the 13 certainty strata were designated, the remaining PSU's in the United States were grouped into 34 strata, and two PSU's were selected per stratum (68 survey locations). The selection was done with PPS and without replacement. The NHANES III sample therefore consists of 81 PSU's or 89 locations.

The 89 locations were randomly divided into two groups, one for each phase. The first group consisted of 44 and the other of 45 locations. One set of PSU's was allocated to the first three-year survey period (1988-91) and the other set to the second three-year period (1991-94). Therefore, unbiased estimates (from the point of view of sample selection) of health and nutrition characteristics can be independently produced for both Phase 1 and Phase 2 as well as for both phases combined.

For most of the sample, the second stage of the design consisted of area segments composed of city or suburban blocks, combinations of blocks, or other area segments in places where block statistics were not produced in the 1980 Census. In the first phase of NHANES III, the area segments were used only for a sample of persons who lived in housing units built before 1980. For units built in 1980 and later, the second stage consisted of sets of addresses selected from building permits issued in 1980 or later. These are referred to as "new construction segments." In the second phase, 1990 Census data and maps were used to define the area segments. Because the second phase followed within a few years of the 1990 Census, new construction did not account for a significant part of the sample, and the entire sample came from the area segments.

The third stage of sample selection consisted of households and certain types of group quarters, such as dormitories. All households and eligible

group quarters in the sample segments were listed, and a subsample was designated for screening to identify potential sample persons. The subsampling rates enabled production of a national, approximately equal-probability sample of households in most of the United States with higher rates for the geographic strata with high Mexican-American populations. Within each geographic stratum, there was a nearly equal-probability sample of households across all 89 stands.

Persons within the sample of households or group quarters were the fourth stage of sample selection. All eligible members within a household were listed, and a subsample of individuals was selected based on sex, age, and race or ethnicity. The definitions of the sex, age, race or ethnic classes, subsampling rates, and designation of potential sample persons within screened households were developed to provide approximately self-weighting samples for each subdomain within geographic strata and at the same time to maximize the average number of sample persons per sample household. Previous NHANES indicated that this increased the overall participation rate. Although the exact sample sizes were not known until data collection was completed, estimates were made. Below is a summary of the sample sizes for the full six-year NHANES III at each stage of selection:

| | |
|--|--------|
| Number of PSU's | 81 |
| Number of stands (survey locations) | 89 |
| Number of segments | 2,144 |
| Number of households screened | 93,653 |
| Number of households with sample persons | 19,528 |
| Number of designated sample persons | 39,695 |
| Number of interviewed sample persons | 33,994 |
| Number of MEC-examined sample persons | 30,818 |
| Number of home-examined sample persons | 493 |

More detailed information on the sample design and weighting and estimation procedures for NHANES III can be found in the Plan and Operation of the Third National Health and Nutrition Examination Survey, 1988-94 (NCHS, 1994; U.S. DHHS, 1996) and in the Analytic and Reporting Guidelines: Third National Health and Nutrition Examination Survey (NHANES III), 1988-94 (U.S. DHHS, 1996).

Analysis Guidelines

Because of the complex survey design used in NHANES III, traditional methods of statistical analysis based on the assumption of a simple random sample are not applicable. Detailed descriptions of this issue and possible analytic methods for analyzing NHANES data have been described earlier (NCHS, 1985; Yetley, 1987; Landis, 1982; Delgado, 1990). Recent analytic and reporting guidelines that should be used for most NHANES III analyses and publications are contained in Analytic and Reporting Guidelines (U.S. DHHS, 1996). These recommendations differ slightly from those used by analysts for previous NHANES surveys. These suggested guidelines provide a framework to users for producing estimates that conform to the analytic design of the survey. All users are strongly urged to review these analytic and reporting guidelines before beginning any analyses of NHANES III data.

It is important to remember that this set of statistical guidelines is not absolute. When conducting analyses, the analyst needs to use his/her subject matter knowledge (including methodological issues) as well as information about the survey design. The more one deviates from the original analytic categories defined in the sample design, the more important it is to evaluate the results carefully and to interpret the findings cautiously.

In NHANES III, 89 survey locations were randomly divided into two sets or phases, the first consisting of 44 and the other of 45 locations. One set of PSU's was allocated to the first three-year survey period (1988-91) and the other set to the second three-year period (1991-94). Therefore, unbiased national estimates of health and nutrition characteristics can be independently produced for each phase as well as for both phases combined. Computation of national estimates from both phases combined (i.e., total NHANES III) is the preferred option; individual phase estimates may be highly variable. In addition, individual phase estimates are not statistically independent. It is also difficult to evaluate whether differences in individual phase estimates are real or due to methodological differences. That is, differences may be due to changes in sampling methods or data collection methodology over time. At this time, there is no valid statistical test for examining differences between Phase 1 and Phase 2. Therefore, although point estimates can be produced separately for each phase, no test is available to test whether those estimates are significantly different from each other.

NHANES III is based on a complex, multi-stage probability sample design. Several aspects of the NHANES design must be taken into account in data analysis, including the sample weights and the complex survey design. Appropriate sample weights are needed to estimate prevalence, means, medians, and other statistics. Sample weights are used to produce correct population estimates because each sample person does not have the same probability of selection. The sample weights incorporate the differential probabilities of selection and include adjustments for noncoverage and nonresponse. A detailed discussion of nonresponse adjustments and issues related to survey coverage have been published (U.S. DHHS, 1996). With the large oversampling of young children, older persons, black persons, and Mexican-Americans in NHANES III, it is essential that the sample weights be used in all analyses. Otherwise, a misinterpretation of results is highly likely. Other aspects of the design that must be taken into account in data analyses are the strata and PSU pairings from the sample design. These pairings should be used to estimate variances and test for statistical significance. For weighted analyses, analysts can use special computer software packages that use an appropriate method for estimating variances for complex samples such as SUDAAN (Shah, 1995) and WesVarPC (Westat, 1996).

Although initial exploratory analyses may be performed on unweighted data using standard statistical packages and assuming simple random sampling, final analyses should be done on weighted data using appropriate sample weights. A summary of the weighting methodology and the type of sample weights developed for NHANES III is included in Weighting and Estimation Methodology (U.S. DHHS, 1996).

The purpose of weighting the sample data is to permit analysts to produce estimates of statistics that would have been obtained if the entire sampling frame (the United States) had been surveyed. Sample weights can be considered as measures of the number of persons the particular sample

observation represents. Weighting takes into account several features of the survey: the specific probabilities of selection for the individual domains that were oversampled as well as nonresponse and differences between the sample and the total U.S. population. Differences between the sample and the population may arise due to sampling variability, differential undercoverage in the survey among demographic groups, and possibly other types of response errors, such as differential response rates or misclassification errors. Sample weighting in NHANES III was used to:

1. Compensate for differential probabilities of selection among subgroups (i.e., age-sex-race-ethnicity subdomains where persons living in different geographic strata were sampled at different rates);
2. Reduce biases arising from the fact that nonrespondents may be different from those who participate;
3. Bring sample data up to the dimensions of the target population totals;
4. Compensate, to the extent possible, for inadequacies in the sampling frame (resulting from omissions of some housing units in the listing of area segments, omissions of persons with no fixed address, etc.); and
5. To reduce variances in the estimation procedure by using auxiliary information that is known with a high degree of accuracy.

In NHANES III, the sample weighting was carried out in three stages. The first stage involved the computation of weights to compensate for unequal probabilities of selection (objective 1, above). The second stage adjusted for nonresponse (objective 2). The third stage used poststratification of the sample weights to Census Bureau estimates of the U.S. population to accomplish the third, fourth, and fifth objectives simultaneously. In NHANES III, several types of sample weights (see the sample weights table that follows) were computed for the interviewed and examined sample and are included in the NHANES III data file. Also, sample weights were computed separately for Phase 1 (1988-91), Phase 2 (1991-94), and total NHANES III (1988-94) to facilitate analysis of items collected only in Phase 1, only in Phase 2, and over six years of the survey. Three sets of pseudo strata and PSU pairings are provided to use with SUDAAN in variance estimation. Since NHANES III is based on a complex, multi-stage sample design, appropriate sample weights should be used in analyses to produce national estimates of prevalence and associated variances while accounting for unequal probability of selection of sample persons. For example, the final interview weight, WTPFQX6, should be used for analysis of the items or questions from the family or household questionnaires, and the final MEC examination weight, WTPFEX6, should be used for analysis of the questionnaires and measurements administered in the MEC. Furthermore, for a combined analysis of measurements from the MEC examinations and associated medical history questions from the household interview, the final MEC examination weight, WTPFEX6, should be used. We recommend using SUDAAN (Shah, 1995) to estimate statistics of interest and the associated variance. However, one can also use other published methods for variance estimation. Application of SUDAAN and alternative methods, such as the average design effect approach, balance repeated replication (BRR) methods, or jackknife methods for variance estimation, are discussed in Weighting and Estimation Methodology (U.S. DHHS, 1996).

Appropriate Uses of the NHANES III Sample Weights

Final interview weight, WTPFQX6

Use only in conjunction with the sample interviewed at home and with items collected during the household interview.

Final examination (MEC only) weight, WTPFEX6

Use only in conjunction with the MEC-examined sample and with interview and examination items collected at the MEC.

Final MEC+home examination weight, WTPFHX6

Use only in conjunction with the MEC+home-examined sample and with items collected at both the MEC and home.

Final allergy weight, WTPFALG6

Use only in conjunction with the allergy subsample and with items collected as part of the allergy component of the exam.

Final CNS weight, WTPFCNS6

Use only in conjunction with the CNS subsample and with items collected as part of the CNS component of the exam.

Final morning examination (MEC only) subsample weight, WTPFSD6

Use only in conjunction with the MEC-examined persons assigned to the morning subsample and only with items collected in the MEC exam.

Final afternoon/evening examination (MEC only) subsample weight, WTPFMD6

Use only in conjunction with the MEC-examined persons assigned to the afternoon/evening subsample and only with items collected in the MEC exam.

Final morning examination (MEC+home) subsample weight, WTPFHSD6

Use only in conjunction with the MEC- and home-examined persons assigned to the morning subsample and with items collected during the MEC and home examinations.

Final afternoon/evening examination (MEC+home) weight, WTPFHMD6

Use only in conjunction with the MEC- and home-examined persons assigned to the afternoon/evening subsample and with items collected during the MEC and home examinations.

DATA PREPARATION AND PROCESSING PROCEDURES

Automated data collection procedures for the survey were introduced in NHANES III. In the mobile examination centers, data for the interview and examination components were recorded directly onto a computerized data collection form. With the exception of a few independently automated systems, the system was centrally integrated. This operation allowed for ongoing monitoring of much of the data. Before the introduction of the computer-assisted personal interview (CAPI), the household questionnaire data were reviewed manually by field editors and interviewers. CAPI (1992-1994 only) questionnaires featured built-in edits to prevent entering inconsistencies and out-of-range responses. The multi-level data collection and quality control systems are discussed in detail in the Plan and Operation of the Third National Health and Nutrition Examination Survey, 1988-1994 (NCHS, 1994; U.S. DHHS, 1996). All interview, laboratory, and examination data were sent to NCHS for final processing.

Guidelines were developed that provided standards for naming variables, filling missing values and coding conventional responses, handling missing records, and standardizing two-part quantity/unit questionnaire variables. NCHS staff, assisted by contract staff, developed data editing specifications that checked data sets for valid codes, ranges, and skip pattern inconsistencies and examined the consistency of values between interrelated variables. Comments, collected in both interviews and examination components, were reviewed and recoded when possible. Responses to "Other" and "Specify" were recoded either to existing code categories or to new categories. The documentation for each data set includes notes for those variables that have been recoded and standardized and for those variables that differ significantly from what appears in the original data collection instrument. While the data have undergone many quality control and editing procedures, there still may be values that appear extreme or illogical. Values that varied considerably from what was expected were examined by analysts who checked for comments or other responses that might help to clarify unusual values. Generally, values were retained unless they could not possibly be true, in which case they were changed to "Blank but applicable." Therefore, the user must review each data set for extreme or inconsistent values and determine the status of each value for analysis.

Several editing conventions were used in the creation of final analytic data sets:

1. Standardized variables were created to replace all two-part quantity/unit questions using standard conversion factors. Standardized variables have the same name as the variable of the two-part question with an "S" suffix. For instance, MAPF18S (Months received WIC benefits) in the MEC Adult Questionnaire was created from the two-part response option to question F18, "How long did you receive benefits from the WIC program?," using the conversion factor 12 months per year.

2. Recoded variables were created by combining responses from two or more like variables, or by collapsing responses to create a summary variable for the purpose of confidentiality. Recoded variables have the original variable name with an R suffix. For example, place of birth variable (HFA6X) in the Family Questionnaire was collapsed to a three level response category (U.S., Mexico, Other) and renamed HFA6XR. Generally, only the recoded variable has been included in the data file.
3. Fill values, a series of one or more digits, were used to represent certain specific conditions or responses. Below is a list of the fill values that were employed. Some of the fill values pertain only to questionnaire data, although 8-fill and blank-fill values are found in all data sets. Other fill values, not included in this list, are used to represent component-specific conditions.

6-fills = Varies/varied. (Questionnaires only)

7-fills = Fewer than the smallest number that could be reported within the question structure (e.g., fewer than one cigarette per day). (Questionnaires only)

8-fills = Blank but applicable/cannot be determined. This means that a respondent was eligible to receive the question, test, or component but did not because of refusal, lack of time, lack of staff, loss of data, broken vial, language barrier, unreliability, or other similar reasons.

9-fills = Don't know. This fill was used only when a respondent did not know the response to a question and said, "I don't know." (Questionnaires only)

Blank fills = Inapplicable. If a respondent was not eligible for a questionnaire, test, or component because of age, gender, or specific reason, the variable was blank-filled. In the questionnaire, if a respondent was not asked a question because of a skip-pattern, variables corresponding to the question were blank-filled. For examination or laboratory components, if a person was excluded by a defined protocol (e.g., screening exclusion questions) and these criteria are included in the data set, then the corresponding variables were blank-filled for that person. For home examinees, variables for examination components and blood tests not performed as part of the home examination protocol were blank-filled.

4. For variables describing discrete data, codes of zero (0) were used to mean "none," "never," or the equivalent. Value labels for which "0" is used include: "has not had," "never regularly," "still taking," or "never stopped using." Unless otherwise labeled, for variables containing continuous data, "zero" means "zero."
5. Where there are logical skip patterns in the flow of the questionnaire or examination component, the skip was indicated by placing the variable label of the skip destination in parentheses as part of the value label of the response generating the skip. For example, in the Physical Function Evaluation, the variable PFPWC (in wheelchair) has a value label, "2 No (PFPSCOOT)" that means that the next item for persons not in a wheelchair would be represented by the variable, PFPSCOOT.

Variable Nomenclature

A unique name was assigned to every NHANES III variable using a standard convention. By following this naming convention, the origin of each variable is clear, and there is no chance of overlaying similar variables across multiple components. Variables range in length from three to eight characters. The first two variable characters represent the topic (e.g., analyte, questionnaire instrument, examination component) and are listed below alphabetically by topic. For questionnaires administered in the household, the remainder of the variable name following the first two

characters indicates the question section and number. For example, data for the response to the Household Adult Questionnaire question B1 are contained in the variable HAB1. For most laboratory and examination variables, as well as some other variables, a "P" in the third position refers to "primary" and the remainder of the variable name is a brief description of the item. For instance, in the Laboratory Data File, information on the length of time the person fasted before the first blood draw is contained in the variable PHPFAST. The variable PHPFAST was derived as follows: characters 1-2 (PH) refer to "phlebotomy," character 3 (P) refers to "primary," characters 4-8 (FAST) refer to an abbreviation for "fasting."

| CODE | TOPIC |
|------|--|
| AT | Alanine aminotransferase (from biochemistry profile) |
| AM | Albumin (from biochemistry profile) |
| AP | Alkaline phosphatase (from biochemistry profile) |
| AL | Allergy skin test |
| AC | Alpha carotene |
| AN | Anisocytosis |
| AA | Apolipoprotein (AI) |
| AB | Apolipoprotein (B) |
| AS | Aspartate aminotransferase (from biochemistry profile) |
| LA | Atypical lymphocyte |
| AU | Audiometry |
| BA | Band |
| BO | Basophil |
| BS | Basophilic stippling |
| BC | Beta carotene |
| BX | Beta cryptoxanthin |
| BL | Blast |
| BU | Blood urea nitrogen (BUN) (from biochemistry profile) |
| BM | Body measurements |
| BD | Bone densitometry |
| C1 | C-peptide (first venipuncture) |
| C2 | C-peptide (second venipuncture) |
| CR | C-reactive protein |
| UD | Cadmium |
| CN | Central nervous system function evaluation |
| CL | Chloride (from biochemistry profile) |
| CO | Cotinine |
| CE | Creatinine (serum)(from biochemistry profile) |

| CODE | TOPIC |
|------|--|
| UR | Creatinine (urine) |
| DM | Demographic |
| DE | Dental examination |
| MQ | Diagnostic interview schedule |
| DR | Dietary recall (total nutrient intakes) |
| EO | Eosinophil |
| EP | Erythrocyte protoporphyrin |
| FR | Ferritin |
| FB | Fibrinogen |
| RB | Folate (RBC) |
| FO | Folate (serum) |
| FH | Follicle stimulating hormone (FSH) |
| FP | Fundus photography |
| GG | Gamma glutamyl transferase (GGT) (from biochemistry profile) |
| GU | Gallbladder ultrasonography |
| GB | Globulin (from biochemistry profile) |
| G1 | Glucose (first venipuncture) |
| G2 | Glucose (second venipuncture) |
| SG | Glucose (from biochemistry profile) |
| GH | Glycated hemoglobin |
| GR | Granulocyte |
| C3 | HCO ₃ (Bicarbonate)(from biochemistry profile) |
| HD | HDL cholesterol |
| HP | Helicobacter pylori antibody |
| HT | Hematocrit |
| HG | Hemoglobin |
| AH | Hepatitis A antibody (HAV) |
| HB | Hepatitis B core antibody (anti-HBc) |
| SS | Hepatitis B surface antibody (anti-HBs) |
| SA | Hepatitis B surface antigen (HBsAg) |
| HC | Hepatitis C antibody (HCV) |
| DH | Hepatitis D antibody (HDV) |
| H1 | Herpes 1 antibody |
| H2 | Herpes 2 antibody |
| HX | Home examination (general) |
| HF | Household family questionnaire |
| HA | Household adult questionnaire |
| HQ | Household questionnaire variables (composite) |
| HS | Household screener questionnaire |
| HY | Household youth questionnaire |
| HZ | Hypochromia |
| I1 | Insulin (first venipuncture) |
| I2 | Insulin (second venipuncture) |
| UI | Iodine (urine) |
| FE | Iron |
| SF | Iron (from biochemistry profile) |
| LD | Lactate dehydrogenase (from biochemistry profile) |
| L1 | Latex antibody |
| LC | LDL cholesterol (calculated) |
| PB | Lead |
| LP | Lipoprotein (a) |
| LH | Luteinizing hormone |

| CODE | TOPIC |
|------|--|
| LU | Lutein/zeaxanthin |
| LY | Lycopene |
| LM | Lymphocyte |
| MR | Macrocyte |
| MC | Mean cell hemoglobin (MCH) |
| MH | Mean cell hemoglobin concentration (MCHC) |
| MV | Mean cell volume (MCV) |
| PV | Mean platelet volume |
| MA | MEC adult questionnaire |
| MX | MEC examination (general) |
| FF | Dietary food frequency (ages 12-16 years) |
| MP | MEC proxy questionnaire |
| MY | MEC youth questionnaire |
| ME | Metamyelocyte |
| MI | Microcyte |
| MO | Monocyte |
| MN | Mononuclear cell |
| ML | Myelocyte |
| IC | Normalized calcium (derived from ionized calcium) |
| OS | Osmolality (from biochemistry profile) |
| PH | Phlebotomy data collected in MEC (e.g., questions) |
| PS | Phosphorus (from biochemistry profile) |
| PF | Physical function evaluation |
| PE | Physician's examination |
| PL | Platelet |
| DW | Platelet distribution width |
| PK | Poikilocytosis |
| PO | Polychromatophilia |
| SK | Potassium (from biochemistry profile) |
| PR | Promyelocyte |
| RC | Red blood cell count (RBC) |
| RW | Red cell distribution width (RDW) |
| RE | Retinyl esters |
| RF | Rheumatoid factor antibody |
| RU | Rubella antibody |
| WT | Sample weights |
| SE | Selenium |
| SI | Sickle cell |
| NA | Sodium (from biochemistry profile) |
| SH | Spherocyte |
| SP | Spirometry |
| SD | Survey design |
| TT | Target cell |
| TE | Tetanus |
| TB | Total bilirubin (from biochemistry profile) |
| CA | Total calcium |
| SC | Total calcium (from biochemistry profile) |
| TC | Total cholesterol |
| CH | Total cholesterol (from biochemistry profile) |
| TI | Total iron binding capacity (TIBC) |
| TP | Total protein (from biochemistry profile) |
| TX | Toxic granulation |
| TO | Toxoplasmosis antibody |
| PX | Transferrin saturation |

| CODE | TOPIC |
|------|---|
| TG | Triglycerides |
| TR | Triglycerides (from biochemistry profile) |
| TY | Tympanometry |
| UA | Uric acid (from biochemistry profile) |
| UB | Urinary albumin |
| VU | Vacuolated cells |
| VR | Varicella antibody |
| VA | Vitamin A |
| VB | Vitamin B12 |
| VC | Vitamin C |
| VE | Vitamin E |
| WC | White blood cell count (WBC) |
| WW | WISC/WRAT cognitive test |

GENERAL REFERENCES

- Delgado JL, Johnson CL, Roy I, Trevino FM. Hispanic Health and Nutrition Examination Survey: methodological considerations. *Amer J Pub Health* 80(suppl.):6-10. 1990.
- Engel A, Murphy RS, Maurer K, Collins E. Plan and operation of the HANES I Augmentation Survey of Adults 25-74 Years, United States, 1974-75. National Center for Health Statistics. *Vital Health Stat* 1(14). 1978.
- Freeman DH, Freeman JL, Brock DB, Koch GG. Strategies in the multivariate analysis of data from complex surveys II: an application to the United States National Health Interview Survey. *Int Stat Rev* 40(3):317-30. 1976.
- Khare M, Mohadjer LK, Ezzati-Rice TM, Waksberg J. An evaluation of nonresponse bias in NHANES III (1988-91). 1994 Proceedings of the Survey Research Methods section of the American Statistical Association. 1994.
- Landis JR, Lepkowski JM, Eklund SA, Stehouwer SA. A statistical methodology for analyzing data from a complex survey, the first National Health and Nutrition Examination Survey. National Center for Health Statistics. *Vital Health Stat* 2(92). 1982.
- McDowell A, Engel A, Massey JT, Maurer K. Plan and operation of the second National Health and Nutrition Examination Survey, 1976-80. National Center for Health Statistics. *Vital Health Stat* 1(15). 1981.
- Miller HW. Plan and operation of the Health and Nutrition Examination Survey, United States, 1971-1973. National Center for Health Statistics. *Vital Health Stat* 1(10a) and (10b). 1973.
- National Center for Health Statistics. Plan and initial program of the Health Examination Survey. *Vital Health Stat* 1(4). 1965.
- National Center for Health Statistics. Plan and operation of a health examination survey of U.S. youths 12-17 years of age. *Vital Health Stat* 1(8). 1969.
- National Center for Health Statistics. Plan and operation of the Hispanic Health and Nutrition Examination Survey, 1982-84. *Vital Health Stat* 1(19). 1985.
- National Center for Health Statistics. Plan and operation of the Third National Health and Nutrition Examination Survey, 1988-94. *Vital Health Stat* 1(32). 1994.
- National Center for Health Statistics. Plan, operation, and response results of a program of children's examinations. *Vital Health Stat* 1(5). 1967.
- Shah BV, Barnwell BG, Bieler GS. SUDAAN User's Manual: Software for Analysis of Correlated Data. Research Triangle Park, NC: Research Triangle Institute. Release 6.04. 1995.

Skinner CJ. Aggregated analysis: standard errors and significance tests. In: Skinner CJ, Holt D, Smith TMF, eds. Analysis of complex surveys. New York: John Wiley and Sons, Inc. 1989.

U.S. Department of Health and Human Services (DHHS). National Center for Health Statistics. NHANES III reference manuals and reports (CD-ROM). Hyattsville, MD: Centers for Disease Control and Prevention, 1996. Available from National Technical Information Service (NTIS), Springfield, VA. Acrobat .PDF format; includes access software: Adobe Systems, Inc. Acrobat Reader 2.1.

Westat, Inc. A User's Guide to WesVarPC. Rockville, MD. Westat, Inc. 1996.

Yetley E, Johnson C. Nutritional applications of the Health and Nutrition Examination Surveys (HANES). Annu Rev Nutr 7:441-63. 1987.

NHANES III LABORATORY DATA FILE

General Information

Introduction

The Laboratory Data File contains data from the urine collection and venipuncture components of the examination, including almost all laboratory test results (blood and urine) available to date. The exceptions are discussed elsewhere in this documentation. In addition, auxiliary information such as how long the examinee fasted, the time of day of the venipuncture, and the conditions precluding venipuncture has been included. This documentation presents information that should be reviewed before proceeding with data analysis.

The documentation pertaining specifically to the Laboratory Data File is divided into four main sections. The first section, "General Information," provides information about the contents of the Laboratory Data File. The second section, "Data File Index," includes a brief description of all the variables on the data set and shows the standard name of each variable and its position in the data set. The third section, "Item Descriptions, Codes, Counts, and Notes" provides for each component a description, the standard variable name and a brief description of the values that variable can take on, a count of the frequency of occurrence of each value, notes by variable, and appendices as necessary. "References" are provided in the fourth section.

Blood and urine specimens were collected on examinees aged one year and older at the mobile examination center (MEC). For those examinees aged one year and older who did not travel to the MEC, only blood specimens were collected during the Home Examination (HE). Hematologic profiles were completed for all examinees, and specified laboratory tests were performed upon each specimen based on the examinee's age and sex. Only a limited number of tests were performed on specimens collected during the Home Examination. Appendix 1 lists the laboratory tests by specimen type, age group, sex, and whether the specimen was collected in the Home Examination.

The analysis of NHANES III laboratory data must be conducted with the key survey design and basic demographic variables. The NHANES III Household Youth Questionnaire Data File (ages two months to 16 years) and the NHANES III Household Adult Questionnaire Data File (ages 17 years and older) contain demographic data, health indicators, and other related information collected during household interviews. They also contain all survey design variables and sample weights for these age groups. These two household questionnaire files may be linked to the laboratory data file using the unique survey participant (sample person) identifier SEQN.

Examinee Screening

Prior to the phlebotomy (venipuncture), a questionnaire was administered to determine an examinee's eligibility for all phlebotomy procedures (including venipuncture and the oral glucose tolerance test). It included questions to determine if it was safe to perform the venipuncture, to document and determine fasting compliance, and to aid in analyzing the results of the laboratory tests performed. Examinees reporting hemophilia or

recent cancer chemotherapy treatment were excluded from the venipuncture. For those examinees, the laboratory test results fields for all blood-based laboratory tests were left blank. Because examinees reporting current insulin therapy were excluded from receiving the oral glucose tolerance test (OGTT), the plasma glucose (G2P), serum insulin (I2P) and serum C-peptide (C2P) results from the second venipuncture were left blank as well.

Although examinees aged 12 years and older were instructed to fast for 10-16 hours prior to the morning examination or for six hours before the afternoon or evening examination, the instructions were not followed uniformly. Laboratory test results and the duration of the fast have been included on the data file regardless of the examinee's fasting compliance. Analysts should consider whether fasting status is crucial before undertaking analyses. Examinees who reported insulin use during the household interview were not instructed to fast.

Specimen Collection and Processing Procedures

Detailed specimen collection and processing instructions are discussed in the Manual for Medical Technicians (U.S. DHHS, 1996). Vials were stored under appropriate refrigerated (4-8 degrees Centigrade) or frozen (-20 degrees Centigrade) conditions until they were shipped to analytical laboratories for testing. The analytical methods used by each of the participating laboratories are described in the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996). The manual contains quality control graphs and statistical summary information for each laboratory test at the end of the laboratory method description.

Oral glucose tolerance testing: During NHANES III, the OGTT was conducted on MEC examinees aged 40-74 years. The protocol included two timed venipunctures and a glucose drink. Two glucose drinks were used to measure an examinee's ability to metabolize glucose -- Dextol(TM) and Trutol(TM). After the first venipuncture, the examinee drank the glucose drink, and a second venipuncture was performed approximately two hours later.

Examiner Training and Quality Control

The NHANES III laboratory staff consisted of medical technologists and phlebotomists. The medical technologists held baccalaureates in medical technology. Both they and the phlebotomists were certified by the American Society for Clinical Pathologists or by a similar organization.

All laboratory staff completed comprehensive training in standardized laboratory procedures before they began working in the MEC. The MEC phlebotomists completed comprehensive training in pediatric phlebotomy techniques, including instruction by a pediatric nurse practitioner. Laboratory team performance was monitored using several techniques. NCHS and contract consultants used a structured quality assurance evaluation during unscheduled visits to evaluate both the quality of the laboratory work and the quality-control procedures. Each laboratory staff person was observed for equipment operation, specimen collection and preparation, and testing procedures, and constructive feedback was given to each team. Formal retraining sessions were conducted annually to ensure that required skill levels were maintained.

Laboratory Protocol Changes from 1988 to 1994

Most laboratory tests were performed for the entire six years of NHANES III. Exceptions are detailed below. Apolipoprotein AI and B tests were included during 1988-1991 only. Lipoprotein(a), Vitamin B12, and antibody tests for immunoglobulin E, rubella, varicella, and toxoplasmosis were conducted during 1991-1994. For the 1991-1994 period, the OGTT procedure was modified to add tests for C-peptide and insulin on specimens from the second venipuncture. For statistical analyses of these laboratory test results, the appropriate Phase 1 or Phase 2 sample weight should be used.

Incomplete Data Release

At the time of this data release, some laboratory test results were not available. Tests for which results were unavailable included vitamin D, immunoglobulin E, diphtheria antitoxin, measles antibody, homocysteine, periodontal pathogens, thyroxine, thyroid stimulating hormone, antithyroglobulin antibody, antimicrosomal antibody, and methylmalonic acid. Cotinine test results for 1988-1991 have been included in this laboratory data file. Cotinine testing is still being carried out for 1991-1994, and the laboratory test results will be released at a future date. Results from urine pregnancy tests are included in the NHANES III Examination Data File, rather than in the Laboratory Data File.

Serologic testing for human immunodeficiency virus (HIV) antibody and urine testing for drugs of abuse were performed anonymously. The drugs of abuse for which examinees were tested were cocaine, marijuana, opiates, phencyclidine, and amphetamines. To maintain anonymity, the examinee's serum and urine were labeled with a random identifying number, and limited demographic data were linked to that number. The new identifier was not linked to the original sample identifier. Therefore, these data cannot be linked to other NHANES III data. The HIV test was performed from 1988 through 1994; the urine drug testing was done from 1991 through 1994. Because of the limited analytic potential of the HIV and drug data, this file is not included in this data release.

Data Preparation and Processing

For laboratory tests with a lower detection limit, results below the lower detection limit were replaced with a value equal to the detection limit divided by the square root of two. This value was created to help the user distinguish a nondetectable laboratory test result from a measured laboratory test result. Appendix 2 documents the detection limit for each laboratory test.

The SI unit (le Systeme International d Unites) is an outgrowth of the metric system that has been used throughout most of the world. In addition to providing a uniform international system of units of measurement, a uniform style is prescribed. Laboratory test results not originally reported in SI units were converted to SI units if applicable. Conversion factors, the format of the NHANES and SI results, and NHANES and SI units of measure are in Appendix 3. In converting NHANES III data to SI units, the goal was to preserve the level of detail reported by the laboratories in the original

laboratory test result. Therefore, the number of significant digits in the laboratory test results data may be different from that in published references.

The Laboratory Data File contains laboratory test results for glucose (G1P), triglycerides (TGP), cholesterol (TCP), and iron (FEP) measured by contract laboratories using reference analytic methods. For these methods, consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996). However, the biochemistry profile also included measurements of these analytes. In general, for most analyses, the appropriate variables to use are G1P, TGP, TCP and FEP. The values from the biochemistry profile (SGP, CHP, TRP, SFP) should not be used routinely.

The definition of a reference method by the National Committee for Clinical Laboratory Standards (NCCLS) is "a thoroughly investigated method in which exact and clear descriptions of the necessary conditions and procedures are given for the accurate determination of one or more property values; the documented accuracy and precision of the method are commensurate with the method's use for assessing the accuracy of other methods for measuring the same property values or for assigning reference method values to reference materials" (NCCLS, 1991).

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|---------------|-----------|
| ----- | | |
| DEMOGRAPHIC DATA | | |
| HOUSEHOLD SCREENER QUESTIONNAIRE (HSQ) | | |
| Sample person identification number | SEQN | 1-5 |
| Family sequence number | DMPFSEQ | 6-10 |
| Examination/interview Status | DMPSTAT | 11 |
| Race-ethnicity | DMARETHN | 12 |
| Race | DMARACER | 13 |
| Ethnicity | DMAETHNR | 14 |
| Sex | HSSEX | 15 |
| Age at interview (Screener) | HSAGEIR | 16-17 |
| Age at interview - unit (Screener) | HSAGEU | 18 |
| Age in months at interview (screener) | HSAITMOR | 19-22 |
| Family size (persons in family) | HSFSIZER | 23-24 |
| Household size (persons in dwelling) | HSHSIZER | 25-26 |
| County code | DMPCNTYR | 27-29 |
| FIPS code for State | DMPFIPSR | 30-31 |
| Rural/urban code based on USDA code | DMPMETRO | 32 |
| Census region, weighting(Texas in south) | DMPCREGN | 33 |
| Poverty Income Ratio (unimputed income) | DMPPIR | 34-39 |
| SURVEY DESIGN DATA | | |
| Phase of NHANES III survey | SDPPHASE | 40 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|---------------|-----------|
| Total NHANES III pseudo-PSU | SDPPSU6 | 41 |
| Total NHANES III pseudo-stratum | SDPSTRA6 | 42-43 |
| Pseudo-PSU for phase 1 | SDPPSU1 | 44 |
| Pseudo-stratum for phase 1 | SDPSTRA1 | 45-46 |
| Pseudo-PSU for phase 2 | SDPPSU2 | 47 |
| Pseudo-stratum for phase 2 | SDPSTRA2 | 48-49 |
| SAMPLING WEIGHTS - TOTAL NHANES III (1988-94) | | |
| Total interviewed sample final weight | WTPFQX6 | 50-58 |
| Total MEC-examined sample final weight | WTPFEX6 | 59-67 |
| Total M+H examined sample final weight | WTPFHX6 | 68-76 |
| Total allergy subsample final weight | WTPFALG6 | 77-85 |
| Total CNS subsample final weight | WTPFCNS6 | 86-94 |
| Total morning subsample final wgt | WTPFSD6 | 95-103 |
| Total afternoon/eve subsample final wgt | WTPFMD6 | 104-112 |
| Total M+H morning subsample final wgt | WTPFHSD6 | 113-121 |
| Total M+H afternoon subsample final wgt | WTPFHMD6 | 122-130 |
| SAMPLING WEIGHTS - NHANES III PHASE 1 (1988-91) | | |
| Phase 1 interviewed sample final wgt | WTPFQX1 | 131-139 |
| Phase 1 MEC examined sample final wgt | WTPFEX1 | 140-148 |
| Phase 1 M+H examined sample final wgt | WTPFHX1 | 149-157 |
| Phase 1 allergy subsample final wgt | WTPFALG1 | 158-166 |
| Phase 1 CNS subsample final wgt | WTPFCNS1 | 167-175 |
| Phase 1 morning sess subsample final wgt | WTPFSD1 | 176-184 |
| Phase 1 aft/eve subsample final wgt | WTPFMD1 | 185-193 |
| Phase 1 morning M+H subsample final wgt | WTPFHSD1 | 194-202 |
| Phase 1 aft/eve M+H subsample final wgt | WTPFHMD1 | 203-211 |
| SAMPLING WEIGHTS - NHANES III PHASE 2 (1991-94) | | |
| Phase 2 interviewed sample final wgt | WTPFQX2 | 212-220 |
| Phase 2 MEC examined sample final wgt | WTPFEX2 | 221-229 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|---------------|-----------|
| Phase 2 M+H examined sample final wgt | WTPFHX2 | 230-238 |
| Phase 2 allergy subsample final wgt | WTPFALG2 | 239-247 |
| Phase 2 CNS subsample final wgt | WTPFCNS2 | 248-256 |
| Phase 2 morning sess subsample final wgt | WTPFSD2 | 257-265 |
| Phase 2 aft/eve subsample final wgt | WTPFMD2 | 266-274 |
| Phase 2 morning M+H subsample final wgt | WTPFHSD2 | 275-283 |
| Phase 2 aft/eve M+H subsample final wgt | WTPFHMD2 | 284-292 |

FAY'S BRR REPLICATE INTERVIEW WEIGHTS - TOTAL NHANES III (1988-94)

| | | |
|---|----------|---------|
| Replicate 1 final interview weight | WTPQRP1 | 293-301 |
| Replicate 2 final interview weight | WTPQRP2 | 302-310 |
| Replicate 3 final interview weight | WTPQRP3 | 311-319 |
| Replicate 4 final interview weight | WTPQRP4 | 320-328 |
| Replicate 5 final interview weight | WTPQRP5 | 329-337 |
| Replicate 6 final interview weight | WTPQRP6 | 338-346 |
| Replicate 7 final interview weight | WTPQRP7 | 347-355 |
| Replicate 8 final interview weight | WTPQRP8 | 356-364 |
| Replicate 9 final interview weight | WTPQRP9 | 365-373 |
| Replicate 10 final interview weight | WTPQRP10 | 374-382 |
| Replicate 11 final interview weight | WTPQRP11 | 383-391 |
| Replicate 12 final interview weight | WTPQRP12 | 392-400 |
| Replicate 13 final interview weight | WTPQRP13 | 401-409 |
| Replicate 14 final interview weight | WTPQRP14 | 410-418 |
| Replicate 15 final interview weight | WTPQRP15 | 419-427 |
| Replicate 16 final interview weight | WTPQRP16 | 428-436 |
| Replicate 17 final interview weight | WTPQRP17 | 437-445 |
| Replicate 18 final interview weight | WTPQRP18 | 446-454 |
| Replicate 19 final interview weight | WTPQRP19 | 455-463 |
| Replicate 20 final interview weight | WTPQRP20 | 464-472 |
| Replicate 21 final interview weight | WTPQRP21 | 473-481 |
| Replicate 22 final interview weight | WTPQRP22 | 482-490 |
| Replicate 23 final interview weight | WTPQRP23 | 491-499 |
| Replicate 24 final interview weight | WTPQRP24 | 500-508 |
| Replicate 25 final interview weight | WTPQRP25 | 509-517 |
| Replicate 26 final interview weight | WTPQRP26 | 518-526 |
| Replicate 27 final interview weight | WTPQRP27 | 527-535 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|---------------|-----------|
| Replicate 28 final interview weight | WTPQRP28 | 536-544 |
| Replicate 29 final interview weight | WTPQRP29 | 545-553 |
| Replicate 30 final interview weight | WTPQRP30 | 554-562 |
| Replicate 31 final interview weight | WTPQRP31 | 563-571 |
| Replicate 32 final interview weight | WTPQRP32 | 572-580 |
| Replicate 33 final interview weight | WTPQRP33 | 581-589 |
| Replicate 34 final interview weight | WTPQRP34 | 590-598 |
| Replicate 35 final interview weight | WTPQRP35 | 599-607 |
| Replicate 36 final interview weight | WTPQRP36 | 608-616 |
| Replicate 37 final interview weight | WTPQRP37 | 617-625 |
| Replicate 38 final interview weight | WTPQRP38 | 626-634 |
| Replicate 39 final interview weight | WTPQRP39 | 635-643 |
| Replicate 40 final interview weight | WTPQRP40 | 644-652 |
| Replicate 41 final interview weight | WTPQRP41 | 653-661 |
| Replicate 42 final interview weight | WTPQRP42 | 662-670 |
| Replicate 43 final interview weight | WTPQRP43 | 671-679 |
| Replicate 44 final interview weight | WTPQRP44 | 680-688 |
| Replicate 45 final interview weight | WTPQRP45 | 689-697 |
| Replicate 46 final interview weight | WTPQRP46 | 698-706 |
| Replicate 47 final interview weight | WTPQRP47 | 707-715 |
| Replicate 48 final interview weight | WTPQRP48 | 716-724 |
| Replicate 49 final interview weight | WTPQRP49 | 725-733 |
| Replicate 50 final interview weight | WTPQRP50 | 734-742 |
| Replicate 51 final interview weight | WTPQRP51 | 743-751 |
| Replicate 52 final interview weight | WTPQRP52 | 752-760 |

FAY'S BRR REPLICATE EXAMINATION WEIGHTS - TOTAL NHANES III (1988-94)

| | | |
|-------------------------------------|---------|---------|
| Replicate 1 final exam weight | WTPXRP1 | 761-769 |
| Replicate 2 final exam weight | WTPXRP2 | 770-778 |
| Replicate 3 final exam weight | WTPXRP3 | 779-787 |
| Replicate 4 final exam weight | WTPXRP4 | 788-796 |
| Replicate 5 final exam weight | WTPXRP5 | 797-805 |
| Replicate 6 final exam weight | WTPXRP6 | 806-814 |
| Replicate 7 final exam weight | WTPXRP7 | 815-823 |
| Replicate 8 final exam weight | WTPXRP8 | 824-832 |
| Replicate 9 final exam weight | WTPXRP9 | 833-841 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|--------------------------------------|---------------|-----------|
| Replicate 10 final exam weight | WTPXRP10 | 842-850 |
| Replicate 11 final exam weight | WTPXRP11 | 851-859 |
| Replicate 12 final exam weight | WTPXRP12 | 860-868 |
| Replicate 13 final exam weight | WTPXRP13 | 869-877 |
| Replicate 14 final exam weight | WTPXRP14 | 878-886 |
| Replicate 15 final exam weight | WTPXRP15 | 887-895 |
| Replicate 16 final exam weight | WTPXRP16 | 896-904 |
| Replicate 17 final exam weight | WTPXRP17 | 905-913 |
| Replicate 18 final exam weight | WTPXRP18 | 914-922 |
| Replicate 19 final exam weight | WTPXRP19 | 923-931 |
| Replicate 20 final exam weight | WTPXRP20 | 932-940 |
| Replicate 21 final exam weight | WTPXRP21 | 941-949 |
| Replicate 22 final exam weight | WTPXRP22 | 950-958 |
| Replicate 23 final exam weight | WTPXRP23 | 959-967 |
| Replicate 24 final exam weight | WTPXRP24 | 968-976 |
| Replicate 25 final exam weight | WTPXRP25 | 977-985 |
| Replicate 26 final exam weight | WTPXRP26 | 986-994 |
| Replicate 27 final exam weight | WTPXRP27 | 995-1003 |
| Replicate 28 final exam weight | WTPXRP28 | 1004-1012 |
| Replicate 29 final exam weight | WTPXRP29 | 1013-1021 |
| Replicate 30 final exam weight | WTPXRP30 | 1022-1030 |
| Replicate 31 final exam weight | WTPXRP31 | 1031-1039 |
| Replicate 32 final exam weight | WTPXRP32 | 1040-1048 |
| Replicate 33 final exam weight | WTPXRP33 | 1049-1057 |
| Replicate 34 final exam weight | WTPXRP34 | 1058-1066 |
| Replicate 35 final exam weight | WTPXRP35 | 1067-1075 |
| Replicate 36 final exam weight | WTPXRP36 | 1076-1084 |
| Replicate 37 final exam weight | WTPXRP37 | 1085-1093 |
| Replicate 38 final exam weight | WTPXRP38 | 1094-1102 |
| Replicate 39 final exam weight | WTPXRP39 | 1103-1111 |
| Replicate 40 final exam weight | WTPXRP40 | 1112-1120 |
| Replicate 41 final exam weight | WTPXRP41 | 1121-1129 |
| Replicate 42 final exam weight | WTPXRP42 | 1130-1138 |
| Replicate 43 final exam weight | WTPXRP43 | 1139-1147 |
| Replicate 44 final exam weight | WTPXRP44 | 1148-1156 |
| Replicate 45 final exam weight | WTPXRP45 | 1157-1165 |
| Replicate 46 final exam weight | WTPXRP46 | 1166-1174 |
| Replicate 47 final exam weight | WTPXRP47 | 1175-1183 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|---------------|-----------|
| Replicate 48 final exam weight | WTPXRP48 | 1184-1192 |
| Replicate 49 final exam weight | WTPXRP49 | 1193-1201 |
| Replicate 50 final exam weight | WTPXRP50 | 1202-1210 |
| Replicate 51 final exam weight | WTPXRP51 | 1211-1219 |
| Replicate 52 final exam weight | WTPXRP52 | 1220-1228 |
| HOUSEHOLD YOUTH QUESTIONNAIRE (HYQ) | | |
| Age in months at youth interview | HYAITMO | 1229-1232 |
| MEC EXAMINATION | | |
| Language used by SP in MEC | MXPLANG | 1233 |
| Session for MEC examination | MXPSESSR | 1234 |
| Day of week of MEC exam | MXPTIDW | 1235 |
| Age in months at MEC exam | MXPAXTMR | 1236-1239 |
| HOME EXAMINATION | | |
| Day of week of home exam | HXPTIDW | 1240 |
| Age in months at home exam | HXPAXTMR | 1241-1244 |
| Session for home examination | HXPSESSR | 1245 |
| PHLEBOTOMY SCREENING QUESTIONNAIRE | | |
| Language | PHPLANG | 1246 |
| Do you have hemophilia? | PHPHEMO | 1247 |
| Recent chemo/within the past four weeks | PHPCHM2 | 1248 |
| Are you currently taking insulin? | PHPINSU | 1249 |
| Time participant last ate | PHPSNTI | 1250-1254 |
| Day participant last ate | PHPSNDA | 1255 |
| Have you had anything to drink? | PHPDRIN | 1256 |
| Time participant last drank | PHPDRTI | 1257-1261 |
| Day participant last drank | PHPDRDA | 1262 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|---------------|-----------|
| Length of calculated fast (in hours) | PHPPFAST | 1263-1267 |
| Time of venipuncture | PHPBEST | 1268-1272 |
| HEMATOLOGY | | |
| White blood cell count | WCP | 1273-1277 |
| White blood cell count: SI | WCPSI | 1278-1282 |
| Lymphocyte percent (Coulter) | LMPPCNT | 1283-1287 |
| Mononuclear percent (Coulter) | MOPPCNT | 1288-1292 |
| Granulocyte percent (Coulter) | GRPPCNT | 1293-1297 |
| Lymphocyte number (Coulter) | LMP | 1298-1302 |
| Mononuclear number (Coulter) | MOP | 1303-1306 |
| Granulocyte number (Coulter) | GRP | 1307-1311 |
| Red blood cell count | RCP | 1312-1315 |
| Red blood cell count: SI | RCPSI | 1316-1319 |
| Hemoglobin (g/dL) | HGP | 1320-1324 |
| Hemoglobin: SI (g/L) | HGPSI | 1325-1329 |
| Hematocrit (%) | HTP | 1330-1334 |
| Hematocrit: SI (L/L=1) | HTPSI | 1335-1339 |
| Mean cell volume: SI (fL) | MVPSI | 1340-1344 |
| Mean cell hemoglobin: SI (pg) | MCPSI | 1345-1349 |
| Mean cell hemoglobin concentration | MHP | 1350-1354 |
| Mean cell hemoglobin concentration: SI | MHPSI | 1355-1359 |
| Red cell distribution width (%) | RWP | 1360-1364 |
| Red cell distribution width:SI(fraction) | RWPSI | 1365-1370 |
| Platelet count | PLP | 1371-1375 |
| Platelet count: SI | PLPSI | 1376-1380 |
| Platelet distribution width (%) | DWP | 1381-1385 |
| Mean platelet volume: SI (fL) | PVPSI | 1386-1390 |
| Segment neutrophil(percent of 100 cells) | GRPDIF | 1391-1393 |
| Lymphocytes (percent of 100 cells) | LMPDIF | 1394-1396 |
| Monocytes (percent of 100 cells) | MOPDIF | 1397-1398 |
| Eosinophils (percent of 100 cells) | EOP | 1399-1400 |
| Basophils (percent of 100 cells) | BOP | 1401-1402 |
| Blasts (percent of 100 cells) | BLP | 1403 |
| Promyelocytes (percent of 100 cells) | PRP | 1404 |
| Metamyelocytes (percent of 100 cells) | MEP | 1405 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|--|---------------|-----------|
| Myelocytes (percent of 100 cells) | MLP | 1406 |
| Bands (percent of 100 cells) | BAP | 1407-1408 |
| Atyp lymphocytes (percent of 100 cells) | LAP | 1409-1410 |
| Anisocytosis (variation of cell size) | ANP | 1411 |
| Basophilic stippling | BSP | 1412 |
| Hypochromia (stain intensity of cell) | HZP | 1413 |
| Poikilocytosis (cell shape variation) | PKP | 1414 |
| Polychromatophilia (bluish color of cell) | POP | 1415 |
| Macrocytosis (large cell prevalence) | MRP | 1416 |
| Microcytosis (small cell prevalence) | MIP | 1417 |
| Sickle cells | SIP | 1418 |
| Spherocytosis | SHP | 1419 |
| Target cells | TTP | 1420 |
| Toxic granulation | TXP | 1421 |
| Vacuolated cells | VUP | 1422 |
| GENERAL BIOCHEMISTRY TESTS | | |
| Lead (ug/dL) | PBP | 1423-1426 |
| Lead: SI (umol/L) | PBPSI | 1427-1431 |
| Erythrocyte protoporphyrin (ug/dL) | EPP | 1432-1435 |
| Erythrocyte protoporphyrin: SI (umol/L) | EPPSI | 1436-1440 |
| Serum iron (ug/dL) | FEP | 1441-1443 |
| Serum iron: SI (umol/L) | FEPSI | 1444-1448 |
| Serum TIBC (ug/dL) | TIP | 1449-1452 |
| Serum TIBC: SI (umol/L) | TIPSI | 1453-1458 |
| Serum transferrin saturation (%) | PXP | 1459-1462 |
| Serum ferritin (ng/mL) | FRP | 1463-1466 |
| Serum ferritin: SI (ug/L) | FRPSI | 1467-1470 |
| Serum folate (ng/mL) | FOP | 1471-1475 |
| Serum folate: SI (nmol/L) | FOPSI | 1476-1480 |
| RBC folate (ng/mL) | RBP | 1481-1484 |
| RBC folate: SI (nmol/L) | RBPSI | 1485-1490 |
| Serum vitamin B12 (pg/mL) | VBP | 1491-1496 |
| Serum vitamin B12: SI (pmol/L) | VBPSI | 1497-1504 |
| Serum vitamin C (mg/dL) | VCP | 1505-1508 |
| Serum vitamin C: SI (mmol/L) | VCPSI | 1509-1514 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|------------------|-----------|
| Serum normalized calcium: SI (mmol/L) | ICPSI | 1515-1518 |
| Serum total calcium: SI (mmol/L) | CAPSI | 1519-1522 |
| Serum selenium (ng/mL) | SEP | 1523-1526 |
| Serum selenium: SI (nmol/L) | SEPSI | 1527-1530 |
| Serum vitamin A (ug/dL) | VAP | 1531-1533 |
| Serum vitamin A: SI (umol/L) | VAPSI | 1534-1537 |
| Serum vitamin E (ug/dL) | VEP | 1538-1542 |
| Serum vitamin E: SI (umol/L) | VEPSI | 1543-1548 |
| Serum alpha carotene (ug/dL) | ACP | 1549-1551 |
| Serum alpha carotene: SI (umol/L) | ACPSI | 1552-1555 |
| Serum beta carotene (ug/dL) | BCP | 1556-1559 |
| Serum beta carotene: SI (umol/L) | BCPSI | 1560-1564 |
| Serum beta cryptoxanthin (ug/dL) | BXP | 1565-1567 |
| Serum beta cryptoxanthin: SI (umol/L) | BXPSI | 1568-1571 |
| Serum lutein/zeaxanthin (ug/dL) | LUP | 1572-1574 |
| Serum lutein/zeaxanthin: SI (umol/L) | LUPSI | 1575-1578 |
| Serum lycopene (ug/dL) | LYP | 1579-1581 |
| Serum lycopene: SI (umol/L) | LYPSI | 1582-1585 |
| Serum sum retinyl esters (ug/dL) | REP | 1586-1588 |
| Serum sum retinyl esters: SI (umol/L) | REPSI | 1589-1592 |
| Serum cotinine (ng/mL) | COP | 1593-1597 |
| Serum cholesterol (mg/dL) | TCP | 1598-1600 |
| Serum cholesterol: SI (mmol/L) | TCPSI | 1601-1605 |
| Serum triglycerides (mg/dL) | TGP | 1606-1609 |
| Serum triglycerides: SI (mmol/L) | TGPSI | 1610-1614 |
| Serum LDL cholesterol (mg/dL) | LCP | 1615-1617 |
| Serum LDL cholesterol: SI (mmol/L) | LCPSI | 1618-1621 |
| Serum HDL cholesterol (mg/dL) | HDP | 1622-1624 |
| Serum HDL cholesterol: SI (mmol/L) | HDPSI | 1625-1628 |
| Serum apolipoprotein AI (mg/dL) | AAP | 1629-1631 |
| Serum apolipoprotein AI: SI (g/L) | AAPSI | 1632-1635 |
| Serum apolipoprotein B (mg/dL) | ABP | 1636-1638 |
| Serum apolipoprotein B: SI (g/L) | ABPSI | 1639-1642 |
| Serum lipoprotein(a) (mg/dL) | LPP | 1643-1645 |
| Serum lipoprotein(a): SI (g/L) | LPPSI | 1646-1649 |
| Serum FSH: SI (IU/L) | FHPSI | 1650-1654 |
| Serum luteinizing hormone: SI (IU/L) | LHPSI | 1655-1658 |
| Plasma fibrinogen (mg/dL) | FBP | 1659-1662 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|--|---------------|-----------|
| Plasma fibrinogen: SI (g/L) | FBPSI | 1663-1666 |
| Serum C-reactive protein (mg/dL) | CRP | 1667-1671 |
| ANTIBODY TESTS | | |
| Serum tetanus antibody (U/mL) | TEP | 1672-1677 |
| Serum hepatitis A antibody | AHP | 1678 |
| Serum hepatitis B core antibody | HBP | 1679 |
| Serum hepatitis B surface antibody | SSP | 1680-1681 |
| Serum hepatitis B surface antigen | SAP | 1682 |
| Serum hepatitis C antibody | HCP | 1683 |
| Serum hepatitis D antibody | DHP | 1684 |
| Serum herpes I antibody | H1P | 1685 |
| Serum herpes II antibody | H2P | 1686 |
| Serum rubella antibody | RUP | 1687-1691 |
| Serum rubella antibody (IU) | RUPUNIT | 1692-1695 |
| Serum varicella antibody | VRP | 1696-1700 |
| Serum toxoplasmosis antibody | TOP | 1701-1703 |
| Serum rheumatoid factor antibody | RFP | 1704-1708 |
| Serum latex antibody (IU/mL) | L1P | 1709-1713 |
| Serum helicobacter pylori antibody | HPP | 1714 |
| BIOCHEMISTRY PROFILE | | |
| Serum sodium: SI (mmol/L) | NAPSI | 1715-1719 |
| Serum potassium: SI (mmol/L) | SKPSI | 1720-1723 |
| Serum chloride: SI (mmol/L) | CLPSI | 1724-1728 |
| Serum bicarbonate: SI (mmol/L) | C3PSI | 1729-1730 |
| Serum total calcium (mg/dL) | SCP | 1731-1734 |
| Serum total calcium: SI (mmol/L) | SCPSI | 1735-1739 |
| Serum phosphorus (mg/dL) | PSP | 1740-1743 |
| Serum phosphorus: SI (mmol/L) | PSPSI | 1744-1748 |
| Serum uric acid (mg/dL) | UAP | 1749-1752 |
| Serum uric acid: SI (umol/L) | UAPSI | 1753-1757 |
| Serum glucose (mg/dL) | SGP | 1758-1760 |
| Serum glucose: SI (mmol/L) | SGPSI | 1761-1765 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|--|---------------|-----------|
| Serum blood urea nitrogen (mg/dL) | BUP | 1766-1768 |
| Serum blood urea nitrogen: SI (mmol/L) | BUPSI | 1769-1773 |
| Serum total bilirubin (mg/dL) | TBP | 1774-1777 |
| Serum total bilirubin: SI (umol/L) | TBPSI | 1778-1783 |
| Serum creatinine (mg/dL) | CEP | 1784-1787 |
| Serum creatinine: SI (umol/L) | CEPSI | 1788-1793 |
| Serum iron (ug/dL) | SFP | 1794-1796 |
| Serum iron: SI (umol/L) | SFPSI | 1797-1800 |
| Serum cholesterol (mg/dL) | CHP | 1801-1804 |
| Serum cholesterol: SI (mmol/L) | CHPSI | 1805-1810 |
| Serum triglycerides (mg/dL) | TRP | 1811-1814 |
| Serum triglycerides: SI (mmol/L) | TRPSI | 1815-1820 |
| Aspartate aminotransferase: SI(U/L) | ASPSI | 1821-1823 |
| Alanine aminotransferase: SI (U/L) | ATPSI | 1824-1826 |
| Gamma glutamyl transferase: SI(U/L) | GGPSI | 1827-1830 |
| Serum lactate dehydrogenase: SI (U/L) | LDPSI | 1831-1834 |
| Serum alkaline phosphatase: SI (U/L) | APPSI | 1835-1838 |
| Serum total protein (g/dL) | TPP | 1839-1842 |
| Serum total protein: SI (g/L) | TPPSI | 1843-1845 |
| Serum albumin (g/dL) | AMP | 1846-1848 |
| Serum albumin: SI (g/L) | AMPSI | 1849-1851 |
| Serum globulin (g/dL) | GBP | 1852-1854 |
| Serum globulin: SI (g/L) | GBPSI | 1855-1857 |
| Serum osmolality: SI (mmol/Kg) | OSPSI | 1858-1860 |
| DIABETES TESTING PROFILE | | |
| Glycated hemoglobin: (%) | GHP | 1861-1864 |
| Glycated hemoglobin: test method | GHPMETH | 1865 |
| Plasma glucose (mg/dL) | G1P | 1866-1870 |
| Plasma glucose: SI (mmol/L) | G1PSI | 1871-1876 |
| Incomplete glucose test (OGTT) code | G1PCODE | 1877-1878 |
| Minutes between drink and second draw | G1PTIM1 | 1879-1881 |
| Minutes between first and second draw | G1PTIM2 | 1882-1884 |
| Second plasma glucose (mg/dL) | G2P | 1885-1889 |
| Second plasma glucose: SI (mmol/L) | G2PSI | 1890-1895 |
| Serum C-peptide (pmol/mL) | C1P | 1896-1900 |

NHANES III Laboratory Data File Index
Whole Blood, Serum, Plasma, and Urine Data

| Description | Variable Name | Positions |
|---|---------------|-----------|
| Serum C-peptide: SI (nmol/L) | C1PSI | 1901-1905 |
| Second serum C-peptide (pmol/mL) | C2P | 1906-1911 |
| Second serum C-peptide: SI (nmol/L) | C2PSI | 1912-1917 |
| Serum insulin (uU/mL) | I1P | 1918-1923 |
| Serum insulin: SI (pmol/L) | I1PSI | 1924-1930 |
| Serum insulin: test kit | I1P2PFLG | 1931 |
| Second serum insulin (uU/mL) | I2P | 1932-1937 |
| Second serum insulin: SI (pmol/L) | I2PSI | 1938-1944 |
| URINE TESTS | | |
| Urinary cadmium (ng/mL) | UDP | 1945-1949 |
| Urinary cadmium: SI (nmol/L) | UDPSI | 1950-1955 |
| Urinary creatinine (mg/dL) | URP | 1956-1960 |
| Urinary creatinine: SI (mmol/L) | URPSI | 1961-1964 |
| Urinary albumin (ug/mL) | UBP | 1965-1970 |
| Urinary iodine (ug/dL) | UIP | 1971-1977 |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

HOUSEHOLD SCREENER QUESTIONNAIRE (HSQ)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|------------------------------------|----------|
| ----- | | | |
| 12 | | Race-ethnicity | See note |
| DMARETHN | 10507 | 1 Non-Hispanic white | |
| | 8756 | 2 Non-Hispanic black | |
| | 8786 | 3 Mexican-American | |
| | 1265 | 4 Other | |
| 13 | | Race | See note |
| DMARACER | 19180 | 1 White | |
| | 9091 | 2 Black | |
| | 1037 | 3 Other | |
| | 6 | 8 Mexican-American of unknown race | |
| 14 | | Ethnicity | See note |
| DMAETHNR | 8786 | 1 Mexican-American | |
| | 788 | 2 Other Hispanic | |
| | 19740 | 3 Not Hispanic | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

HOUSEHOLD SCREENER QUESTIONNAIRE (HSQ)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|-----------------------------------|----------|
| | 15 | Sex | |
| HSSEX | 13980 | 1 Male | |
| | 15334 | 2 Female | |
| | 16-17 | Age at interview (Screeners) | See note |
| HSAGEIR | 29165 | 01-89 | |
| | 149 | 90 90+ | |
| | 18 | Age at interview-unit (Screeners) | |
| HSAGEU | 29314 | 2 Years | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

HOUSEHOLD SCREENER QUESTIONNAIRE (HSQ)

| Positions | | Item description | |
|-----------|--------|--------------------------|----------|
| SAS name | Counts | and code | Notes |
| | | | |
| 19-22 | | Age in months (Screener) | See note |
| HSAITMOR | 29157 | 0012-1079 | |
| | 147 | 1080 1080+ months | |
| | 10 | 9999 Don't know | |
| 23-24 | | Family size | See note |
| HSFSIZER | 3076 | 01 | |
| | 5411 | 02 | |
| | 5006 | 03 | |
| | 5950 | 04 | |
| | 4313 | 05 | |
| | 2312 | 06 | |
| | 1236 | 07 | |
| | 821 | 08 | |
| | 428 | 09 | |
| | 761 | 10 10+ | |
| 25-26 | | Household size | See note |
| HSHSIZER | 2478 | 01 | |
| | 5473 | 02 | |
| | 5040 | 03 | |
| | 6041 | 04 | |
| | 4337 | 05 | |
| | 2393 | 06 | |
| | 1301 | 07 | |
| | 893 | 08 | |
| | 459 | 09 | |
| | 899 | 10 10+ | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

HOUSEHOLD SCREENER QUESTIONNAIRE (HSQ)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|----------|
| 27-29 DMPCNTYR | | County FIPS codes for United States counties with populations >= 500,000 | See note |
| | 13799 | 001-439 | |
| | 15515 | Blank | |
| 30-31 DMPFIPSR | | State FIPS codes for United States counties with populations >= 500,000 | See note |
| | 359 | 04 | |
| | 4531 | 06 | |
| | 1090 | 12 | |
| | 900 | 17 | |
| | 242 | 25 | |
| | 676 | 26 | |
| | 312 | 29 | |
| | 1662 | 36 | |
| | 625 | 39 | |
| | 724 | 42 | |
| | 276 | 44 | |
| | 2044 | 48 | |
| | 358 | 53 | |
| | 15515 | Blank | |
| 32 DMPMETRO | | Urbanization classification based on USDA Rural/Urban continuum codes. | See note |
| | 14615 | 1 Central counties of metro areas of 1 million population or more, OR, Fringe counties of metro areas of 1 million population or more | |
| | 14699 | 2 All other areas | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

HOUSEHOLD SCREENER QUESTIONNAIRE (HSQ)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|------------------------------|----------|
| | 33 | Census region | See note |
| DMPCREGN | 3740 | 1 Northeast | |
| | 5498 | 2 Midwest | |
| | 12639 | 3 South | |
| | 7437 | 4 West | |
| | 34-39 | Poverty Income Ratio | See note |
| DMPPIR | 82 | 00.000 No reported income | |
| | 26503 | 000.02-11.889 | |
| | 2729 | 888888 Blank but applicable | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

SURVEY DESIGN DATA

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------------------|--|----------|
| 40 SDPPHASE | 14833 14481 | Phase of NHANES III survey 1 1988-1991 2 1991-1994 | See note |
| 41 SDPPSU6 | 14630 14684 | Total NHANES III Pseudo-PSU 1 2 | See note |
| 42-43 SDPSTRA6 | 29314 | Total NHANES III Pseudo-stratum 01-49 | See note |
| 44 SDPPSU1 | 7633 7200 14481 | Phase 1 Pseudo-PSU 1 2 Blank | See note |
| 45-46 SDPSTRA1 | 14833 14481 | Phase 1 Pseudo-stratum 01-23 Blank | See note |
| 47 SDPPSU2 | 7080 7401 14833 | Phase 2 Pseudo-PSU 1 2 Blank | See note |
| 48-49 SDPSTRA2 | 14481 14833 | Phase 2 Pseudo-stratum 01-23 Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

SAMPLING WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|----------------------|---|----------|
| 50-58 WTPFQX6 | 29314 | Total NHANES III interviewed sample final weight 000215.53-0132278.9 | See note |
| 59-67 WTPFEX6 | 457 28857 | Total NHANES III MEC-examined sample final weight 000000.00 000213.45-140778.72 | See note |
| 68-76 WTPFHX6 | 29314 | Total NHANES III MEC and home- examined final weight 000214.25-139744.91 | See note |
| 77-85 WTPFALG6 | 23 12106 17185 | Total NHANES III allergy subsample final weight 000000.00 000213.45-288897.91 Blank | See note |
| 86-94 WTPFCNS6 | 12 5662 23640 | Total NHANES III central nervous system (CNS) subsample final weight 000000.00 001316.46-295826.48 Blank | See note |
| 95-103 WTPFSD6 | 920 9127 19267 | Total NHANES III morning session MEC-examined subsample final weight 000000.00 000450.95-292590.96 Blank | See note |
| 104-112 WTPFMD6 | 697 9497 19120 | Total NHANES III afternoon/evening session MEC-examined subsample final weight 000000.00 000495.13-256201.99 Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

SAMPLING WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|----------|
| 113-121 WTPFHSD6 | | Total NHANES III morning session MEC and home-examined subsample final weight | See note |
| | 791 | 000000.00 | |
| | 9254 | 000446.49-291479.91 | |
| | 19269 | Blank | |
| 122-130 WTPFHMD6 | | Total NHANES III afternoon/evening session MEC and home-examined subsample final weight | See note |
| | 562 | 000000.00 | |
| | 9630 | 000503.56-256245.36 | |
| | 19122 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

SAMPLING WEIGHTS - NHANES III PHASE 1 (1988-91)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|----------|
| 131-139 WTPFQX1 | | Phase 1 interviewed sample final weight | See note |
| | 14833 | 000461.29-264557.81 | |
| | 14481 | Blank | |
| 140-148 WTPFEX1 | | Phase 1 MEC-examined sample final weight | See note |
| | 229 | 000000.00 | |
| | 14604 | 000527.01-281557.44 | |
| | 14481 | Blank | |
| 149-157 WTPFHX1 | | Phase 1 MEC and home-examined sample final weight | See note |
| | 14833 | 000513.14-279489.83 | |
| | 14481 | Blank | |
| 158-166 WTPFALG1 | | Phase 1 allergy subsample final weight | See note |
| | 14 | 000000.00 | |
| | 6097 | 000821.62-577795.82 | |
| | 23203 | Blank | |
| 167-175 WTPFCNS1 | | Phase 1 central nervous system (CNS) subsample final weight | See note |
| | 8 | 000000.00 | |
| | 2751 | 002699.84-591652.96 | |
| | 26555 | Blank | |
| 176-184 WTPFSD1 | | Phase 1 morning session MEC-examined subsample final weight | See note |
| | 451 | 000000.00 | |
| | 4462 | 001111.36-585181.93 | |
| | 24401 | Blank | |
| 185-193 WTPFMD1 | | Phase 1 afternoon/evening session MEC- examined subsample final weight | See note |
| | 322 | 000000.00 | |
| | 4726 | 001104.11-506697.07 | |
| | 24266 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

SAMPLING WEIGHTS - NHANES III PHASE 1 (1988-91)

| Positions | | Item description | |
|-----------|--------|--|----------|
| SAS name | Counts | and code | Notes |
| 194-202 | | Phase 1 morning session MEC and home- | See note |
| WTPFHSD1 | | examined subsample final weight | |
| | 373 | 000000.00 | |
| | 4540 | 0001091.8-582959.83 | |
| | 24401 | Blank | |
| 203-211 | | Phase 1 afternoon/evening session MEC | See note |
| WTPFHMD1 | | and home-examined subsample final weight | |
| | 264 | 000000.00 | |
| | 4784 | 001085.73-507417.05 | |
| | 24266 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

SAMPLING WEIGHTS - NHANES III PHASE 2 (1991-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|----------|
| 212-220 WTPFQX2 | | Phase 2 interviewed sample final weight | See note |
| | 14481 | 000431.06-243267.38 | |
| | 14833 | Blank | |
| 221-229 WTPFEX2 | | Phase 2 MEC-examined sample final weight | See note |
| | 228 | 000000.00 | |
| | 14253 | 000426.91-262887.56 | |
| | 14833 | Blank | |
| 230-238 WTPFHX2 | | Phase 2 MEC and home-examined sample final weight | See note |
| | 14481 | 0000428.5-262188.52 | |
| | 14833 | Blank | |
| 239-247 WTPFALG2 | | Phase 2 allergy subsample final weight | See note |
| | 9 | 000000.00 | |
| | 6009 | 000426.91-552445.57 | |
| | 23296 | Blank | |
| 248-256 WTPFCNS2 | | Phase 2 central nervous system (CNS) subsample final weight | See note |
| | 4 | 000000.00 | |
| | 2911 | 002632.92-518040.33 | |
| | 26399 | Blank | |
| 257-265 WTPFSD2 | | Phase 2 morning session MEC-examined subsample final weight | See note |
| | 469 | 000000.00 | |
| | 4665 | 0000901.9-550430.69 | |
| | 24180 | Blank | |
| 266-274 WTPFMD2 | | Phase 2 afternoon/evening session MEC- examined subsample final weight | See note |
| | 375 | 000000.00 | |
| | 4771 | 000990.26-512403.98 | |
| | 24168 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

SAMPLING WEIGHTS - NHANES III PHASE 2 (1991-94)

| Positions | | Item description | |
|-----------|--------|--|----------|
| SAS name | Counts | and code | Notes |
| 275-283 | | Phase 2 morning session MEC and home- | See note |
| WTPFHSD2 | | examined subsample final weight | |
| | 418 | 000000.00 | |
| | 4714 | 000892.98-552545.64 | |
| | 24182 | Blank | |
| 284-292 | | Phase 2 afternoon/evening session MEC | See note |
| WTPFHMD2 | | and home-examined subsample final weight | |
| | 298 | 000000.00 | |
| | 4846 | 001007.13-512490.71 | |
| | 24170 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE INTERVIEW WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 293-301 WTPQRP1 | 29314 | Replicate 1 final interview weight 000053.27-148435.02 | See note |
| 302-310 WTPQRP2 | 29314 | Replicate 2 final interview weight 000067.13-143746.82 | See note |
| 311-319 WTPQRP3 | 29314 | Replicate 3 final interview weight 000047.49-152075.62 | See note |
| 320-328 WTPQRP4 | 29314 | Replicate 4 final interview weight 000062.62-137241.93 | See note |
| 329-337 WTPQRP5 | 29314 | Replicate 5 final interview weight 000048.42-147700.94 | See note |
| 338-346 WTPQRP6 | 29314 | Replicate 6 final interview weight 0000053.1-146803.63 | See note |
| 347-355 WTPQRP7 | 29314 | Replicate 7 final interview weight 000058.18-145261.07 | See note |
| 356-364 WTPQRP8 | 29314 | Replicate 8 final interview weight 000048.23-161126.44 | See note |
| 365-373 WTPQRP9 | 29314 | Replicate 9 final interview weight 000053.27-147301.59 | See note |
| 374-382 WTPQRP10 | 29314 | Replicate 10 final interview weight 000073.37-0148125.5 | See note |
| 383-391 WTPQRP11 | 29314 | Replicate 11 final interview weight 000058.31-146940.58 | See note |
| 392-400 WTPQRP12 | 29314 | Replicate 12 final interview weight 000053.67-153958.72 | See note |
| 401-409 WTPQRP13 | 29314 | Replicate 13 final interview weight 000067.93-147395.78 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE INTERVIEW WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 410-418 WTPQRP14 | 29314 | Replicate 14 final interview weight 000065.08-138456.05 | See note |
| 419-427 WTPQRP15 | 29314 | Replicate 15 final interview weight 000062.35-140673.55 | See note |
| 428-436 WTPQRP16 | 29314 | Replicate 16 final interview weight 000040.28-147603.74 | See note |
| 437-445 WTPQRP17 | 29314 | Replicate 17 final interview weight 000045.36-154057.83 | See note |
| 446-454 WTPQRP18 | 29314 | Replicate 18 final interview weight 000070.42-138896.98 | See note |
| 455-463 WTPQRP19 | 29314 | Replicate 19 final interview weight 000050.96-139447.18 | See note |
| 464-472 WTPQRP20 | 29314 | Replicate 20 final interview weight 000045.79-156365.73 | See note |
| 473-481 WTPQRP21 | 29314 | Replicate 21 final interview weight 000049.79-146241.31 | See note |
| 482-490 WTPQRP22 | 29314 | Replicate 22 final interview weight 000047.25-0154848.6 | See note |
| 491-499 WTPQRP23 | 29314 | Replicate 23 final interview weight 000037.18-148309.04 | See note |
| 500-508 WTPQRP24 | 29314 | Replicate 24 final interview weight 000057.42-141344.14 | See note |
| 509-517 WTPQRP25 | 29314 | Replicate 25 final interview weight 000044.13-145105.09 | See note |
| 518-526 WTPQRP26 | 29314 | Replicate 26 final interview weight 0000066.1-146773.53 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE INTERVIEW WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 527-535 WTPQRP27 | 29314 | Replicate 27 final interview weight 000044.88-142455.25 | See note |
| 536-544 WTPQRP28 | 29314 | Replicate 28 final interview weight 000000046-148272.41 | See note |
| 545-553 WTPQRP29 | 29314 | Replicate 29 final interview weight 000079.38-153624.57 | See note |
| 554-562 WTPQRP30 | 29314 | Replicate 30 final interview weight 000058.09-151140.25 | See note |
| 563-571 WTPQRP31 | 29314 | Replicate 31 final interview weight 000051.39-159963.39 | See note |
| 572-580 WTPQRP32 | 29314 | Replicate 32 final interview weight 000066.17-132356.37 | See note |
| 581-589 WTPQRP33 | 29314 | Replicate 33 final interview weight 0000057.8-136762.37 | See note |
| 590-598 WTPQRP34 | 29314 | Replicate 34 final interview weight 000062.28-140628.16 | See note |
| 599-607 WTPQRP35 | 29314 | Replicate 35 final interview weight 000063.73-154630.49 | See note |
| 608-616 WTPQRP36 | 29314 | Replicate 36 final interview weight 000067.29-153648.69 | See note |
| 617-625 WTPQRP37 | 29314 | Replicate 37 final interview weight 000043.47-135065.98 | See note |
| 626-634 WTPQRP38 | 29314 | Replicate 38 final interview weight 000054.55-152122.87 | See note |
| 635-643 WTPQRP39 | 29314 | Replicate 39 final interview weight 000050.55-152941.69 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE INTERVIEW WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 644-652 WTPQRP40 | 29314 | Replicate 40 final interview weight 000054.45-146815.92 | See note |
| 653-661 WTPQRP41 | 29314 | Replicate 41 final interview weight 000059.62-141514.78 | See note |
| 662-670 WTPQRP42 | 29314 | Replicate 42 final interview weight 000068.97-0140162.4 | See note |
| 671-679 WTPQRP43 | 29314 | Replicate 43 final interview weight 000044.04-150981.83 | See note |
| 680-688 WTPQRP44 | 29314 | Replicate 44 final interview weight 000040.36-144080.03 | See note |
| 689-697 WTPQRP45 | 29314 | Replicate 45 final interview weight 000054.74-0142465.6 | See note |
| 698-706 WTPQRP46 | 29314 | Replicate 46 final interview weight 000078.43-137838.21 | See note |
| 707-715 WTPQRP47 | 29314 | Replicate 47 final interview weight 000052.71-145055.34 | See note |
| 716-724 WTPQRP48 | 29314 | Replicate 48 final interview weight 000046.91-148787.77 | See note |
| 725-733 WTPQRP49 | 29314 | Replicate 49 final interview weight 0000072.4-148375.43 | See note |
| 734-742 WTPQRP50 | 29314 | Replicate 50 final interview weight 000070.53-159394.39 | See note |
| 743-751 WTPQRP51 | 29314 | Replicate 51 final interview weight 000054.73-0144964.3 | See note |
| 752-760 WTPQRP52 | 29314 | Replicate 52 final interview weight 000072.04-149087.24 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE EXAMINATION WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------------|--|----------|
| 761-769 WTPXRP1 | 457 28857 | Replicate 1 final exam weight 000000.00 000054.73-164698.81 | See note |
| 770-778 WTPXRP2 | 457 28857 | Replicate 2 final exam weight 000000.00 0000067.3-164887.24 | See note |
| 779-787 WTPXRP3 | 457 28857 | Replicate 3 final exam weight 000000.00 0000048.2-0161201.8 | See note |
| 788-796 WTPXRP4 | 457 28857 | Replicate 4 final exam weight 000000.00 000067.24-149561.18 | See note |
| 797-805 WTPXRP5 | 457 28857 | Replicate 5 final exam weight 000000.00 000055.97-146312.81 | See note |
| 806-814 WTPXRP6 | 457 28857 | Replicate 6 final exam weight 000000.00 000051.48-156250.53 | See note |
| 815-823 WTPXRP7 | 457 28857 | Replicate 7 final exam weight 000000.00 000060.06-0157694.3 | See note |
| 824-832 WTPXRP8 | 457 28857 | Replicate 8 final exam weight 000000.00 0000053.1-169111.97 | See note |
| 833-841 WTPXRP9 | 457 28857 | Replicate 9 final exam weight 000000.00 000052.31-156939.22 | See note |
| 842-850 WTPXRP10 | 457 28857 | Replicate 10 final exam weight 000000.00 000072.13-0165805.2 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE EXAMINATION WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------------|--|----------|
| 851-859 WTPXRP11 | 457 28857 | Replicate 11 final exam weight 000000.00 000053.54-154918.93 | See note |
| 860-868 WTPXRP12 | 457 28857 | Replicate 12 final exam weight 000000.00 000055.35-164023.88 | See note |
| 869-877 WTPXRP13 | 457 28857 | Replicate 13 final exam weight 000000.00 0000067.9-147355.32 | See note |
| 878-886 WTPXRP14 | 457 28857 | Replicate 14 final exam weight 000000.00 000067.04-154034.72 | See note |
| 887-895 WTPXRP15 | 457 28857 | Replicate 15 final exam weight 000000.00 000062.21-156384.73 | See note |
| 896-904 WTPXRP16 | 457 28857 | Replicate 16 final exam weight 000000.00 000000040-157994.12 | See note |
| 905-913 WTPXRP17 | 457 28857 | Replicate 17 final exam weight 000000.00 000048.34-160889.46 | See note |
| 914-922 WTPXRP18 | 457 28857 | Replicate 18 final exam weight 000000.00 0000075.2-153937.93 | See note |
| 923-931 WTPXRP19 | 457 28857 | Replicate 19 final exam weight 000000.00 000056.83-149483.14 | See note |
| 932-940 WTPXRP20 | 457 28857 | Replicate 20 final exam weight 000000.00 0000045.1-165457.71 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE EXAMINATION WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------------|--|----------|
| ----- | | | |
| 941-949 WTPXRP21 | 457 28857 | Replicate 21 final exam weight 000000.00 000055.15-152305.97 | See note |
| 950-958 WTPXRP22 | 457 28857 | Replicate 22 final exam weight 000000.00 000045.53-159746.13 | See note |
| 959-967 WTPXRP23 | 457 28857 | Replicate 23 final exam weight 000000.00 000037.51-158016.62 | See note |
| 968-976 WTPXRP24 | 457 28857 | Replicate 24 final exam weight 000000.00 000054.91-153043.54 | See note |
| 977-985 WTPXRP25 | 457 28857 | Replicate 25 final exam weight 000000.00 000043.77-155179.51 | See note |
| 986-994 WTPXRP26 | 457 28857 | Replicate 26 final exam weight 000000.00 000071.23-168273.22 | See note |
| 995-1003 WTPXRP27 | 457 28857 | Replicate 27 final exam weight 000000.00 000043.82-153212.25 | See note |
| 1004-1012 WTPXRP28 | 457 28857 | Replicate 28 final exam weight 000000.00 000045.61-147920.01 | See note |
| 1013-1021 WTPXRP29 | 457 28857 | Replicate 29 final exam weight 000000.00 000083.17-159279.49 | See note |
| 1022-1030 WTPXRP30 | 457 28857 | Replicate 30 final exam weight 000000.00 000059.05-162389.35 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE EXAMINATION WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------------|--|----------|
| ----- | | | |
| 1031-1039 WTPXRP31 | 457 28857 | Replicate 31 final exam weight 000000.00 000052.61-163894.16 | See note |
| 1040-1048 WTPXRP32 | 457 28857 | Replicate 32 final exam weight 000000.00 000067.05-0149876.8 | See note |
| 1049-1057 WTPXRP33 | 457 28857 | Replicate 33 final exam weight 000000.00 000055.58-153417.47 | See note |
| 1058-1066 WTPXRP34 | 457 28857 | Replicate 34 final exam weight 000000.00 000063.45-156981.83 | See note |
| 1067-1075 WTPXRP35 | 457 28857 | Replicate 35 final exam weight 000000.00 000064.47-157897.09 | See note |
| 1076-1084 WTPXRP36 | 457 28857 | Replicate 36 final exam weight 000000.00 000067.68-171875.06 | See note |
| 1085-1093 WTPXRP37 | 457 28857 | Replicate 37 final exam weight 000000.00 000045.36-153137.39 | See note |
| 1094-1102 WTPXRP38 | 457 28857 | Replicate 38 final exam weight 000000.00 000055.94-159979.02 | See note |
| 1103-1111 WTPXRP39 | 457 28857 | Replicate 39 final exam weight 000000.00 000057.47-151920.72 | See note |
| 1112-1120 WTPXRP40 | 457 28857 | Replicate 40 final exam weight 000000.00 000057.86-157191.41 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE EXAMINATION WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------------|--|----------|
| 1121-1129 WTPXRP41 | 457 28857 | Replicate 41 final exam weight 000000.00 0000061.4-000146023 | See note |
| 1130-1138 WTPXRP42 | 457 28857 | Replicate 42 final exam weight 000000.00 000069.57-154624.02 | See note |
| 1139-1147 WTPXRP43 | 457 28857 | Replicate 43 final exam weight 000000.00 000044.35-159439.04 | See note |
| 1148-1156 WTPXRP44 | 457 28857 | Replicate 44 final exam weight 000000.00 000044.16-155951.73 | See note |
| 1157-1165 WTPXRP45 | 457 28857 | Replicate 45 final exam weight 000000.00 000059.87-147941.67 | See note |
| 1166-1174 WTPXRP46 | 457 28857 | Replicate 46 final exam weight 000000.00 000074.92-150980.02 | See note |
| 1175-1183 WTPXRP47 | 457 28857 | Replicate 47 final exam weight 000000.00 000050.64-151763.92 | See note |
| 1184-1192 WTPXRP48 | 457 28857 | Replicate 48 final exam weight 000000.00 0000045.8-156115.62 | See note |
| 1193-1201 WTPXRP49 | 457 28857 | Replicate 49 final exam weight 000000.00 000082.17-159609.54 | See note |
| 1202-1210 WTPXRP50 | 457 28857 | Replicate 50 final exam weight 000000.00 000071.97-168153.71 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

FAY'S BRR REPLICATE EXAMINATION WEIGHTS - TOTAL NHANES III (1988-94)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------------|--|----------|
| 1211-1219 WTPXRP51 | 457 28857 | Replicate 51 final exam weight 000000.00 000054.04-158632.23 | See note |
| 1220-1228 WTPXRP52 | 457 28857 | Replicate 52 final exam weight 000000.00 000073.26-158493.21 | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

HOUSEHOLD YOUTH QUESTIONNAIRE (HYQ)

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|----------------------|--|----------|
| 1229-1232 HYAITMO | 11138 14 18162 | Age in months at household youth interview 0012-0204 8888 Blank but applicable Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

MEC EXAMINATION

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---------------------------------------|----------|
| | 1233 | Language used by sample person in MEC | See note |
| MXPLANG | 23936 | 1 English | |
| | 3906 | 2 Spanish | |
| | 3 | 3 Other | |
| | 1469 | Blank | |
| | 1234 | Examination session for MEC | See note |
| MXPSESSR | | examinees | |
| | 13643 | 1 Morning | |
| | 9419 | 2 Afternoon | |
| | 5795 | 3 Evening | |
| | 457 | Blank | |
| | 1235 | Day of week of MEC exam | |
| MXPTIDW | 2884 | 1 Sunday | |
| | 2618 | 2 Monday | |
| | 2503 | 3 Tuesday | |
| | 2914 | 4 Wednesday | |
| | 5466 | 5 Thursday | |
| | 5082 | 6 Friday | |
| | 7390 | 7 Saturday | |
| | 457 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

MEC EXAMINATION

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|------------------------------|----------|
| 1236-1239 | | Age in months at MEC exam | See note |
| MXPAXTMR | 28751 | 0012-1079 | |
| | 106 | 1080 1080+ months | |
| | 457 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DEMOGRAPHIC DATA

HOME EXAMINATION

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 1240 | | Day of week of home exam | |
| HXPTIDW | 22 | 1 Sunday | |
| | 111 | 2 Monday | |
| | 6 | 3 Tuesday | |
| | 16 | 4 Wednesday | |
| | 123 | 5 Thursday | |
| | 119 | 6 Friday | |
| | 60 | 7 Saturday | |
| | 28857 | Blank | |
| 1241-1244 | | Age in months at home exam | See note |
| HXPAXTMR | 410 | 0252-1079 | |
| | 47 | 1080 1080+ months | |
| | 28857 | Blank | |
| 1245 | | Examination session for home examinees | See note |
| HXPSESSR | 203 | 1 Morning | |
| | 212 | 2 Afternoon | |
| | 38 | 3 Evening | |
| | 4 | 8 Blank but applicable | |
| | 28857 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

PHLEBOTOMY SCREENING QUESTIONNAIRE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------|---|----------|
| | 1246 | Language | See note |
| PHPLANG | 25009 | 1 English | |
| | 2736 | 2 Spanish | |
| | 1569 | 8 Blank but applicable | |
| | 1247 | Do you have hemophilia? This is a hereditary blood-clotting disorder | See note |
| PHPHEMO | 9 | 1 Yes, subsequent fields blank | |
| | 27736 | 2 No | |
| | 1569 | 8 Blank but applicable | |
| | 1248 | Within the past four weeks have you received any cancer chemotherapy treatment? | See note |
| PHPCHM2 | 19 | 1 Yes, subsequent fields blank | |
| | 27717 | 2 No | |
| | 1569 | 8 Blank but applicable | |
| | 9 | Blank | |
| | 1249 | Are you currently taking insulin? | See note |
| PHPINSU | 418 | 1 Yes | |
| | 27298 | 2 No | |
| | 1570 | 8 Blank but applicable | |
| | 28 | Blank | |
| | 1250-1254 | Including your last meal and any snacks, at what time did you last have anything at all to eat? | |
| PHPSNTI | 27701 | 00:00-23:59 | |
| | 1585 | 88888 Blank but applicable | |
| | 28 | Blank | |
| | 1255 | Day participant last ate | |
| PHPSNDA | 12604 | 1 Yesterday | |
| | 15081 | 2 Today | |
| | 16 | 3 Before yesterday | |
| | 1585 | 8 Blank but applicable | |
| | 28 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

PHLEBOTOMY SCREENING QUESTIONNAIRE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 1256 PHPDRIN | | Have you had anything to drink, other than water, after the time you last ate? | |
| | 3947 | 1 Yes | |
| | 23754 | 2 No, subsequent drink fields blank | |
| | 1585 | 8 Blank but applicable | |
| | 28 | Blank | |
| 1257-1261 PHPDRTI | | At what time did you last have anything at all to drink other than water? | |
| | 3947 | 00:00-23:57 | |
| | 1585 | 88888 Blank but applicable | |
| | 23782 | Blank | |
| 1262 PHPDRDA | | Day participant last drank | |
| | 1094 | 1 Yesterday | |
| | 2853 | 2 Today | |
| | 1585 | 8 Blank but applicable | |
| | 23782 | Blank | |
| 1263-1267 PHPFAST | | Computed number of hours since last ate or drank | See note |
| | 27700 | 00000-39.13 | |
| | 1586 | 88888 Blank but applicable | |
| | 28 | Blank | |
| 1268-1272 PHPBEST | | Time of venipuncture | See note |
| | 27703 | 07:32-22:02 | |
| | 1583 | 88888 Blank but applicable | |
| | 28 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

HEMATOLOGY

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|---------------------|---|----------|
| 1273-1277 WCP | 26372 2914 28 | White blood cell count 01.75-71.35 88888 Blank but applicable Blank | See note |
| 1278-1282 WCPSI | 26372 2914 28 | White blood cell count: SI 01.75-71.35 88888 Blank but applicable Blank | |
| 1283-1287 LMPPCNT | 26370 2916 28 | Lymphocyte percent (Coulter) 003.2-083.2 88888 Blank but applicable Blank | |
| 1288-1292 MOPPCNT | 25924 3362 28 | Mononuclear percent (Coulter) 00000-37.55 88888 Blank but applicable Blank | |
| 1293-1297 GRPPCNT | 25925 3361 28 | Granulocyte percent (Coulter) 010.9-093.9 88888 Blank but applicable Blank | |
| 1298-1302 LMP | 26370 2916 28 | Lymphocyte number (Coulter) 00.35-048.1 88888 Blank but applicable Blank | See note |
| 1303-1306 MOP | 25924 3362 28 | Mononuclear number (Coulter) 0000-06.4 8888 Blank but applicable Blank | See note |
| 1307-1311 GRP | 25925 3361 28 | Granulocyte number (Coulter) 000.2-023.4 88888 Blank but applicable Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

HEMATOLOGY

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|---------------------|---|----------|
| 1312-1315 RCP | 26370 2916 28 | Red blood cell count 1.69-6.84 8888 Blank but applicable Blank | See note |
| 1316-1319 RCPSI | 26370 2916 28 | Red blood cell count: SI 1.69-6.84 8888 Blank but applicable Blank | |
| 1320-1324 HGP | 26372 2914 28 | Hemoglobin (g/dL) 04.95-019.6 88888 Blank but applicable Blank | See note |
| 1325-1329 HGPSI | 26372 2914 28 | Hemoglobin: SI (g/L) 049.5-00196 88888 Blank but applicable Blank | |
| 1330-1334 HTP | 26370 2916 28 | Hematocrit (%) 016.6-057.6 88888 Blank but applicable Blank | See note |
| 1335-1339 HTPSI | 26370 2916 28 | Hematocrit: SI (L/L=1) 0.166-0.576 88888 Blank but applicable Blank | |
| 1340-1344 MVPSI | 26371 2915 28 | Mean cell volume: SI (fL) 051.2-122.8 88888 Blank but applicable Blank | See note |
| 1345-1349 MCPSI | 26369 2917 28 | Mean cell hemoglobin: SI (pg) 013.6-053.6 88888 Blank but applicable Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

HEMATOLOGY

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|----------|
| 1350-1354 MHP | | Mean cell hemoglobin concentration (g/dL) | See note |
| | 26369 | 25.95-52.35 | |
| | 2917 | 88888 Blank but applicable | |
| | 28 | Blank | |
| 1355-1359 MHPSI | | Mean cell hemoglobin concentration: SI (g/L) | |
| | 26369 | 259.5-523.5 | |
| | 2917 | 88888 Blank but applicable | |
| | 28 | Blank | |
| 1360-1364 RWP | | Red cell distribution width (%) | |
| | 26372 | 007.8-31.95 | |
| | 2914 | 88888 Blank but applicable | |
| | 28 | Blank | |
| 1365-1370 RWPSI | | Red cell distribution width: SI (fraction) | |
| | 26372 | 00.078-0.3195 | |
| | 2914 | 888888 Blank but applicable | |
| | 28 | Blank | |
| 1371-1375 PLP | | Platelet count | See note |
| | 26367 | 014.5-00981 | |
| | 2919 | 88888 Blank but applicable | |
| | 28 | Blank | |
| 1376-1380 PLPSI | | Platelet count: SI | |
| | 26367 | 014.5-00981 | |
| | 2919 | 88888 Blank but applicable | |
| | 28 | Blank | |
| 1381-1385 DWP | | Platelet distribution width (%) | |
| | 26200 | 005.8-24.65 | |
| | 3086 | 88888 Blank but applicable | |
| | 28 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

HEMATOLOGY

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------------------|--|----------|
| 1386-1390 PVPSI | 26373 2913 28 | Mean platelet volume: SI (fL) 00003-00043 88888 Blank but applicable Blank | |
| 1391-1393 GRPDIF | 8150 2276 18888 | Segmented neutrophils (percent of 100 cells) 007-090 888 Blank but applicable Blank | See note |
| 1394-1396 LMPDIF | 8150 2276 18888 | Lymphocytes (percent of 100 cells) 004-088 888 Blank but applicable Blank | See note |
| 1397-1398 MOPDIF | 8150 2276 18888 | Monocytes (percent of 100 cells) 00-23 88 Blank but applicable Blank | See note |
| 1399-1400 EOP | 8150 2276 18888 | Eosinophils (percent of 100 cells) 00-51 88 Blank but applicable Blank | See note |
| 1401-1402 BOP | 8150 2276 18888 | Basophils (percent of 100 cells) 00-22 88 Blank but applicable Blank | See note |
| 1403 BLP | 8150 2276 18888 | Blasts (percent of 100 cells) 0 8 Blank but applicable Blank | See note |
| 1404 PRP | 8150 2276 18888 | Promyelocytes (percent of 100 cells) 0 8 Blank but applicable Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

HEMATOLOGY

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------|---|----------|
| MEP | 1405 | Metamyelocytes (percent of 100 cells) | See note |
| | 8150 | 0-2 | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| MLP | 1406 | Myelocytes (percent of 100 cells) | See note |
| | 8150 | 0-1 | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| BAP | 1407-1408 | Bands (percent of 100 cells) | See note |
| | 8150 | 00-22 | |
| | 2276 | 88 Blank but applicable | |
| | 18888 | Blank | |
| LAP | 1409-1410 | Atypical lymphocytes (percent of 100 cells) | See note |
| | 8150 | 00-28 | |
| | 2276 | 88 Blank but applicable | |
| | 18888 | Blank | |
| ANP | 1411 | Anisocytosis (variation of cell size) | See note |
| | 6120 | 0 Normal | |
| | 2030 | 1-4 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| BSP | 1412 | Basophilic stippling | See note |
| | 8047 | 0 Normal | |
| | 103 | 1-3 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| HZP | 1413 | Hypochromia (stain intensity of cell) | See note |
| | 6891 | 0 Normal | |
| | 1259 | 1-4 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

HEMATOLOGY

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|----------|
| ----- | | | |
| PKP | 1414 | Poikilocytosis (cell shape variation) | See note |
| | 7067 | 0 Normal | |
| | 1083 | 1-4 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| POP | 1415 | Polychromatophilia (bluish color of cell) | See note |
| | 7231 | 0 Normal | |
| | 919 | 1-3 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| MRP | 1416 | Macrocytosis (large cell prevalence) | See note |
| | 7569 | 0 Normal | |
| | 581 | 1-3 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| MIP | 1417 | Microcytosis (small cell prevalence) | See note |
| | 6873 | 0 Normal | |
| | 1277 | 1-4 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| SIP | 1418 | Sickle cells | See note |
| | 8135 | 0 Normal | |
| | 15 | 1-3 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| SHP | 1419 | Spherocytosis | See note |
| | 7479 | 0 Normal | |
| | 671 | 1-4 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |
| TTP | 1420 | Target cells | See note |
| | 7620 | 0 Normal | |
| | 530 | 1-4 Gradation to abnormal | |
| | 2276 | 8 Blank but applicable | |
| | 18888 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

HEMATOLOGY

| Positions | | | Item description | |
|-----------|--------|-----|-----------------------|----------|
| SAS name | Counts | | and code | Notes |
| | 1421 | | Toxic granulation | See note |
| TXP | 7839 | 0 | Normal | |
| | 311 | 1-4 | Gradation to abnormal | |
| | 2276 | 8 | Blank but applicable | |
| | 18888 | | Blank | |
| | 1422 | | Vacuolated cells | See note |
| VUP | 8150 | 0 | Normal | |
| | 2276 | 8 | Blank but applicable | |
| | 18888 | | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------------------------|---|----------|
| 1423-1426 PBP | 2342 24476 2468 28 | Lead (ug/dL) 00.7 Below level of detection 0001-71.8 8888 Blank but applicable Blank | |
| 1427-1431 PBPSI | 2342 24476 2468 28 | Lead: SI (umol/L) 0.034 Below level of detection 0.048-3.465 88888 Blank but applicable Blank | |
| 1432-1435 EPP | 26706 2580 28 | Protoporphyrin (ug/dL RBC) 0003-1008 8888 Blank but applicable Blank | |
| 1436-1440 EPPSI | 26706 2580 28 | Protoporphyrin: SI (umol/L RBC) 00.05-17.94 88888 Blank but applicable Blank | |
| 1441-1443 FEP | 26479 2807 28 | Serum iron (ug/dL) 004-338 888 Blank but applicable Blank | See note |
| 1444-1448 FEPSI | 26479 2807 28 | Serum iron: SI (umol/L) 00.72-60.54 88888 Blank but applicable Blank | |
| 1449-1452 TIP | 25802 3484 28 | Serum TIBC (ug/dL) 0069-0866 8888 Blank but applicable Blank | |
| 1453-1458 TIPSI | 25802 3484 28 | Serum TIBC: SI (umol/L) 012.36-0155.1 888888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|----------------------------|---|----------|
| 1459-1462 PXP | 25770 3516 28 | Serum transferrin saturation (%) 00.8-98.5 8888 Blank but applicable Blank | See note |
| 1463-1466 FRP | 13 26380 2893 28 | Serum ferritin (ng/mL) 0002 Below level of detection 0003-3059 8888 Blank but applicable Blank | |
| 1467-1470 FRPSI | 13 26380 2893 28 | Serum ferritin: SI (ug/L) 0002 Below level of detection 0003-3059 8888 Blank but applicable Blank | |
| 1471-1475 FOP | 1 23704 1937 3672 | Serum folate (ng/mL) 000.1 Below level of detection 000.4-00199 88888 Blank but applicable Blank | See note |
| 1476-1480 FOPSI | 1 23704 1937 3672 | Serum folate: SI (nmol/L) 000.2 Below level of detection 000.9-450.9 88888 Blank but applicable Blank | |
| 1481-1484 RBP | 23404 2238 3672 | RBC folate (ng/mL) 0007-1755 8888 Blank but applicable Blank | See note |
| 1485-1490 RBPSI | 23404 2238 3672 | RBC folate: SI (nmol/L) 0015.9-3976.8 888888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions | | Item description | |
|-----------|--------|---------------------------------------|----------|
| SAS name | Counts | and code | Notes |
| 1491-1496 | | Serum vitamin B12 (pg/mL) | |
| VBP | 12024 | 000033-099999 | |
| | 722 | 888888 Blank but applicable | |
| | 16568 | Blank | |
| 1497-1504 | | Serum vitamin B12: SI (pmol/L) | |
| VBPSI | 12024 | 00024.35-73779.26 | |
| | 722 | 88888888 Blank but applicable | |
| | 16568 | Blank | |
| 1505-1508 | | Serum vitamin C (mg/dL) | See note |
| VCP | 20636 | 0000-4.72 | |
| | 2408 | 8888 Blank but applicable | |
| | 6270 | Blank | |
| 1509-1514 | | Serum vitamin C: SI (mmol/L) | |
| VCPSI | 20636 | 000000-000268 | |
| | 2408 | 888888 Blank but applicable | |
| | 6270 | Blank | |
| 1515-1518 | | Serum normalized calcium: SI (mmol/L) | See note |
| ICPSI | 16737 | 0.81-1.95 | |
| | 3022 | 8888 Blank but applicable | |
| | 9555 | Blank | |
| 1519-1522 | | Serum total calcium: SI (mmol/L) | |
| CAPSI | 4 | 1.06 Below level of detection | |
| | 18490 | 1.57-3.29 | |
| | 1265 | 8888 Blank but applicable | |
| | 9555 | Blank | |
| 1523-1526 | | Serum selenium (ng/mL) | See note |
| SEP | 18597 | 0039-0622 | |
| | 1619 | 8888 Blank but applicable | |
| | 9098 | Blank | |
| 1527-1530 | | Serum selenium: SI (nmol/L) | |
| SEPSI | 18597 | 00.5-07.9 | |
| | 1619 | 8888 Blank but applicable | |
| | 9098 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|----------------------------|--|----------|
| 1531-1533 VAP | 23274 2368 3672 | Serum vitamin A (ug/dL) 002-259 888 Blank but applicable Blank | |
| 1534-1537 VAPSI | 23274 2368 3672 | Serum vitamin A: SI (umol/L) 0.07-9.04 8888 Blank but applicable Blank | |
| 1538-1542 VEP | 23274 2368 3672 | Serum vitamin E (ug/dL) 00028-09999 88888 Blank but applicable Blank | See note |
| 1543-1548 VEPSI | 23274 2368 3672 | Serum vitamin E: SI (umol/L) 000.65-232.18 888888 Blank but applicable Blank | |
| 1549-1551 ACP | 23274 2368 3672 | Serum alpha carotene (ug/dL) 000-202 888 Blank but applicable Blank | |
| 1552-1555 ACPSI | 23274 2368 3672 | Serum alpha carotene: SI (umol/L) 0000-3.76 8888 Blank but applicable Blank | |
| 1556-1559 BCP | 5 23269 2368 3672 | Serum beta carotene (ug/dL) 0000 Below level of detection 0001-0674 8888 Blank but applicable Blank | See note |
| 1560-1564 BCPSI | 5 23269 2368 3672 | Serum beta carotene: SI (umol/L) 00.00 Below level of detection 00.02-12.56 88888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------------------------|--|----------|
| 1565-1567 BXP | 23272 2370 3672 | Serum beta cryptoxanthin (ug/dL) 000-144 888 Blank but applicable Blank | |
| 1568-1571 BXPSI | 23272 2370 3672 | Serum beta cryptoxanthin: SI (umol/L) 0000-02.6 8888 Blank but applicable Blank | |
| 1572-1574 LUP | 3 23271 2368 3672 | Serum lutein/zeaxanthin (ug/dL) 000 Below level of detection 001-478 888 Blank but applicable Blank | See note |
| 1575-1578 LUPSI | 3 23271 2368 3672 | Serum lutein/zeaxanthin: SI (umol/L) 0.00 Below level of detection 0.02-08.4 8888 Blank but applicable Blank | |
| 1579-1581 LYP | 25 23249 2368 3672 | Serum lycopene (ug/dL) 000 Below level of detection 001-124 888 Blank but applicable Blank | See note |
| 1582-1585 LYPSI | 25 23249 2368 3672 | Serum lycopene: SI (umol/L) 0.00 Below level of detection 0.02-2.31 8888 Blank but applicable Blank | |
| 1586-1588 REP | 23274 2368 3672 | Serum sum retinyl esters (ug/dL) 000-269 888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|--|-----------------------|--|----------|
| 1589-1592 REPSI | 23274 2368 3672 | Serum sum retinyl esters: SI (umol/L) 0000-9.39 8888 Blank but applicable Blank | |
| 1593-1597 COP | 29314 | Serum cotinine (ng/mL) Blank | See note |
| *Note: See LAB2 file for Updated Serum Cotinine Data | | | |
| 1598-1600 TCP | 23561 2081 3672 | Serum cholesterol (mg/dL) 059-702 888 Blank but applicable Blank | |
| 1601-1605 TCPSI | 23561 2081 3672 | Serum cholesterol: SI (mmol/L) 01.53-18.15 88888 Blank but applicable Blank | |
| 1606-1609 TGP | 23515 2127 3672 | Serum triglycerides (mg/dL) 0013-3616 8888 Blank but applicable Blank | See note |
| 1610-1614 TGPSI | 23515 2127 3672 | Serum triglycerides: SI (mmol/L) 00.15-40.82 88888 Blank but applicable Blank | |
| 1615-1617 LCP | 7891 2254 19169 | Serum LDL cholesterol (mg/dL) 020-380 888 Blank but applicable Blank | See note |
| 1618-1621 LCPSI | 7891 2254 19169 | Serum LDL cholesterol: SI (mmol/L) 0.52-9.83 8888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|------------------------|---|----------|
| 1622-1624 HDP | 23409 2233 3672 | Serum HDL cholesterol (mg/dL) 008-196 888 Blank but applicable Blank | |
| 1625-1628 HDPSI | 23409 2233 3672 | Serum HDL cholesterol: SI (mmol/L) 0.21-5.07 8888 Blank but applicable Blank | |
| 1629-1631 AAP | 11432 1464 16418 | Serum apolipoprotein AI (mg/dL) 059-300 888 Blank but applicable Blank | See note |
| 1632-1635 AAPSI | 11432 1464 16418 | Serum apolipoprotein AI: SI (g/L) 0.59-0003 8888 Blank but applicable Blank | |
| 1636-1638 ABP | 11483 1413 16418 | Serum apolipoprotein B (mg/dL) 040-260 888 Blank but applicable Blank | See note |
| 1639-1642 ABPSI | 11483 1413 16418 | Serum apolipoprotein B: SI (g/L) 00.4-02.6 8888 Blank but applicable Blank | |
| 1643-1645 LPP | 12018 728 16568 | Serum lipoprotein (a) (mg/dL) 000-276 888 Blank but applicable Blank | |
| 1646-1649 LPPSI | 12018 728 16568 | Serum lipoprotein (a): SI (g/L) 0000-2.76 8888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

GENERAL BIOCHEMISTRY TESTS

| Positions | | Item description | |
|-----------|--------|--|-------|
| SAS name | Counts | and code | Notes |
| ----- | | | |
| 1650-1654 | | Serum follicle stimulating hormone: SI | |
| FHPSI | | (IU/L) | |
| | 6 | 000.1 Below level of detection | |
| | 3116 | 000.2-00170 | |
| | 253 | 88888 Blank but applicable | |
| | 25939 | Blank | |
| 1655-1658 | | Serum luteinizing hormone: SI (IU/L) | |
| LHPSI | 2 | 00.1 Below level of detection | |
| | 3118 | 00.2-67.1 | |
| | 255 | 8888 Blank but applicable | |
| | 25939 | Blank | |
| 1659-1662 | | Plasma fibrinogen (mg/dL) | |
| FBP | 9350 | 0019-0957 | |
| | 810 | 8888 Blank but applicable | |
| | 19154 | Blank | |
| 1663-1666 | | Plasma fibrinogen: SI (g/L) | |
| FBPSI | 9350 | 0.19-9.57 | |
| | 810 | 8888 Blank but applicable | |
| | 19154 | Blank | |
| 1667-1671 | | Serum C-reactive protein (mg/dL) | |
| CRP | 16218 | 00.21 Below level of detection | |
| | 6249 | 000.3-025.2 | |
| | 3175 | 88888 Blank but applicable | |
| | 3672 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

ANTIBODY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 1672-1677 | | Serum tetanus antibody (U/mL) | |
| TEP | 19336 | 000000-074.67 | |
| | 5849 | 888888 Blank but applicable | |
| | 4129 | Blank | |
| 1678 | | Serum hepatitis A antibody (anti-HAV) | |
| AHP | 9872 | 1 Positive | |
| | 11376 | 2 Negative | |
| | 12 | 3 Borderline | |
| | 1784 | 8 Blank but applicable | |
| | 6270 | Blank | |
| 1679 | | Serum hepatitis B core antibody (anti-HBc) | See note |
| HBP | 1368 | 1 Positive | |
| | 19886 | 2 Negative | |
| | 11 | 3 Borderline | |
| | 1779 | 8 Blank but applicable | |
| | 6270 | Blank | |
| 1680-1681 | | Serum hepatitis B surface antibody (anti-HBs) | See note |
| SSP | 593 | 01 Positive | |
| | 160 | 02 Negative | |
| | 108 | 03 Borderline | |
| | 340 | 11 > 10 mIU | |
| | 155 | 22 < 10 mIU | |
| | 1856 | 88 Blank but applicable | |
| | 26102 | Blank | |
| 1682 | | Serum hepatitis B surface antigen (HBsAg) | See note |
| SAP | 82 | 1 Positive | |
| | 1292 | 2 Negative | |
| | 1 | 3 Borderline | |
| | 1837 | 8 Blank but applicable | |
| | 26102 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

ANTIBODY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------|--|----------|
| HCP | 1683 | Serum hepatitis C antibody (anti-HCV) | |
| | 402 | 1 Positive | |
| | 20796 | 2 Negative | |
| | 43 | 4 Indeterminate | |
| | 1803 | 8 Blank but applicable | |
| | 6270 | Blank | |
| DHP | 1684 | Serum hepatitis D antibody (anti-HDV) | See note |
| | 3 | 1 Positive | |
| | 76 | 2 Negative | |
| | 4 | 8 Blank but applicable | |
| | 29231 | Blank | |
| H1P | 1685 | Serum herpes I antibody | |
| | 9843 | 1 Positive | |
| | 3205 | 2 Negative | |
| | 50 | 3 Indeterminate | |
| | 3184 | 8 Blank but applicable | |
| H2P | 1686 | Serum herpes II antibody | |
| | 3532 | 1 Positive | |
| | 9476 | 2 Negative | |
| | 86 | 3 Indeterminate | |
| | 3188 | 8 Blank but applicable | |
| RUP | 1687-1691 | Serum rubella antibody | See note |
| | 21288 | 00000-72.91 | |
| | 2213 | 88888 Blank but applicable | |
| RUPUNIT | 1692-1695 | Serum rubella antibody (IU) | See note |
| | 21288 | 0000-1224 | |
| | 2213 | 8888 Blank but applicable | |
| | 5813 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

ANTIBODY TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 1696-1700 VRP | 21288 | Serum varicella antibody 00000-29.64 | See note |
| | 2213 | 88888 Blank but applicable | |
| | 5813 | Blank | |
| 1701-1703 TOP | 17658 | Serum toxoplasmosis antibody 000-240 | See note |
| | 2558 | 888 Blank but applicable | |
| | 9098 | Blank | |
| 1704-1708 RFP | 5271 | Serum rheumatoid factor antibody 00000-40960 | |
| | 437 | 88888 Blank but applicable | |
| | 23606 | Blank | |
| 1709-1713 L1P | 5524 | Serum latex antibody (IU/mL) 00000-47.48 | See note |
| | 23790 | Blank | |
| 1714 HPP | 848 | Serum helicobacter pylori antibody 1 Positive | See note |
| | 1733 | 2 Negative | |
| | 125 | 3 Equivocal | |
| | 26608 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

BIOCHEMISTRY PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------------------|--|-------|
| 1715-1719 NAPSI | 18723 1493 9098 | Serum sodium: SI (mmol/L) 123.4-177.5 88888 Blank but applicable Blank | |
| 1720-1723 SKPSI | 18723 1493 9098 | Serum potassium: SI (mmol/L) 2.51-6.94 8888 Blank but applicable Blank | |
| 1724-1728 CLPSI | 18723 1493 9098 | Serum chloride: SI (mmol/L) 076.2-121.6 88888 Blank but applicable Blank | |
| 1729-1730 C3PSI | 18721 1495 9098 | Serum bicarbonate: SI (mmol/L) 04-53 88 Blank but applicable Blank | |
| 1731-1734 SCP | 18722 1494 9098 | Serum total calcium (mg/dL) 06.6-15.4 8888 Blank but applicable Blank | |
| 1735-1739 SCPSI | 18722 1494 9098 | Serum total calcium: SI (mmol/L) 01.65-03.85 88888 Blank but applicable Blank | |
| 1740-1743 PSP | 18723 1493 9098 | Serum phosphorus (mg/dL) 01.5-10.5 8888 Blank but applicable Blank | |
| 1744-1748 PSPSI | 18723 1493 9098 | Serum phosphorus: SI (mmol/L) 0.484-03.39 88888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

BIOCHEMISTRY PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|-----------------------|--|----------|
| 1749-1752 UAP | 18723 1493 9098 | Serum uric acid (mg/dL) 00.2-15.9 8888 Blank but applicable Blank | |
| 1753-1757 UAPSI | 18723 1493 9098 | Serum uric acid: SI (umol/L) 011.9-945.7 88888 Blank but applicable Blank | |
| 1758-1760 SGP | 18719 1497 9098 | Serum glucose (mg/dL) 037-571 888 Blank but applicable Blank | See note |
| 1761-1765 SGPSI | 18719 1497 9098 | Serum glucose: SI (mmol/L) 02.05-031.7 88888 Blank but applicable Blank | |
| 1766-1768 BUP | 18723 1493 9098 | Serum blood urea nitrogen (mg/dL) 002-104 888 Blank but applicable Blank | |
| 1769-1773 BUPSI | 18723 1493 9098 | Serum blood urea nitrogen: SI (mmol/L) 00.71-37.13 88888 Blank but applicable Blank | |
| 1774-1777 TBP | 18723 1493 9098 | Serum total bilirubin (mg/dL) 0000-10.4 8888 Blank but applicable Blank | |
| 1778-1783 TBPSI | 18723 1493 9098 | Serum total bilirubin: SI (umol/L) 000000-177.84 888888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

BIOCHEMISTRY PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|------------------------|---|----------|
| 1784-1787 CEP | 18722 1494 9098 | Serum creatinine (mg/dL) 00.3-13.9 8888 Blank but applicable Blank | |
| 1788-1793 CEPSI | 18722 1494 9098 | Serum creatinine: SI (umol/L) 0026.5-1228.8 888888 Blank but applicable Blank | |
| 1794-1796 SFP | 14056 1493 13765 | Serum iron (ug/dL) 000-464 888 Blank but applicable Blank | See note |
| 1797-1800 SFPSI | 14056 1493 13765 | Serum iron: SI (umol/L) 0000-83.1 8888 Blank but applicable Blank | |
| 1801-1804 CHP | 18721 1495 9098 | Serum cholesterol (mg/dL) 0039-0748 8888 Blank but applicable Blank | See note |
| 1805-1810 CHPSI | 18721 1495 9098 | Serum cholesterol: SI (mmol/L) 01.009-19.343 888888 Blank but applicable Blank | |
| 1811-1814 TRP | 14056 1493 13765 | Serum triglycerides (mg/dL) 0003-3900 8888 Blank but applicable Blank | See note |
| 1815-1820 TRPSI | 14056 1493 13765 | Serum triglycerides: SI (mmol/L) 00.034-44.031 888888 Blank but applicable Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

BIOCHEMISTRY PROFILE

| Positions | | Item description | |
|-----------|--------|--------------------------------------|----------|
| SAS name | Counts | and code | Notes |
| 1821-1823 | | Serum aspartate aminotransferase: SI | |
| ASPSI | | (U/L) | |
| | 18723 | 006-517 | |
| | 1493 | 888 Blank but applicable | |
| | 9098 | Blank | |
| 1824-1826 | | Serum alanine aminotransferase: SI | |
| ATPSI | | (U/L) | |
| | 18723 | 001-486 | |
| | 1493 | 888 Blank but applicable | |
| | 9098 | Blank | |
| 1827-1830 | | Serum gamma glutamyl transferase: SI | See note |
| GGPSI | | (U/L) | |
| | 14549 | 0001-1342 | |
| | 1495 | 8888 Blank but applicable | |
| | 13270 | Blank | |
| 1831-1834 | | Serum lactate dehydrogenase: SI | |
| LDPSI | | (U/L) | |
| | 18721 | 0029-0970 | |
| | 1495 | 8888 Blank but applicable | |
| | 9098 | Blank | |
| 1835-1838 | | Serum alkaline phosphatase: SI (U/L) | |
| APPSI | | | |
| | 18721 | 0017-0952 | |
| | 1495 | 8888 Blank but applicable | |
| | 9098 | Blank | |
| 1839-1842 | | Serum total protein (g/dL) | |
| TPP | | | |
| | 18723 | 04.6-10.4 | |
| | 1493 | 8888 Blank but applicable | |
| | 9098 | Blank | |
| 1843-1845 | | Serum total protein: SI (g/L) | |
| TPPSI | | | |
| | 18723 | 046-104 | |
| | 1493 | 888 Blank but applicable | |
| | 9098 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

BIOCHEMISTRY PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|------------------------|--|----------|
| 1846-1848 AMP | 18723 1493 9098 | Serum albumin (g/dL) 0.9-6.1 888 Blank but applicable Blank | |
| 1849-1851 AMPSI | 18723 1493 9098 | Serum albumin: SI (g/L) 009-061 888 Blank but applicable Blank | |
| 1852-1854 GBP | 14056 1493 13765 | Serum globulin (g/dL) 1.5-6.6 888 Blank but applicable Blank | See note |
| 1855-1857 GBPSI | 14056 1493 13765 | Serum globulin: SI (g/L) 015-066 888 Blank but applicable Blank | |
| 1858-1860 OSPSI | 14056 1493 13765 | Serum osmolality: SI (mmol/Kg) 241-352 888 Blank but applicable Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DIABETES TESTING PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--|---|----------|
| 1861-1864 GHP | 23476 2166 3672 | Glycated hemoglobin: (%) 02.8-16.2 8888 Blank but applicable Blank | See note |
| 1865 GHPMETH | 13892 4811 2549 2224 2166 3672 | Glycated hemoglobin: test method 1 Diamat method (instrument 1) 2 Diamat method (instrument 2) 3 Diamat method (instrument 3) 4 Affinity method 8 Blank but applicable Blank | See note |
| 1866-1870 G1P | 15877 674 12763 | Plasma glucose - first venipuncture (mg/dL) 035.4-642.6 88888 Blank but applicable Blank | See note |
| 1871-1876 G1PSI | 15877 674 12763 | Plasma glucose - first venipuncture: SI (mmol/L) 01.965-35.671 888888 Blank but applicable Blank | |
| 1877-1878 G1PCODE | 2 11 301 98 42 142 12 187 368 28151 | Incomplete glucose test (OGTT) code 20 Hemophiliac 21 Chemotherapy within 4 weeks 22 Diabetic on insulin 23 Refused venipuncture 24 Ill/faint during test 25 Venipuncture unsuccessful 26 Physician canceled test 27 Refused glucose challenge 99 All remaining reasons Blank | See note |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DIABETES TESTING PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|--|----------|
| 1879-1881 G1PTIM1 | | Minutes between glucose challenge and second venipuncture | See note |
| | 6640 | 086-178 | |
| | 855 | 888 Blank but applicable | |
| | 21819 | Blank | |
| 1882-1884 G1PTIM2 | | Minutes between first and second venipuncture | See note |
| | 6637 | 094-184 | |
| | 858 | 888 Blank but applicable | |
| | 21819 | Blank | |
| 1885-1889 G2P | | Plasma glucose - second venipuncture (mg/dL) | See note |
| | 6652 | 033.7-755.1 | |
| | 843 | 88888 Blank but applicable | |
| | 21819 | Blank | |
| 1890-1895 G2PSI | | Plasma glucose - second venipuncture: SI (mmol/L) | |
| | 6652 | 01.871-41.916 | |
| | 843 | 888888 Blank but applicable | |
| | 21819 | Blank | |
| 1896-1900 C1P | | Serum C-peptide - first venipuncture (pmol/mL) | See note |
| | 63 | 0.021 Below level of detection | |
| | 15730 | 00.03-12.77 | |
| | 758 | 88888 Blank but applicable | |
| | 12763 | Blank | |
| 1901-1905 C1PSI | | Serum C-peptide - first venipuncture: SI (nmol/L) | |
| | 63 | 0.021 Below level of detection | |
| | 15730 | 00.03-12.77 | |
| | 758 | 88888 Blank but applicable | |
| | 12763 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DIABETES TESTING PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|----------|
| 1906-1911 C2P | | Serum C-peptide - second venipuncture (pmol/mL) | See note |
| | 1 | 00.021 Below level of detection | |
| | 3365 | 00.075-15.363 | |
| | 381 | 888888 Blank but applicable | |
| | 25567 | Blank | |
| 1912-1917 C2PSI | | Serum C-peptide - second venipuncture: SI (nmol/L) | |
| | 1 | 00.021 Below level of detection | |
| | 3365 | 00.075-15.363 | |
| | 381 | 888888 Blank but applicable | |
| | 25567 | Blank | |
| 1918-1923 I1P | | Serum insulin - first venipuncture (uU/mL) | See note |
| | 65 | 001.76 Below level of detection | |
| | 15689 | 002.51-002367 | |
| | 797 | 888888 Blank but applicable | |
| | 12763 | Blank | |
| 1924-1930 I1PSI | | Serum insulin - first venipuncture: SI (pmol/L) | |
| | 65 | 0010.56 Below level of detection | |
| | 15689 | 0015.06-0014202 | |
| | 797 | 8888888 Blank but applicable | |
| | 12763 | Blank | |
| 1931 I1P2PFLG | | Serum insulin - first venipuncture: test kit | See note |
| | 2693 | 1 Kit 1 | |
| | 1906 | 2 Kit 2 | |
| | 11156 | 3 Kit 3 | |
| | 796 | 8 Blank but applicable | |
| | 12763 | Blank | |
| 1932-1937 I2P | | Serum insulin - second venipuncture (uU/mL) | See note |
| | 3378 | 0002.7-823.01 | |
| | 369 | 888888 Blank but applicable | |
| | 25567 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

DIABETES TESTING PROFILE

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|---|-------|
| 1938-1944 I2PSI | | Serum insulin - second venipuncture: SI (pmol/L) | |
| | 3378 | 00016.2-4938.06 | |
| | 369 | 8888888 Blank but applicable | |
| | 25567 | Blank | |

NHANES III Laboratory Data File
Whole Blood, Serum, Plasma, and Urine Data

URINE TESTS

| Positions SAS name | Counts | Item description and code | Notes |
|-----------------------|--------|----------------------------------|----------|
| 1945-1949 UDP | | Urinary cadmium (ng/mL) | |
| | 22321 | 00.01-16.65 | |
| | 749 | 888888 Blank but applicable | |
| | 6244 | Blank | |
| 1950-1955 UDPSI | | Urinary cadmium: SI (nmol/L) | |
| | 22321 | 000.09-148.14 | |
| | 749 | 888888 Blank but applicable | |
| | 6244 | Blank | |
| 1956-1960 URP | | Urinary creatinine (mg/dL) | See note |
| | 83 | 007.9 Below level of detection | |
| | 22162 | 011.3-682.1 | |
| | 825 | 88888 Blank but applicable | |
| | 6244 | Blank | |
| 1961-1964 URPSI | | Urinary creatinine: SI (mmol/L) | |
| | 83 | 00.7 Below level of detection | |
| | 22162 | 0001-60.3 | |
| | 825 | 8888 Blank but applicable | |
| | 6244 | Blank | |
| 1965-1970 UBP | | Urinary albumin (ug/mL) | |
| | 386 | 0000.4 Below level of detection | |
| | 21859 | 0000.5-015700 | |
| | 825 | 888888 Blank but applicable | |
| | 6244 | Blank | |
| 1971-1977 UIP | | Urinary iodine (ug/dL) | |
| | 5 | 00000.1 Below level of detection | |
| | 22085 | 00000.5-0019750 | |

| | | |
|------|---------|----------------------|
| 980 | 8888888 | Blank but applicable |
| 6244 | Blank | |

DEMOGRAPHIC DATA: NOTES

Screener Questionnaire

DMPFSEQ: Family sequence number

This variable can be used to determine all family members who participated in the survey. Sample persons who have identical family sequence numbers (i.e. match on all 5 digits) are members of the same family.

DMPSTAT: Examination/interview status

This variable identifies the interview or examination status of all persons selected for the NHANES III sample. Interviewed persons completed preselected questions in specific sections of the Household Adult or Youth Questionnaires. Mobile examination center (MEC)-examined persons were interviewed and successfully completed at least one examination component in the MEC. Home-examined persons were interviewed and successfully completed at least one home examination component. The home examination was an option for frail older adults, infants 2-11 months of age, and other adults who were unable to come to the MEC.

DMARETHN: Race-ethnicity

This key analytic variable, based on the NHANES III survey design, was derived from many sources of data and is based on reported race and ethnicity. The other category includes all Hispanics, regardless of race, who were not Mexican-American and also includes all non-Hispanics from racial groups other than white or black.

DMARACER: Race

This variable was obtained from two primary sources: the Screener and the Family Questionnaires. Prior to the selection of the sample, race (Black, White, Other) was self-reported or reported by proxy in the Screener Questionnaire. During the administration of the Family Questionnaire, race was self-reported or reported by the respondent of the Family Questionnaire from five categories (Aleut, Eskimo, American Indian, Asian or Pacific Islander, Black, White, Other). Responses from the two sources were adjudicated, as necessary, to create a three level variable (Black, White, Other).

DMAETHNR: Ethnicity

This variable was obtained from two primary sources: the Screener and the Family Questionnaires. As part of both interviews, hand cards were used to determine Mexican/Mexican-American or Other Latin American/Spanish ancestry or national origin. Responses of non-Hispanic ancestry or national origin were categorized as other. Responses from the two interviews were adjudicated, as necessary, and this three level variable was created.

HSAGEIR: Age (Screener Questionnaire)

Age was calculated using the birth date which was obtained from the Screener Questionnaire. The variable HSAGEU provides the age unit (months or years) for HSAGEIR. Ages of 90 years or greater were recoded into a single category of 90+ years to help protect the confidentiality of survey participants.

HSAITMOR: Age in months (Screener Questionnaire)

Age in months was calculated by computing number of months between the Screener Questionnaire date and date of birth. This variable was created for analyses where exact age at the interview may be needed. HSAITMOR differs slightly from the age in years (HSAGEIR), the variable most often used for analyses. Ages of 1080 months and older (90 years and older) were recoded into a single category of 1080+ months to protect the confidentiality of survey participants.

HSFSIZER: Family Size

Family size represents the total number of related persons living in a household (single dwelling unit). All household members were rostered by family during the Screener interview. Household members who were related to the family reference person (knowledgeable household member 17 years or older who owned or rented the dwelling unit) by blood or marriage were considered part of the family. Adopted children, foster- and god-children were also included, if they were living in the dwelling unit. However, family members who were away at college, or living independently were not included. Other household members who were unrelated to the reference person were considered members of separate families. Families with 10 members or more were recoded into a single response category of 10+ persons to help protect confidentiality. See note for Household Size (HSHSIZER).

HSHSIZER: Household Size

Household size represents the total number of persons living in a single dwelling unit, both related and unrelated. All permanent household members were rostered according to their family as part the Screener interview. This was done in order to obtain a complete list of all persons living or staying in the dwelling unit, and to distinguish household and family members. Households with 10 members or more were recoded into a single response category of 10+ persons to help protect confidentiality. See note for Family Size (HFHSIZER).

DMPCNTYR: County FIPS codes for United States counties with populations of
500,000 and more

These county FIPS codes identify large counties with populations of 500,000 and more that were sampled in the survey. Counties with

population less than 500,000 are not included to prevent identification of these locations. See Appendix 1 for listing of codes.

DMPFIPSR: State FIPS codes for United States counties with populations of 500,000 and more

These state FIPS codes identify counties with populations of 500,000+ that were sampled in the survey. Counties with population less than 500,000 are not included to prevent identification of these locations. See Appendix 1 for listing of codes.

DMPMETRO: Urbanization classification based on USDA Rural-Urban continuum codes

These classifications are based on the USDA Rural-Urban codes (Butler and Beale, 1993) that describe metro and nonmetro counties by degree of urbanization and nearness to metro areas. The USDA codes were recoded into two categories to prevent identification of counties that were sampled in the survey.

DMPCREGN: Census region

The United States was divided into four broad geographic regions as defined by the Bureau of Census. Because all states were not included in the selected sample, regional estimates may not be representative for a given region.

DMPPIR: Poverty income ratio (or poverty index)

The poverty income ratio (PIR) was computed as a ratio of two components. The numerator was the midpoint of the observed family income category in the Family Questionnaire variable:HFF19R. The denominator was the poverty threshold, the age of the family reference person, and the calendar year in which the family was interviewed.

Poverty threshold values (in dollars) are produced annually by the Census Bureau (Series P-60). These threshold values are based on calendar years and adjusted for changes caused by inflation between calendar years. Reports for each of the calendar years in the survey (1988-94) were used in the calculation of PIR. For the years 1991 and 1994, data from preliminary reports were used. The poverty income ratio allows income data to be analyzed in a comparable manner across the six years of the survey and with previous NHANES.

Persons who reported having had no income and were assigned a zero value for PIR. A substantial proportion of persons refused to report their income or income category during the Family Questionnaire. Due to the income nonresponse the potential for bias in PIR may be high. Users are cautioned to examine potential nonresponse bias for PIR and other income variables.

Survey Design Data

SDPPHASE: Phase of NHANES III survey

For operational purposes, 81 primary sampling units were divided into 89 survey locations (or stands) and randomly allocated to two three-year phases. Phase 1 data were collected from October 1988 through October 1991 and Phase 2 data were collected from October 1991 through October 1994.

SDPSTRA6, SDPSTRA1, SDPSTRA2, and SDPPSU6, SDPPSU1, SDPPSU2: Pseudo strata codes and pseudo PSU pair codes

Because NHANES III was based upon a complex sample design, the assumptions of many statistical tests and routinely available statistical programs are not met. For this reason, when estimates of the variances of statistics are computed, the technique of estimation must be based upon complex sampling theory. In order to provide users with the capability of estimating the complex sample variances, 49 pseudo strata and a pair of Primary Sampling Unit (PSU) codes per stratum were designed.

A software package, "SUDAAN- Software for the Statistical Analysis of Correlated Data" (Shah, 1995), was developed by the Research Triangle Institute to analyze complex sample design data like NHANES. SUDAAN uses strata and PSU codes to conduct analysis with two PSU per stratum design. Therefore, definition of pseudo strata and PSU provided in this data file should be used to compute complex sample variances in analyses. Other software available for estimation of complex sample variance may also be used. For further discussion of methods of variance estimation in NHANES III, see additional information on this subject in Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

Sampling Weights

WTPFQX6, WTPFQX1, WTPFQX2: Total NHANES III and phase-specific final interview weights

These sampling weights should be used only for items collected during the household interviews. To compute final interview weights, final basic weights were first adjusted for nonresponse to household interview, then post-stratified to the unpublished Current Population Survey 1990 (Phase 1) and 1993 (Phase 2) population control estimates of the U.S. population adjusted for undercount. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFEX6, WTPFEX1, WTPFEX2: Total NHANES III and phase-specific final MEC examination weights

These MEC sampling weights should be used for analysis of measurements or interview items collected in the MEC. Persons who were not examined in the MEC have a sampling weight of zero and should be excluded from analyses. To compute final MEC examination weights, final interview weights were first adjusted for nonresponse to MEC examinations, then post-stratified to the unpublished Current Population Survey 1990 (Phase 1) and 1993 (Phase 2) population control estimates of the U.S. population adjusted for undercount. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFHX6, WTPFHX1, WTPFHX2: Total NHANES III and phase-specific MEC+home examination weights

These MEC+home sampling weights should be used for analysis of the examination items where measurements or interview items were collected in the MEC and home. Persons who were not examined in the MEC or home have a sampling weight of zero and should be excluded from analyses. To compute final MEC+home examination weights, final interview weights were first adjusted for nonresponse to MEC and home examinations, then post-stratified to unpublished Current Population Survey 1990 (Phase 1) and 1993 (Phase 2) population control estimates of the U.S. population adjusted for undercount. No separate sampling weights were computed for home examinees. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFALG6, WTPFALG1, WTPFALG2: Total NHANES III and phase-specific allergy examination subsample weights

These subsample weights are for analysis of allergy measurements. Allergy skin reactivity tests were administered to all MEC-examined persons aged 6-19 years and a random half-sample of the adults aged 20-59 years. Eligible MEC-examined persons who did not complete the allergy tests have a sampling weight of zero and should be excluded from the analyses. Final MEC examination weights were first adjusted for selection of the half-sample among adults (20-59 years), and post-stratified to the unpublished Current Population Survey 1990 (Phase 1) and 1993 (Phase 2) population control estimates of the U.S. population adjusted for undercount in the final step. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFCNS6, WTPFCNS1, WTPFCNS2: Total NHANES III and phase-specific central nervous system (CNS) examination subsample final weights

These subsample weights are for analysis of measurements from the Central Nervous System (CNS) test. The CNS examination was administered to a random half-sample of the adults aged 20-69 years. Eligible MEC-examined persons who did not complete CNS testing have a sampling weight of zero and should be excluded from the analyses. Final MEC examination weights were first adjusted for selection of half sample among adults (20-59 years), and post-stratified to unpublished Current

Population Survey 1990 (Phase 1) and 1993 (Phase 2) population control estimates of the U.S. population adjusted for undercount in the final step. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFSD6, WTPFSD1, WTPFSD2: Total NHANES III and phase-specific morning session MEC examination subsample final weights

These subsample weights are for special analyses where fasting time may be an important factor. They were computed for persons aged 12 years and older who were scheduled and examined in the MEC morning session. Sampled households in the survey were randomly assigned to one of two groups -- morning session ("standard") or afternoon/evening session ("modified") assignments. All sample persons from a household received the same session assignment and were requested to schedule examinations for the assigned session. Fasting instructions varied by age and session assignment (Plan and Operation of The Third National Health and Nutrition Examination Survey, 1988-94, U.S. DHHS, 1996). It should be noted that actual fasting time may have differed from the instructed fasting time and can be obtained from the variable PHPFAST in the NHANES III Laboratory Data File. To compute these weights, final MEC examination weights were first adjusted for the random half selection, then adjusted for the non-response to assigned session, and finally, post-stratified to the unpublished Current Population Survey 1990 and 1993 Population control estimates of the U.S. population adjusted for undercount. Eligible MEC-examined persons who were assigned to the morning session and examined in another session have a sampling weight of zero and should be excluded in analyses. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFMD6, WTPFMD1, WTPFMD2: Total NHANES III and phase-specific afternoon/evening session MEC examination subsample final weights

These subsample weights are for special analyses where fasting time might be an important factor. They were computed for MEC examined persons aged 12 years and older who were scheduled and examined in the afternoon or evening sessions. Sampled households in the survey were randomly assigned to one of two groups -- morning session ("standard") or afternoon/evening session ("modified") assignments. All sample persons from a household received the same session assignment and were requested to schedule examinations for the assigned session. Fasting instruction varied by age and session assignments (Plan and Operation of the Third National Health and Nutrition Examination Survey, 1988-94, U.S. DHHS, 1996). It should be noted that actual fasting time may have differed from the instructed fasting time and can be obtained from the variable PHPFAST in the NHANES III Laboratory Data File.) compute these weights, final MEC examination weights were first adjusted for the random half selection, then adjusted for the nonresponse to assigned session, and finally, post-stratified to the unpublished Current Population Survey 1990 and 1993 population control estimates of the U.S. population adjusted for undercount. Eligible MEC examined persons who were

assigned to the afternoon or evening sessions and examined in another session have a sampling weight of zero and should be excluded in analyses. For details, see Weighting and Estimation Methodology (U.S.DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFHSD6, WTPFHSD1, WTPFHSD2: Total NHANES III and phase-specific morning session MEC+home examination subsample final post stratified weights

These subsample weights are for special analyses where fasting time may be an important factor. They were computed for MEC+home examined persons aged 12 years and older who were scheduled and examined in the morning session. Sampled households in the survey were randomly assigned to one of two groups -- morning session ("standard") or afternoon/evening session ("modified") assignments. All sample persons from a household received the same session assignment and were requested to schedule examinations for the assigned session. Fasting instruction varied by age and session assignments (Plan and Operations of the Third National Health and Nutrition Examination Survey, 1988-94, U.S. DHHS, 1996). It should be noted that actual fasting time may have differed from the instructed fasting time and can be obtained from the variable PHPFAST in the NHANES III Laboratory Data File. To compute these weights, final MEC+home examination weights were first adjusted for the random half selection, then adjusted for the nonresponse to assigned session, and finally, post-stratified to the unpublished Current Population Survey 1990 and 1993 population control estimates of the U.S. population adjusted for undercount. Eligible MEC+home examined persons who were assigned to the morning session and examined in another session have a sampling weight of zero and should be excluded in analyses. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPFHMD6, WTPFHMD1, WTPFHMD2: Total NHANES III and phase-specific afternoon/evening MEC+home examination subsample final weights

These subsample weights are for special analyses where fasting time may be an important factor. They were computed for MEC+home examined persons aged 12 years and older who were scheduled and examined in the afternoon or evening sessions. Sampled households in the survey were randomly assigned to one of two groups -- morning session ("standard") or afternoon/evening session ("modified") assignments. All sample persons from a household received the same session assignment and were requested to schedule examinations for the assigned session. Fasting instruction varied by age and session assignments (Plan and Operation of the Third National Health and Nutrition Examination Survey, U.S. DHHS, 1996). It should be noted that actual fasting time may have differed from the instructed fasting time. The actual fasting time can be obtained from the variable PHPFAST in the NHANES III Laboratory Data File. To compute these weights, final MEC+home examination weights were first adjusted for the random half selection, then adjusted for the nonresponse to assigned session, and finally, post-stratified to the unpublished Current Population Survey 1990 and 1993 population control estimates of the U.S. population adjusted for undercount. Eligible MEC+home examined persons who were

assigned to the afternoon or evening sessions and examined in another session have a sampling weight of zero and should be excluded in analyses. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPQRP1--WTPQRP52: Fay's BRR Replicate interview sample

To allow for alternative methods to estimate variance, 52 replicate weights were computed using repeated sampling method where WESVAR or other software that use repeated samples, can be used for estimating variance. Fay's method (see Fay, 1990; Judkins, 1990) was used to draw half samples and adjust sampling weights in each of the random half samples. Sampling weights in one half sample were multiplied by the factor $k=1.7$ and in the other half sample by $k=0.3$ using the Fay's method. After this adjustment, sampling weights were further adjusted for non-response and post-stratified using the same procedure as the final full sample interview weights. These weights should be used only for estimating variance of items from the household adult and youth interviews. For details, see Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

WTPXRP1--WTPXRP52: Fay's BRR Replicate weights for MEC- examined sample

To allow for alternative methods to estimate variance, 52 replicate weights were computed using repeated sampling method where WESVAR or other BRR type software can be used to estimate variance. Fay's method (see Fay, 1990; Judkins, 1990) was used to draw half samples and adjust sampling weights in each of the random half samples. Sampling weights in one half sample were multiplied by the factor $k=1.7$ and in the other half sample by $k=0.3$ using Fay's method. After this adjustment, weights were further adjusted for nonresponse and were post-stratified using the same procedure as the full sample final weights. These weights should be used only for estimating variance of outcome measurements or interview items from the MEC Examination. For details, see additional information on this subject in Weighting and Estimation Methodology (U.S. DHHS, 1996) and NHANES III Analytic and Reporting Guidelines (U.S. DHHS, 1996).

Household Youth Questionnaire

HYAITMO: Age in months (Household Youth Interview)

Age in months was calculated by computing number of months between Household Youth Interview date and the date of birth. It was created for special analyses where exact age at the interview may be needed. This computed age may be different from the self-reported age in HSAGEIR and HSAGEU, or HSAITMOR. For most analyses, age reported in HSAGEIR (and HSAGEU) should be used.

MEC Examination

MXPLANG: Language of MEC examination

This variable designates the language of conduct for the MEC examination. Questionnaires were designed to be implemented in a bilingual (English/Spanish) format so that respondents could be interviewed in their preferred language. When it was necessary to conduct an interview in another language, a translator assisted the interviewer in administering the questionnaires. These interviews were coded as other.

MXPSESSR: Examination session for MEC examinees

This variable designates the period during the day that the examination occurred. To increase response rates and allow flexibility, examinations were scheduled in three sessions: morning, afternoon and evening. On occasion, more than one session was attended in order to complete the full examination. In such a situation, the session was coded as the one when most of the examinations were completed.

MXPAXTMR: Age in months at MEC examination

Age in total months was created for special analyses where exact age at the examination may be needed (e.g., computation of growth charts). It was calculated by computing number of months between examination date and the date of birth. Some examinees may have had a birthday between household interview and examination so that this computed age at examination may differ slightly from the age reported in HSAGEIR (and HSAGEU), or HSAITMOR. For most analyses age reported in HSAGEIR (and HSAGEU) should be used. Ages of 1080 months and older (90 years and older) were recoded into a single category of 1080+ months to protect the confidentiality of survey participants.

Home Examination

HXPAXTMR: Age in months at home examination

Age in total months was created for special analyses where exact age at the examination may be needed (e.g., computation of growth charts). It was calculated by computing number of months between examination date and the date of birth. Some examinees may have had a birthday between household interview and examination so that this computed age at examination may differ slightly from the age reported in HSAGEIR (and HSAGEU), or HSAITMOR. For most analyses age reported in HSAGEIR (and HSAGEU) should be used. Ages of 1080 months and older (90 years and older) were recoded into a single category of 1080+ months to protect the confidentiality of survey participants.

HXPSESSR: Examination session for home examinees

This variable designates the period during the day that the examination occurred. To increase response rates and allow flexibility, examinations were scheduled in three sessions: morning, afternoon and evening. On occasion, more than one session was attended in order to complete the full examination. In such a situation, the session was coded as the one when most of the examinations were completed.

References

Butler MA, Beale CL. Rural-urban continuum codes for metro and nonmetro counties, 1993. Agriculture and Rural Economy Division, Economic Research Services, U.S. Department of Agriculture, Staff Report No. AGES-9425, 1993.

Fay RE. VPLX: Variance Estimates for Complex Surveys. In: Proceedings of the Survey Research Methods section of the American Statistical Association, pp. 266-271, 1990.

Judkins DR. Fay's method for variance estimation. Journal of Official Statistics 6 (3):223-239. 1990.

U.S. Department of Health and Human Services(DHHS). National Center for Health Statistics. NHANES III Reference Manuals and Reports (CD-ROM). Hyattsville,Md.: Centers for Disease Control and Prevention, 1996. Available from National Technical Information Service (NTIS), Springfield,Va. (Acrobat.PDF format; includes access software: Adobe Systems Inc. Acrobat Reader 2.1)

Shah BV, Barnwell BG, Bieler GS. SUDAAN User's Manual: Software for Analysis of Correlated Data, Release 6.04. Research Triangle Park, North Carolina. 1995.

Appendix 1. State and county FIPS codes for areas with populations of 500,000 or more.

| DMPFIPSR | State | DMPCNTYR | County |
|----------|---------------|----------|----------------|
| 4 | Arizona | 13 | Maricopa |
| 6 | California | 1 | Alameda |
| 6 | California | 19 | Fresno |
| 6 | California | 37 | Los Angeles |
| 6 | California | 59 | Orange |
| 6 | California | 71 | San Bernardino |
| 6 | California | 73 | San Diego |
| 6 | California | 85 | Santa Clara |
| 6 | California | 111 | Ventura |
| 12 | Florida | 25 | Dade |
| 12 | Florida | 31 | Duval |
| 12 | Florida | 99 | Palm Beach |
| 17 | Illinois | 31 | Cook |
| 25 | Massachusetts | 17 | Middlesex |
| 26 | Michigan | 125 | Oakland |
| 26 | Michigan | 163 | Wayne |
| 29 | Missouri | 189 | St Louis |
| 36 | New York | 29 | Erie |
| 36 | New York | 47 | Kings |
| 36 | New York | 59 | Nassau |
| 36 | New York | 61 | New York |
| 36 | New York | 81 | Queens |
| 36 | New York | 119 | Westchester |
| 39 | Ohio | 35 | Cuyahoga |
| 39 | Ohio | 61 | Hamilton |
| 42 | Pennsylvania | 3 | Allegheny |
| 42 | Pennsylvania | 45 | Delaware |
| 42 | Pennsylvania | 101 | Philadelphia |
| 44 | Rhode Island | 7 | Providence |
| 48 | Texas | 29 | Bexar |
| 48 | Texas | 113 | Dallas |
| 48 | Texas | 141 | El Paso |
| 48 | Texas | 201 | Harris |
| 48 | Texas | 439 | Tarrant |
| 53 | Washington | 33 | King |

LABORATORY DATA: NOTES

AAP: Serum apolipoprotein AI

Apolipoprotein AI and apolipoprotein B results were measured only during 1988-1991. Three different methods were used at different times to measure apolipoprotein AI and apolipoprotein B. These were radial immunodiffusion (RID), rate immunonephelometry (INA), and the World Health Organization -International Federation of Clinical Chemistry (WHO-IFCC) method (Bachorik, 1994; Marcovina, 1991; Albers, 1989). Results using the RID and INA methods were adjusted to the WHO-IFCC method.

ABP: Serum apolipoprotein B

See note for AAP.

ANP: Anisocytosis

Microscopic examination (manual differential) of the peripheral blood spread on a glass slide utilized a stained blood film to perform a differential leukocyte count, evaluate red cell morphology, and estimate number of platelets. Manual differential variables include segmented neutrophils, lymphocytes, monocytes, eosinophils, basophils, blasts, promyelocytes, metamyelocytes, myelocytes, bands, atypical lymphocytes, anisocytosis, basophilic stippling, hypochromia, poikilocytosis, polychromatophilia, macrocytosis, microcytosis, sickle cells, spherocytosis, target cells, toxic granulation, and vacuolated cells (GRPDIF, LMPDIF, MOPDIF, EOP, BAP, BOP, BLP, PRP, MEP, MLP, BAP, LAP, ANP, BSP, HZP, PKP, POP, MRP, MIP, SIP, SHP, TTP, TXP, and VUP).

In NHANES III, a manual differential was performed on a special subsample of examinees aged one year and older. This manual differential was used for internal quality control purposes and to confirm abnormal hematology results. This subsample was defined as a random 10-percent sample of all examined persons plus all examinees who had a predetermined high or low value for one or more of the following hematologic assessments: white blood cell count (WBC), red blood cell count (RBC), hemoglobin, hematocrit, mean cell volume (MCV), mean cell hemoglobin (MCH), mean cell hemoglobin concentration (MCHC), red blood cell distribution width (RDW), platelet count, mean platelet volume (MPV), lymphocyte percentage, mononuclear percentage, or granulocyte percentage. A table of predetermined high and low values for WBC, RBC, hemoglobin, hematocrit, MCV, MCH, MCHC, RDW, platelet count, MPV, lymphocyte percentage, mononuclear percentage, and granulocyte percentage is located in the Manual for Medical Technicians (U.S. DHHS, pp. 5-54 and 5-55, 1996).

BAP: Band cells

See note for ANP.

BCP: Serum beta carotene

The lower limit of detection (LOD) for beta carotene was 0.67 ug/dL. Using the LOD coding formula (detection limit divided by the square root of two), the calculated value to indicate that the serum beta carotene results were below the level of detection would be 0.48. After rounding, the value of 0 (zero) was placed in the results field to indicate that the serum beta carotene was below 0.67 ug/dL.

BLP: Blast cells

See note for ANP.

BOP: Basophil cells

See note for ANP.

BSP: Basophilic stippling

See note for ANP.

C1P: Serum C-peptide (first venipuncture)

The specimen for this assay was obtained at the time of the initial venipuncture. This result is available for all six years of the survey.

Examinees aged 40-74 years who used insulin were excluded from the OGTT. A first venipuncture was obtained, but the glucose challenge and second venipuncture were canceled. In these instances, the variables G1P, C1P and I1P have a value, but the results G2P, C2P and I2P from the second venipuncture are blank-filled to indicate a medical exclusion.

C2P: Serum C-peptide (second venipuncture)

Post-glucose challenge levels of C-peptide and insulin for examinees who had an OGTT were measured only during 1991-1994.

CHP: Serum cholesterol

This value was obtained from the standard battery of biochemical assessments. Use of the laboratory test result from the reference method (TCP), rather than the CHP value, is generally recommended. For most analyses of serum cholesterol, the appropriate variable to use will be TCP. The value from the biochemistry profile (CHP) should not be used routinely. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996) for the details.

COP: Serum cotinine

Only cotinine results from 1988-1991 are included in this field.

DHP: Serum hepatitis D antibody

Hepatitis B virus testing scheme: From 1988-1991, all sera were tested for the core antibody to hepatitis B virus (anti-HBc). If this test was positive, the sera were tested for the hepatitis B surface antigen (HBsAg) and hepatitis B surface antibody (anti-HBs). If the HBsAg test was positive and the anti-HBs test was <10 mIU, then the antibody to hepatitis D virus (anti-HDV) test was performed. If the HBsAg test was negative and the anti-HBs test was <10 mIU, then the anti-HBc test was repeated for confirmation.

In June 1993, all sera were tested for both anti-HBc and anti-HBs. Sera testing positive for anti-HBc were tested further for HBsAg, and positive HBsAg samples were tested for anti-HDV.

EOP: Eosinophil cells

See note for ANP.

FEP: Serum iron

Laboratory methods differed between NHANES III and previous surveys. Therefore, results may not be comparable between surveys. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996).

FOP: Serum folate

Laboratory methods differed between NHANES III and previous surveys. Therefore, results may not be comparable between surveys. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996).

G1P: Plasma glucose (first venipuncture)

Plasma glucose was measured using the reference method on examinees aged 20 years and older. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996) for details.

During NHANES III, OGTT testing was conducted on MEC examinees aged 40-74 years. A random assignment was made prior to conducting the OGTT to determine who should receive a morning examination (NCHS, 1994; U.S. DHHS, 1996). As a result, approximately half of the OGTT examinees received the morning OGTT after an overnight fast. This subsample most closely conformed to the World Health Organization (WHO) criteria for OGTT testing to identify diabetes (WHO, 1995). Therefore, this morning subsample is the NHANES III subsample that should be used to estimate the prevalence of diabetes and impaired glucose tolerance.

People who reported a medical history of diabetes but who were not using insulin therapy were asked to conform to the fasting instructions for their examination session and were eligible for an OGTT if the age criteria were satisfied. The morning sample weights (WTPFHSD6) for total NHANES III weights for the morning OGTT subsample should be used when weighting these data to generate national estimates. Data from the afternoon and evening OGTTs do not conform to the WHO protocol for diagnosing diabetes or IGT and should not be used for these purposes.

If an examinee was given an OGTT during an examination session other than the session assigned, that examinee's sample weight for the assigned session will be zero. For example, if an examinee was selected for a morning OGTT but was tested in the afternoon, the examinee's morning sample weight for the OGTT will be zero.

G1PCODE: Reasons for an incomplete glucose tolerance test

The reason for which an examinee aged 40-74 years did not complete the OGTT was entered in this field. This field either will contain an incomplete OGTT code or will be blank. Examinees who responded affirmatively to the hemophilia question (code 20) or who received chemotherapy within the past four weeks (code 21) were excluded from venipuncture. Examinees who reported on their examination day that they used insulin therapy (code 22) were excluded from the OGTT. Codes 23-27 were recoded from comments and notations on the questionnaires and may not include complete data on these reasons.

G1PTIM1: Interval between glucose drink and second venipuncture in minutes

If an examinee was aged 40-74 years and received the OGTT, the time that the glucose drink was consumed and the time of the second venipuncture were recorded. This variable contains the calculated time difference between the glucose drink consumption and the second venipuncture.

G1PTIM2: Interval between first and second venipuncture in minutes

If an examinee was aged 40-74 years and received the OGTT, two timed venipunctures were performed. This variable contains the calculated time difference between the first and second venipunctures.

G2P: Plasma glucose (second venipuncture)

See notes for C1P and G1P.

GBP: Serum globulin

Globulin results were added to the protocol after NHANES III began. This result field was blank-filled for examinees who were examined prior to the start of testing.

GGPSI: Serum gamma glutamyl transferase

Gamma glutamyl transferase results were added to the protocol after NHANES III began. This result field was blank-filled for examinees who were examined prior to the start of testing.

GHP: Glycated hemoglobin (HbA1c)

Glycohemoglobin measurements for NHANES III were performed by the Diabetes Diagnostic Laboratory at the University of Missouri -- Columbia using the Diamat Analyzer System (Bio-Rad Laboratories, Hercules, CA). This ion-exchange HPLC system measures HbA1c (a specific glycohemoglobin) and has demonstrated excellent, long-term precision (interassay CV's 2.0). It was standardized to the reference method that was used for the Diabetes Control and Complications Trial (DCCT). Variant hemoglobins, including hemoglobin C, D, F, and elevated HbF, can interfere with HbA1c measurement by the Diamat HPLC. Hemoglobin S in its heterozygous state does not interfere with this assay. Although interferences usually can be detected by an abnormal Diamat chromatogram, HbA1c results for these specimens were not considered valid. Therefore, samples containing hemoglobin variants or elevated HbF or samples that produce chromatograms indicating hemoglobin degradation were analyzed by an alternate method that used affinity chromatography to separate glycohemoglobin. Affinity chromatographic methods were not affected by the presence of hemoglobin variants and were less sensitive to hemoglobin degradation due to improper sample handling. The affinity method used also was standardized to the DCCT reference method. Reasons for using the affinity method for an examinee's test included an extra peak on the chromatogram, hemoglobin C, elevated hemoglobin F, or other abnormal hemoglobin.

GHPMETH: Glycated hemoglobin method

See note for GHP.

GRP: Granulocyte number

Consult the Manual for Medical Technicians for the Coulter granulocyte number, lymphocyte number, mononuclear number, white blood cell count, red blood cell count, and platelet count units (U.S. DHHS, 1996).

GRPDIF: Segmented neutrophil cells

See note for ANP.

HBP: Serum hepatitis B core antibody

See note for DHP.

HGP: Hemoglobin

In NHANES I, NHANES II, and HHANES, determinations of red and white blood cell counts were made using a semiautomated cell counter (Coulter model FN). Determinations of hemoglobin concentration (Hb) were made using a Coulter hemoglobinometer, and determinations of packed

cell volume (PCV) were made using the microhematocrit centrifuge method. The hematologic indices MCH, MCHC, and MCV were calculated as follows:

$$\begin{aligned} \text{MCH} &= \text{Hb/RBC} \\ \text{MCHC} &= \text{Hb/PCV} \\ \text{MCV} &= \text{PCV/RBC} \end{aligned}$$

In NHANES III, these hematologic parameters were determined by using a fully automated Coulter S+JR hematology analyzer. These analyzers measured the mean (red) cell volume (MCV) directly, utilizing a process of continuous integration of pulse heights divided by the pulse number; PCV values were calculated through the multiplication of MCV and RBC.

Although it has been shown that identified errors in the microhematocrit method caused by plasma trapping and red cell dehydration approximately compensate each other (Bull, 1990), packing errors can occur in macrocytic anemia and can be considerable in sickle cell anemia, spherocytosis, and thalassemias (NCCLS, 1993). Therefore, individual values for MCV, PCV ("hematocrit"), and MCHC from NHANES III cannot be compared directly to values from the previous NHANES.

HPP: Serum Helicobacter pylori antibody

H. pylori antibody testing was performed on surplus sera from children and adolescents aged 6-19 years. This result field was blank-filled for examinees aged 6-19 years for whom surplus specimens were not available for testing. Due to variability in the laboratory test (Pylori Stat, Whittaker Bioproducts, Inc.), 50 percent of the assays were repeated randomly. There was a seven-percent error rate in which the first result (HPP) did not match the repeat result (HPQ). The original result was kept if the controls on the ELISA plate were within the acceptable range. Testing on adults will be performed at a later date using the same assay.

HTP: Hematocrit

See note for HGP.

HZP: Hypochromia

See note for ANP.

I1P: Serum insulin (first venipuncture)

This is the adjusted insulin value for examinees. Most of the insulin

values in NHANES III (1988-1991) were adjusted because the manufacturer of the laboratory testing kits changed during that period. An indicator of the kit number is located in the I1P2PFLG field (i.e., 1 = Kit 1, 2 = Kit 2, and 3 = Kit 3). All insulin values from Kit 1 and Kit 2 assays were adjusted linearly to match the Kit 3 numbers. Further information on this adjustment procedure is available in the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996).

The equations used to adjust the data were:

$$\text{Kit 3} = 0.787 (\text{Kit 1}) + 0.832 \quad \text{Equation 1}$$

$$\text{Kit 3} = 0.597 (\text{Kit 2}) + 1.746 \quad \text{Equation 2}$$

The following steps were used to make the adjustment:

1. Equation 1 was applied to group 1 (Kit 1) data
2. Equation 2 was applied to group 2 (Kit 2) data
3. Group 3 data (Kit 3) were left unchanged.

The time periods for the insulin kits were as follows:

| Group | Assay Period | Assay Method |
|-------|-----------------------|--------------|
| 1 | 10/88-01/05/90 | Kit 1 |
| 2 | 01/06/90-09/06/90 | Kit 2 |
| 3 | 11/01/90-end of study | Kit 3 |

See note for C1P.

I1P2PFLG: Insulin adjustment flag

This field shows which kit was used for the original insulin measurement.

I2P: Serum insulin (second venipuncture)

See notes for C1P, C2P and I1P.

ICPSI: Serum normalized calcium

This variable contains the normalized calcium value derived from adjusting the measured ionized calcium for pH. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996) for details.

L1P: Serum latex antibody

Latex antibody testing was performed on surplus sera from persons ages 17-60 years who were examined in phase 1 (1988-91). This result field was blank-filled for examinees ages 17-60 years for whom surplus specimens were not available for testing.

LAP: Atypical lymphocyte cells

See note for ANP.

LCP: Serum LDL cholesterol calculation

The value for LDL was calculated by the Friedewald equation as follows:

$LDL = \text{total cholesterol} - \text{high density cholesterol} - \text{triglyceride}/5.$

Because the equation is not valid when serum triglyceride values exceed 400 mg/dL, the LDL is missing when serum triglyceride (TGP) exceeds 400 mg/dL.

Serum LDL was calculated on examinees who were instructed to fast (ages 12 and older) and who did fast at least nine hours, were examined in the morning, and were randomly assigned to the morning fasting sample (WTPFHSD6 > 0). Therefore, LDL would be blank if examinees were aged less than 12 years, fasted fewer than nine hours, were examined in an afternoon or evening session, or were not randomly assigned to the morning session. For the purpose of this calculation, the number of hours fasted was rounded to the nearest whole integer.

For more information regarding this equation, refer to the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996).

LMP: Lymphocyte number

See note for GRP.

LMPDIF: Lymphocyte cells

See note for ANP.

LUP: Serum lutein/zeaxanthin

The lower limit of detection (LOD) for lutein/zeaxanthin was 0.43 ug/dL. Using the LOD coding formula (detection limit divided by the square root of two), the calculated value indicating that the serum lycopene results were below the level of detection would be 0.30. After rounding, the value of 0 (zero) was placed in the results field to indicate that the serum lutein/zeaxanthin was below 0.43 ug/dL.

LYP: Serum lycopene

The lower limit of detection (LOD) for lycopene was 0.63 ug/dL. Using the LOD coding formula (detection limit divided by the square root of two), the calculated value indicating that the serum lycopene results were below the level of detection would be 0.44. After rounding, the value of 0 (zero) was placed in the results field to indicate that the serum lycopene was below 0.63 ug/dL.

MCPSI: Mean cell hemoglobin

See note for HGP.

MEP: Metamyelocyte cells

See note for ANP.

MHP: Mean cell hemoglobin concentration

See note for HGP.

MIP: Microcytosis

See note for ANP.

MLP: Myelocyte cells

See note for ANP.

MOP: Mononuclear number

See note for GRP.

MOPDIF: Monocyte cells

See note for ANP.

MRP: Macrocytosis

See note for ANP.

MVPSI: Mean cell volume

See note for HGP.

OSPSI: Serum osmolality

Results for osmolality were added to the protocol after NHANES III began. This result field is blank-filled for examinees who were examined prior to the start of testing.

PHPBEST: Time of venipuncture

The time of venipuncture is expressed using the 24-hour clock system (military time) in which 01:00 corresponds to 1:00 a.m., 12:00 corresponds to 12 noon, 13:00 corresponds to 1:00 p.m., and 00:00 corresponds to 12 midnight.

PHPCHM2: Within the past four weeks have you received any cancer chemotherapy treatment?

All examinees who indicated at the time of venipuncture that they had received cancer chemotherapy treatment in the past two weeks (later this was changed to four weeks) were excluded from venipuncture. For these examinees, results fields for blood-based analyses are blank-filled.

PHPFAST: Calculated fasting time in hours

The fasting time was calculated using the time of venipuncture and the time the examinee last ate or drank (other than water). This was determined using the snack/drink time and the corresponding day variables. Fasting time is the elapsed interval between the time the examinee last ate or drank and the time of venipuncture.

The following variables were used to calculate this variable: PHPSNTI, PHPSNDA, PHPDRIN, PHPDRTI, PHPDRDA, and PHPBEST. If the examinee drank only water since he/she last ate (PHPDRIN = 2), then the time and day the examinee last ate (PHPSNTI and PHPSNDA) were subtracted from the time and day of the venipuncture (PHPBEST). The difference was the number of hours between the time the examinee last ate and the time of the venipuncture.

If the examinee drank anything other than water (PHPDRIN = 1), then the time and day the examinee last drank (PHPDRTI and PHPDRDA) were subtracted from the time and day of the venipuncture (PHPBEST). The difference was the number of hours between the time the examinee last drank and the time of the venipuncture.

PHPHEMO: Do you have hemophilia?

All examinees who indicated at the time of venipuncture that they had hemophilia, a hereditary blood-clotting disorder, were excluded from the venipuncture. Results for blood analyses were blank-filled.

PHPINSU: Are you currently taking insulin?

See note for G1P and G1PCODE.

PHPLANG: Language of the venipuncture screening questionnaire

Both English and Spanish versions of the venipuncture screening questionnaire were used. The language used depended on the preference of the examinee. Translators, either hired or friends/family members, were available for examinees who spoke neither Spanish nor English.

PKP: Poikilocytosis

See note for ANP.

PLP: Platelet count

See note for GRP.

POP: Polychromatophilia

See note for ANP.

PRP: Promyelocyte cells

See note for ANP.

PXP: Serum transferrin saturation

This value was calculated as $(FEP / TIP) * 100$.

RBP: RBC folate

See note for FOP.

RCP: Red blood cell count

See notes for HGP and GRP.

RUP: Serum rubella antibody

Rubella antibody data are reported both as an optical density index and in International Units. The index was calculated by subtracting the absorbance of the control well from the absorbance of the antigen well (AG-NS) and dividing the difference by the cut-off value. The cut-off value was calculated as the mean AG-NS value of duplicate 10 IU standards. The equation used was:

O.D. index = (AG-NS)/Cut-off value

An O.D. index greater than or equal to one indicates the presence of antibody.

RUPUNIT: Serum rubella antibody (IU)

Rubella antibody data are reported both as an optical density index and in International Units. International Units were calculated based on a standard curve using a regression analysis of duplicate AG-NS values of 10, 40, & 100 IU standards and their squares. An International Unit value greater than or equal to 10 indicates the presence of antibody.

SAP: Serum hepatitis B surface antigen

See note for HBP.

SEP: Serum selenium

Selenium values were measured on two Perkin-Elmer graphite furnace atomic absorption spectrophotometers (model 3030 and model 5100) during the six-year study. Based on a comparability study using linear models, the results generated using the Model 5100 instrument (from 12/07/90 to 1/13/95) were on average 4.3 percent higher than those from the Model 3030 instrument (used from 10/1/88 to 12/06/90). Since the Model 5100 represented more precise measurements, the model 3030 data were adjusted to make them comparable to the Model 5100. Perkin-Elmer Model 5100 Zeeman-corrected graphite furnace atomic absorption spectrophotometer testing began on 12/07/90. All selenium values measured prior to 12/07/90 were adjusted to the AA5100 values. The formula used was:

New value = 16.795 + 0.902 * original value.

SFP: Serum iron

This value was obtained from the standard battery of biochemical assessments. Use of the laboratory test result from the reference method (FEP), rather than the SFP value, is generally recommended. For most analyses of serum iron, the appropriate variable to use will be FEP. The value from the biochemistry profile (SFP) should not be used routinely. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996) for details. Laboratory test results for SFP were added to the protocol after NHANES III began. This result field was blank-filled for examinees who were examined prior to the start of testing.

SGP: Serum glucose

This value was obtained from the standard battery of biochemical assessments. Use of the laboratory test result for plasma glucose

from the reference method (G1P), rather than the SGP value, is generally recommended. For most analyses, the appropriate variable to use will be G1P. The value from the biochemistry profile (SGP) should not be used routinely. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996) for details.

SHP: Spherocytosis

See note for ANP.

SIP: Sickle cells

See note for ANP.

SSP: Serum hepatitis B surface antibody

See note for HBP.

TGP: Serum triglycerides

Serum triglyceride levels were measured regardless of the examinee's fasting status. Mean serum triglycerides and the distribution of serum triglycerides should be estimated only on examinees who did fast at least nine hours, were examined in the morning, and were randomly assigned to the morning fasting sample (WTPFHSD6 > 0). For the purpose of this calculation, the number of hours fasted was rounded to the nearest whole integer. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996) for details.

TOP: Serum toxoplasmosis antibody

The presence and quantity of antibody to *Toxoplasma gondii* in the test sample were determined by comparing the optical density of the test sample to a standard curve. A standard curve was constructed using optical density readings from positive control sera obtained from a kit; these readings were calibrated to WHO Toxo 60 serum and read as International Units (IU/mL). Those test samples exhibiting titer below 7 IU/mL indicated a non-significant level of antibody according to this technique; thus, they were considered to be negative, indicating no infection. Those test samples with results greater than 6 IU/mL were considered to be positive, indicating infection at some undetermined time.

TRP: Serum triglycerides

This value was obtained from the standard battery of biochemical assessments. Use of the laboratory test result from the reference method (TGP), rather than the TRP value, is generally recommended. For most analyses, the appropriate variable to use is TGP. The value from the biochemistry profile (TRP) should not be used routinely. Consult the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996)

for details. Results for TRP were added to the protocol after NHANES III began. This result field was blank-filled for examinees who were examined prior to the start of testing.

TTP: Target cells

See note for ANP.

TXP: Toxic granulation

See note for ANP.

URP: Urinary creatinine

Although the laboratory method detection limit for urinary creatinine is 1 mg/dL, all values below 10 mg/dL were considered "statistically suspect" and were coded as "below level of detection".

VCP: Serum vitamin C

For NHANES III, serum concentrations of vitamin C were measured using a total vitamin C, fully reduced method using high-performance liquid chromatography with electrochemical detection (HPLC-EC) analysis. This method differed from the 2,4-dinitrophenyl hydrazine colorimetric method used in the NHANES II study. A comparison study of the two methods was carried out. Linear regression analysis, by an error in both variables' technique, was used to compare the results obtained by the two methods; values for slope, intercept, and correlation coefficient were 0.881, 0.036, and 0.927, respectively, for 138 singlet analyses.

Serum concentrations obtained by HPLC-EC were lower than those obtained by the 2,4-DNPH method. This difference was expected due to the increased specificity of the HPLC method. Unlike colorimetric methods, HPLC separates uric acid and other potential interferers from ascorbate, thereby increasing accuracy and specificity. The 2,4-DNPH method also measured endogenous diketogulonate, the product of the irreversible oxidation of dehydroascorbic acid. This species was not measured by most HPLC methods and generally was not included in total vitamin C measurements since it has no vitamin C activity. Because the laboratory method differed between NHANES III and NHANES II, the results from the two surveys are not comparable.

Blocks of vitamin C data are missing due to an inadvertent misdilution of the ascorbic acid-serum ratio.

VEP: Serum vitamin E

The vitamin E value of 9999 was confirmed.

VRP: Serum varicella antibody

Varicella antibody data were reported as an optical density index. See note RUP for the index calculation. The equation used was:

$$\text{O.D. index} = (\text{AG-NS}) / \text{Cut-off value}$$

The cut-off value was 0.1. An O.D. index equal to or greater than one indicates the presence of antibody.

VUP: Vacuolated cells

See note for ANP.

WCP: White blood cell count

See note for HGP and GRP.

Appendix 1. Blood and Urine Assessments by Specimen Type and Age Group

| AGE GROUP | | |
|--------------------|-------------------------|-------------------------|
| 1-3 years | 4-5 years | 6-11 years |
| | Whole blood | |
| CBC (1)(5) | CBC (1) (5) | CBC (1) (5) |
| Differential smear | Differential smear | Differential smear |
| Lead (5) | Lead (5) | Lead (5) |
| Protoporphyrin (5) | Protoporphyrin (5) | Protoporphyrin (5) |
| | RBC folate | RBC folate |
| | Glycated hemoglobin (5) | Glycated hemoglobin (5) |
| | Serum | |
| Iron (5) | Iron (5) | Iron (5) |
| TIBC (5) | TIBC (5) | TIBC (5) |
| Ferritin (5) | Ferritin (5) | Ferritin (5) |
| | Folate (5) | Folate (5) |
| | Apolipoprotein AI(4)(5) | Apolipoprotein AI(4)(5) |
| | Apolipoprotein B(4)(5) | Apolipoprotein B(4)(5) |
| | Cholesterol (5) | Cholesterol (5) |
| | HDL/LDL (5) | HDL/LDL (5) |
| | Triglycerides (5) | Triglycerides (5) |
| | Lp(a)(2)(5) | Lp(a)(2)(5) |
| | Cotinine (4) | Cotinine (4) |
| | C-reactive protein (5) | C-reactive protein (5) |
| | Vitamin A (5) | Vitamin A (5) |
| | Carotenes (5) | Carotenes (5) |
| | Retinyl esters (5) | Retinyl esters (5) |
| | Vitamin E (5) | Vitamin E (5) |
| | Vitamin B12 (2) | Vitamin B12 (2) |
| | | Helicobacter pylori (4) |
| | Tetanus | Tetanus |
| | | Vitamin C |
| | | Hepatitis A |

Appendix 1. Blood and Urine Assessments by Specimen Type and Age Group
(continued)

AGE GROUP

1-3 years

4-5 years
Serum (continued)

6-11 years

Hepatitis B/delta
Hepatitis C
Hepatitis E
Rubella (5)
Varicella (5)

Urine

Cadmium
Creatinine
Albumin
Iodine

Appendix 1. Blood and Urine Assessments by Specimen Type and Age Group
(continued)

AGE GROUP

12-19 years

20 years and older

Whole blood

CBC (1)(5)

Differential smear

Lead (5)

Protoporphyrin (5)

Glycated hemoglobin (5)

CBC (1)(5)

Differential smear

Lead (5)

Protoporphyrin (5)

RBC folate

Glycated hemoglobin (5)

Serum

Iron (5)

TIBC (5)

Ferritin (5)

Folate (5)

Apolipoprotein AI(4)(5)

Apolipoprotein B(4)(5)

Cholesterol (5)

HDL/LDL (5)

Triglycerides (5)

Lp(a)(2)(5)

Cotinine (4)

C-reactive protein (5)

Vitamin A (5)

Carotenes (5)

Retinyl esters (5)

Vitamin E (5)

Vitamin B12 (2)

Helicobacter pylori (4)

Tetanus

Vitamin C

Hepatitis A

Hepatitis B/delta

Hepatitis C

Hepatitis E

Rubella (5)

Varicella (5)

Iron (5)

TIBC (5)

Ferritin (5)

Folate (5)

Apolipoprotein AI(4)(5)

Apolipoprotein B(4)(5)

Cholesterol (5)

HDL/LDL (5)

Triglycerides (5)

Lp(a)(2)(5)

Cotinine (4)

C-reactive protein (5)

Rheumatoid factor (60+)

Vitamin A (5)

Carotenes (5)

Retinyl esters (5)

Vitamin E (5)

Vitamin B12 (2)

Tetanus

Vitamin C

Hepatitis A

Hepatitis B/delta

Hepatitis C

Hepatitis E

Rubella (5)

Varicella (5)

Appendix 1. Blood and Urine Assessments by Specimen Type and Age Group
(continued)

AGE GROUP

12-19 years

20 years and older

Serum

| | |
|------------------------------|-----------------------------------|
| Diphtheria | Diphtheria |
| Herpes simplex I and II | Herpes simplex I and II |
| HIV I (ages 18+)(3)(5) | HIV I (ages 18+)(3)(5) |
| Toxoplasmosis (5) | Toxoplasmosis (5) |
| Vitamin D (OHD) | Vitamin D (OHD) |
| Total/normalized calcium | Total/normalized calcium |
| Selenium (5) | Selenium (5) |
| Thyroxine (T4) | Thyroxine (T4) |
| Thyroid-stimulating hormone | Thyroid-stimulating hormone |
| Antithyroglobulin antibodies | Antithyroglobulin antibodies |
| Antimicrosomal antibodies | Antimicrosomal antibodies |
| | FSH/LH (females aged 35-60 years) |
| | Insulin (6) |
| | C-peptide (6) |
| Biochemistry profile (5) | Biochemistry profile (5) |
| Bicarbonate | Bicarbonate |
| Blood urea nitrogen | Blood urea nitrogen |
| Total bilirubin | Total bilirubin |
| Alkaline phosphatase | Alkaline phosphatase |
| Cholesterol | Cholesterol |
| AST | AST |
| ALT | ALT |
| LDH | LDH |
| GGT | GGT |
| Total protein | Total protein |
| Albumin | Albumin |
| Creatinine | Creatinine |
| Glucose | Glucose |
| Calcium | Calcium |
| Chloride | Chloride |
| Uric acid | Uric acid |
| Phosphorus | Phosphorus |
| Sodium | Sodium |
| Potassium | Potassium |
| Triglycerides | Triglycerides |
| Globulin | Globulin |
| Iron | Iron |
| Osmolality | Osmolality |

Appendix 1. Blood and Urine Assessments by Specimen Type and Age Group
(continued)

AGE GROUP

12-19 years

20 years and older

Plasma

Glucose (examinees aged 20-39 years and 75 years and older)
OGTT (examinees aged 40-74 years)
Fibrinogen (examinees aged 40 years and older)(5)

Urine

Cadmium
Creatinine
Albumin
Iodine
Urine drug (ages 18 years and over)(2)(3)
Cocaine
Opiates
Phencyclidine
Amphetamines
Marijuana

Cadmium
Creatinine
Albumin
Iodine
Urine drug (examinees aged 18 years and over)(2)(3)
Cocaine
Opiates
Phencyclidine
Amphetamines
Marijuana
Pregnancy test (females aged 20-59 years)

White Cells

Storage/banking (5)

Storage/banking (5)

(1) Includes hematocrit, hemoglobin, red, white and platelet cell counts, mean cell volume, mean cell hemoglobin, mean cell hemoglobin concentration, red cell distribution width, platelet distribution width, mean platelet volume, and 3-cell differential

(2) Phase 2 only

(3) Anonymous

(4) Phase 1 only

(5) Home examination also

(6) In phase 2, also from second venipuncture for examinees aged 40-74 years

Appendix 2. Laboratory Test Detection Limits

| Test | Detection limit |
|------------------------------------|-------------------|
| Albumin (urine) | 0.5 ug/mL |
| Alpha carotene | 0 ug/dL |
| Antimicrosomal antibody (AMA) | 0.5 U/mL |
| Antithyroglobulin antibody (ATA) | 1.0 U/mL |
| Beta carotene | 0.67 ug/dL |
| Beta cryptoxanthin | 0 ug/dL |
| C-peptide | 0.03 pmol/mL |
| C-reactive protein | 0.3 mg/dL |
| Cadmium (urine) | 0.01 ng/mL |
| Cotinine | 0.05 ng/mL |
| Creatinine (urine) | 1 mg/dL |
| Erythrocyte protoporphyrin | 2.5 ug/dL RBC |
| Ferritin | 3 ng/mL |
| Folate (serum) | 0.2 ng/mL |
| Follicle stimulating hormone (FSH) | 0.15 IU/L |
| Glucose | 2 mg/dL |
| Glycated hemoglobin | 0 % |
| Hematology parameters | |
| Granulocyte | 0 % |
| Granulocyte (1) | 0 number |
| Hematocrit | 0 % |
| Hemoglobin | 0 g/dL |
| Lymphocyte | 0 % |
| Lymphocyte (1) | 0 number |
| Mean cell hemoglobin | 0 pg |
| Mean cell hemoglobin concentration | 0 g/dL |
| Monocyte | 0 % |
| Monocyte (1) | 0 number |
| Platelet count (1) | 0 |
| Platelet distribution width | 0 % |
| Red blood cell count (RBC) (1) | 0 |
| Red blood cell distribution width | 0 % |
| White blood cell count (WBC) (1) | 0 |
| Hepatitis profile | Qualitative tests |
| Herpes | Qualitative tests |
| High density lipoprotein (HDL) | 10 mg/dL |
| Human immunodeficiency virus (HIV) | Qualitative tests |
| Insulin | 2.5 uU/mL |
| Iodine (urine) | 0.2 ug/dL |
| Iron | 3.0 ug/dL |
| Lead | 1 ug/dL |
| Lipoprotein(a) | 0 mg/dL |
| Lutein/zeaxanthin | 0.43 ug/dL |

Appendix 2. Laboratory Test Detection Limits (continued)

| Test | Detection limit |
|------------------------------------|-------------------|
| Luteinizing hormone (LH) | 0.15 IU/L |
| Lycopene | 0.63 ug/dL |
| Normalized calcium | 0.5 mmol/L |
| RBC folate | 4.4 ng/mL |
| Retinyl esters | 0 ug/dL |
| Rheumatoid factor | Qualitative tests |
| Rubella | 0 IU |
| Selenium | 8 ng/mL |
| Tetanus | 0 U/mL |
| Thyroid stimulating hormone (TSH) | 0.01 mU/mL |
| Thyroxine (T4) | 1.0 ug/dL |
| Total iron binding capacity (TIBC) | 9 ug/dL |
| Total cholesterol | 10 mg/dL |
| Total calcium | 1.5 mmol/L |
| Toxoplasmosis | 0 IU |
| Triglycerides | 10 mg/dL |
| Varicella | 0 |
| Vitamin B12 | 20 pg/mL |
| Vitamin E | 20 ug/dL |
| Vitamin C | 0 mg/dL |
| Vitamin A | 0.5 ug/dL |
| Vitamin D | 5.0 ng/mL |

(1) Units for white blood cell count, red blood cell count, platelet count, lymphocyte number, granulocyte number, and mononuclear number are referenced in the Manual for Medical Technicians p. 5-1 (U.S. DHHS, 1996).

Note: Lower detection limits for analytes included in the general "biochemistry profile" are found in the Laboratory Procedures Used for NHANES III (U.S. DHHS, 1996).

Appendix 3. NHANES III SI Table

| Test (1) | NHANES Unit | NHANES Format | Conversion Factor | SI Unit | SI Format |
|--------------------------------|-------------|---------------|-------------------|---------|-----------|
| Alanine aminotransferase(2) | N/A | N/A | N/A | U/L | XXX |
| Albumin (serum) (2) | g/dL | X.X | 10 | g/L | XX |
| Albumin (urine) | ug/mL | XXXXX.XX | N/A | N/A | N/A |
| Alkaline phosphatase (2) | N/A | N/A | N/A | U/L | XXX |
| Alpha carotene | ug/dL | XXX | 0.01863 | umol/L | X.XX |
| Antimicrosomal antibody | N/A | N/A | N/A | N/A | N/A |
| Antithyroglobulin antibody | N/A | N/A | N/A | N/A | N/A |
| Apolipoprotein AI | mg/dL | XXX | 0.01 | g/L | X.XX |
| Apolipoprotein B | mg/dL | XXX | 0.01 | g/L | X.XX |
| Aspartate aminotransferase (2) | N/A | N/A | N/A | U/L | XXX |
| Beta carotene | ug/dL | XXX | 0.01863 | umol/L | XX.XX |
| Beta cryptoxanthin | ug/dL | XXX | 0.01809 | umol/L | X.XX |
| Bicarbonate (2) | N/A | N/A | N/A | mmol/L | XX |
| Bilirubin (total)(2) | mg/dL | XX.X | 17.1 | umol/L | XXX.XX |
| Blood urea nitrogen (2) | mg/dL | XXX | 0.357 | mmol/L | XX.XX |
| C-peptide | pmol/mL | XX.XXX | 1 | nmol/L | XX.XXX |
| C-reactive protein | N/A | N/A | N/A | N/A | N/A |
| Cadmium (urine) | ng/mL | XX.XX | 8.897 | nmol/L | XXX.XX |
| Calcium (total) | N/A | N/A | N/A | mmol/L | X.XX |
| Calcium (normalized) | N/A | N/A | N/A | mmol/L | X.XX |
| Calcium (2) | mg/dL | XX.X | 0.25 | mmol/L | X.XXX |
| Chloride (2) | N/A | N/A | N/A | mmol/L | XXX.X |
| Cholesterol | mg/dL | XXX | 0.02586 | mmol/L | XX.XX |
| Cholesterol (HDL) | mg/dL | XXX | 0.02586 | mmol/L | X.XX |
| Cholesterol (LDL) | mg/dL | XXX | 0.02586 | mmol/L | X.XX |
| Cholesterol (2) | mg/dL | XXX | 0.02586 | mmol/L | XX.XXX |
| Cotinine | ng/mL | XXXX.XXX | N/A | N/A | N/A |
| Creatinine (2) | mg/dL | XX.X | 88.4 | umol/L | XXXX.X |
| Creatinine (urine) | mg/dL | XXX.X | 0.0884 | mmol/L | XX.X |
| Diphtheria | N/A | N/A | N/A | N/A | N/A |
| Ferritin | ng/mL | XXXX | 1 | ug/L | XXXX |
| Fibrinogen | mg/dL | XXX | 0.01 | g/L | X.XX |
| Folate | ng/mL | XXX.X | 2.266 | nmol/L | XXX.X |
| Folate (RBC) | ng/mL | XXXX | 2.266 | nmol/L | XXXX.X |
| Follicle-stimulating hormone | N/A | N/A | N/A | IU/L | XXX.X |
| GGT (2) | N/A | N/A | N/A | U/L | XXXX |

Appendix 3. NHANES III SI Table

| Test (1) | NHANES Unit | NHANES Format | Conversion Factor | SI Unit | SI Format |
|--|----------------|------------------|----------------------|------------|--------------|
| Globulin (2) | g/dL | X.X | 10 | g/L | XX |
| Glucose (2) | mg/dL | XXX | 0.05551 | mmol/L | XX.XX |
| Glucose (plasma) | mg/dL | XXX.X | 0.05551 | mmol/L | XX.XXX |
| Glycated hemoglobin | % | XX.X | N/A | N/A | N/A |
| Helicobacter pylori | N/A | N/A | N/A | N/A | N/A |
| Hematocrit | % | XX.XX | 0.01 | L/L=1 | 0.XXX |
| Hemoglobin | g/dL | XX.XX | 10 | g/L | XXX.X |
| Hepatitis A virus | N/A | N/A | N/A | N/A | N/A |
| Hepatitis B core antibody (anti-HBc) | N/A | N/A | N/A | N/A | N/A |
| Hepatitis B surface antigen (HbsAg) | N/A | N/A | N/A | N/A | N/A |
| Hepatitis C virus | N/A | N/A | N/A | N/A | N/A |
| Hepatitis D virus | N/A | N/A | N/A | N/A | N/A |
| Hepatitis B surface antibody (anti-HBs) | N/A | N/A | N/A | N/A | N/A |
| Herpes I & II | N/A | N/A | N/A | N/A | N/A |
| Homocysteine | N/A | N/A | N/A | umol/L | XX.X |
| Human immuno- deficiency virus | N/A | N/A | N/A | N/A | N/A |
| Insulin | uU/mL | XXX.XX | 6.0 | pmol/L | XXX.XX |
| Iodine (urine) | ug/dL | XXX.X | N/A | N/A | N/A |
| Iron | ug/dL | XXX | 0.1791 | umol/L | XX.XX |
| Iron (2) | ug/dL | XXX | 0.1791 | umol/L | XX.X |
| LDH (2) | N/A | N/A | N/A | U/L | XXX |
| Latex antibody | IU/mL | XXXX.XX | N/A | N/A | N/A |
| Lead | ug/dL | XX.X | 0.04826 | umol/L | X.XXX |
| Lipoprotein(a) | mg/dL | XXX | 0.01 | g/L | X.XX |
| Lutein/zeaxanthin | ug/dL | XXX | 0.01758 | umol/L | X.XX |
| Luteinizing hormone | N/A | N/A | N/A | IU/L | XX.X |
| Lycopene | ug/dL | XXX | 0.01863 | umol/L | X.XX |
| Mean cell hemoglobin | N/A | N/A | N/A | pg | XX.XX |
| Mean cell volume | N/A | N/A | N/A | fL | XXX.XX |
| Mean cell hemoglobin concentration | g/dL | XX.XX | 10 | g/L | XXX.X |
| Mean platelet volume | N/A | N/A | N/A | fL | XX.XX |
| Methylmalonic acid | ug/dL | N/A | 0.085 | umol/L | N/A |

Appendix 3. NHANES III SI Table (continued)

| Test (1) | NHANES Unit | NHANES Format | Conversion Factor | SI Unit | SI Format |
|--------------------------------------|----------------|------------------|----------------------|------------|--------------|
| Osmolality (2) | N/A | N/A | N/A | mmol/kg | XXX |
| Phosphorus (2) | mg/dL | XX.X | 0.3229 | mmol/L | X.XXX |
| Platelet count (3) | N/A | XXX.X | 1 | N/A | XXX.X |
| Potassium (2) | N/A | N/A | N/A | mmol/L | X.XX |
| Protein (total)(2) | g/dL | XX.X | 10 | g/L | XXX |
| Protoporphyrin | ug/dL | XXXX | 0.0178 | umol/L | XX.XX |
| Red blood cell distribution width | % | XX.XX | 0.01 | fraction | X.XXXX |
| Red blood cell count (3) | N/A | X.XX | 1 | N/A | X.XX |
| Retinyl esters | ug/dL | XXX | 0.03491 | umol/L | X.XX |
| Rheumatoid factor | N/A | N/A | N/A | N/A | N/A |
| Rubella | N/A | N/A | N/A | N/A | N/A |
| Selenium | ng/mL | XXX | 0.0127 | nmol/L | X.XX |
| Sodium (2) | N/A | N/A | N/A | mmol/L | XXX.X |
| Tetanus | U/mL | N/A | N/A | N/A | N/A |
| Thyroid stimulating hormone | uU/mL | XXX.XX | 1 | mU/L | XXX.XX |
| Thyroxine | ug/dL | XX.X | 12.87 | nmol/L | XXX.X |
| Total iron binding capacity | ug/dL | XXX | 0.1791 | umol/L | XXX.XX |
| Toxoplasmosis | N/A | N/A | N/A | N/A | N/A |
| Triglycerides | mg/dL | XXXX | 0.01129 | mmol/L | XX.XX |
| Triglycerides (2) | mg/dL | XXXX | 0.01129 | mmol/L | XX.XXX |
| Uric acid (2) | mg/dL | XX.X | 59.48 | umol/L | XXX.X |
| Varicella | N/A | N/A | N/A | N/A | N/A |
| Vitamin A | ug/dL | XXX | 0.03491 | umol/L | X.XX |
| Vitamin B12 | pg/mL | XXXXXX | 0.7378 | pmol/L | XXXXXX.XX |
| Vitamin C | mg/dL | X.XX | 56.78 | mmol/L | XXX.XX |
| Vitamin D | ng/mL | XXX.X | 2.496 | nmol/L | XXX.X |
| Vitamin E | ug/dL | XXXX | 0.02322 | umol/L | XXX.XX |
| White blood cell count (3) | N/A | XX.XX | 1 | N/A | XX.XX |

(1) Results are based on a serum sample unless otherwise noted.

(2) Biochemistry profile

(3) Units for white blood cell count, red blood cell count, platelet count, lymphocyte number, granulocyte number, and mononuclear number are referenced in the Manual for Medical Technicians p. 5-1 (U.S. DHHS, 1996).

References

- Albers JJ, Marcovina SM. Standardization of apolipoprotein B and A1 measurements. *Clin Chem* 35:1357-61. 1989.
- Bachorik PS, Lovejoy K, Carroll MD, Johnson CL, Albers JL, Marcovina SM. Measurement of apolipoprotein A1 and B during the Health and Nutrition Examination Survey (NHANES III). *Clin Chem* 40(110):1915-1920. 1994.
- Bull BS, Rittenbach JD. A proposed reference hematocrit derived from multiple MCHC determinations via haemoglobin measurements. *Clin Lab Haematol* 12 (suppl 1):43-53. 1990.
- Lewis SA, Hardison NW, Veillon C. Comparison of isotope dilution mass spectrometry and graphite furnace atomic absorption spectrometry with Zeeman background correction for determination of plasma selenium. *Analytical Chemistry* 58:1272-82. 1986.
- Marcovina SM, Albers JJ, Dati F, Ledue TB, Richie RF. International Federation of Clinical Chemistry standardization project for measurements of apolipoprotein a1 and b. *Clin Chem* 37:1676-82. 1991.
- National Center for Health Statistics. Plan and operation of the Third National Health and Nutrition Examination Survey, 1988-94. *Vital Health Stat* 1(32). Hyattsville, Md.: NCHS. 1994.
- National Committee for Clinical Laboratory Standards. Procedure for determining packed cell volume by the microhematocrit method -- second edition: approved standard. NCCLS document H7-32. Wayne, PA: NCCLS. 1993.
- U.S. Department of Health and Human Services (DHHS). National Center for Health Statistics. NHANES III reference manuals and reports (CD-ROM). Hyattsville, Md.: Centers for Disease Control and Prevention, 1996. Available from National Technical Information Service (NTIS), Springfield, Va. (Acrobat .PDF format; includes access software: Adobe Systems Inc. Acrobat Reader 2.1).
- World Health Organization. Diabetes Mellitus: Report of a WHO study group, WHO Technical Report Series 727. Geneva, Switzerland: WHO. 1995.