
Number of Kentucky youth projected to die prematurely from their smoking: 87,902

#### Adult Tobacco Use in Kentucky

Current Cigarette Smoking Among Adults Aged 18 and Older, 1999



## Youth Tobacco Use

GRADES 6–8			GRADES 9–12	
Current Cigarette		Current Any	Current Cigarette	Current Any
Smoking		Tobacco Use	Smoking	Tobacco Use
National*	9.2%	12.8%	28.5%	34.8%
KY†	21.5%	28.3%	37.4%	46.2%
Boys†	21.9%	33.0%	35.6%	50.2%
Girls†	21.1%	23.2%	39.0%	41.8%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. \*Source: National Youth Tobacco Survey, 1999.

†Source: Kentucky Youth Tobacco Survey, 2000.


## Health Impact and Costs

Average Annual Deaths Related		Average Annual Years of	Medical Costs	
to Smoking, 1990–1994		Potential Life Lost,* 1990–1994	Related to Smoking, 1993	
Overall Men Women Death Rate 4 Rank (No. 1 is lowest	7,953 5,357 2,596 44/100,000 50 c death rate)	112,695 years or an average of 14.2 years for each death due to smoking. *Calculated to life expectancy	Ambulatory Hospital Nursing Home† Drug Other <b>Total</b>	\$158,450,000 \$276,560,000 \$197,870,000 \$55,890,000 \$109,470,000 <b>\$798,230,000</b>

Smoking-attributable Medicaid expenditures, Kentucky, fiscal year 1993: \$200,740,000

†Preliminary estimates

# Lung Cancer Death Rate\*



\*1997 deaths per 100,000 population, adjusted to the 1970 total U.S. population.

#### Scheduled 2001 settlement payment to Kentucky: \$121,936,632.68

### **Tobacco Control Funding, 2001**

At 15% of the CDC Best Practices recommended lower estimate funding level, Kentucky ranks 36th (No. 1 is the highest) for tobacco control funding. Kentucky ranks 50th (No. 1 is the low-est death rate) for average annual deaths related to smoking.

FY01 Amount	Funding Cycle
\$2,527,000	7/00–6/01
\$0	
\$0	
\$2,527,000	
\$1,128,413	6/00–5/01
n/a	
\$0	
n/a	
\$1,128,413	
\$3,655,413 \$0.90	
	\$2,527,000 \$0 \$0 \$2,527,000 \$1,128,413 n/a \$0 n/a \$1,128,413 \$3,655,413

An additional \$2,500,000 was appropriated to the Kentucky Agency for Substance Abuse Policy to implement substance abuse treatment services, including smoking cessation.

## Funding as a Percentage of CDC Best Practices Recommendations

	Recommended for Total Program Annual Cost	Recommended Per Capita Funding Level	PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS
Lower Estimate	\$25,090,000	\$6.42	15%
Upper Estimate	\$69,895,000	\$17.88	5%

#### **Excise Tax**

Cigarette tax per pack
Number of packages of cigarettes sold and taxed, per capita, 1999
Federal and state taxes as a percentage of retail price  11.1%    Annual gross tax revenue from cigarettes  \$19,384,000
Smokeless tobacco tax