
Number of Oklahoma youth projected to die prematurely from their smoking: 63,837

Adult Tobacco Use in Oklahoma

Current Cigarette Smoking Among Adults Aged 18 and Older, 1999



Youth Tobacco Use

GRADES 6–8			GRADES 9–12	
Current Cigarette		Current Any	Current Cigarette	Current Any
Smoking		Tobacco Use	Smoking	Tobacco Use
National*	9.2%	12.8%	28.5%	34.8%
OK†	16.9%	21.0%	33.0%	42.0%
Boys†	17.9%	23.8%	31.8%	47.3%
Girls†	15.8%	17.8%	34.0%	36.4%

Current Cigarette Smoking = smoked cigarettes on ≥1 of the 30 days preceding the survey.

 $Current\ Any\ Tobacco\ Use = current\ use\ of\ cigarettes\ or\ smokeless\ tobacco\ or\ pipes\ or\ bidis\ or\ cigars\ or\ kreteks\ on\ \ge 1\ of\ the\ 30\ days\ preceding\ the\ survey.$


Health Impact and Costs

AVERAGE ANNUAL DEATHS RELATED	AVERAGE ANNUAL YEARS OF	MEDICAL COSTS	
TO SMOKING, 1990–1994	POTENTIAL LIFE LOST,* 1990–1994	RELATED TO SMOKING, 1993	
Overall 6,255 Men 4,150 Women 2,106 Death Rate 387/100,000 Rank 42 (No. 1 is lowest death rate)	85,650 years or an average of 13.7 years for each death due to smoking. *Calculated to life expectancy	Ambulatory \$118,780,000 Hospital \$180,130,000 Nursing Home† \$181,940,000 Drug \$39,550,000 Other \$93,050,000 Total \$613,450,000	

Smoking-attributable Medicaid expenditures, Oklahoma, fiscal year 1993: \$80,105,000

†Preliminary estimates

Lung Cancer Death Rate*



^{*1997} deaths per 100,000 population, adjusted to the 1970 total U.S. population.



^{*}Source: National Youth Tobacco Survey, 1999.

[†]Source: Oklahoma Youth Tobacco Survey, 1999.

Scheduled 2001 settlement payment to Oklahoma: \$71,738,602.00

Tobacco Control Funding, 2001

At 18% of the CDC Best Practices recommended lower estimate funding level, Oklahoma ranks 35th (No. 1 is the highest) for tobacco control funding. Oklahoma ranks 42nd (No. 1 is the lowest death rate) for average annual deaths related to smoking.

Funding Source	FY01 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$1,833,333	7/00–6/011
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$1,833,333	
Federal—CDC Office on Smoking and Health	\$1,299,907	6/00–5/01
Federal—SAMHSA	n/a	
Non-Government Source—American Legacy Foundation	\$750,000	10/00-9/01
Non-Government Source—RWJF/AMA	n/a	
Subtotal: Federal/National Sources	\$2,049,907	
Total	\$3,883,240	
Per Capita Funding	\$1.13	

Funding as a Percentage of CDC Best Practices Recommendations

	RECOMMENDED FOR TOTAL PROGRAM ANNUAL COST	RECOMMENDED PER CAPITA FUNDING LEVEL	PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS
Lower Estimate	\$21,825,000	\$6.58	18%
Upper Estimate	\$56,310,000	\$16.98	7%

Excise Tax

Cigarette tax per pack
Number of packages of cigarettes sold and taxed, per capita, 1999
Federal and state taxes as a percentage of retail price
Smokeless tobacco tax