


Number of Washington youth projected to die prematurely from their smoking: 107,799

Adult Tobacco Use in Washington

Current Cigarette Smoking Among Adults Aged 18 and Older, 1999



Youth Tobacco Use

| GRADES 6-8 | | GRADES 9-12 | |
|---------------------------------------|-------------------------|---------------------------|-------------------------|
| Current Cigarette Smoking | Current Any Tobacco Use | Current Cigarette Smoking | Current Any Tobacco Use |
| National 9.2% | 12.8% | 28.5% | 34.8% |
| State-specific data are not available | | | |

Current Cigarette Smoking = smoked cigarettes on ≥1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥1 of the 30 days preceding the survey.
Source: National Youth Tobacco Survey, 1999.


Health Impact and Costs

| AVERAGE ANNUAL DEATHS RELATED TO SMOKING, 1990-1994 | | AVERAGE ANNUAL YEARS OF POTENTIAL LIFE LOST,* 1990-1994 | MEDICAL COSTS RELATED TO SMOKING, 1993 | |
|---|-------------|--|--|------------------------|
| Overall | 7,892 | 102,769 years or an average of 13.0 years for each death due to smoking. | Ambulatory | \$281,590,000 |
| Men | 4,939 | | Hospital | \$335,380,000 |
| Women | 2,953 | | Nursing Home† | \$316,750,000 |
| Death Rate | 351/100,000 | | Drug | \$62,740,000 |
| Rank | 26 | | Other | \$158,820,000 |
| (No. 1 is lowest death rate) | | *Calculated to life expectancy | Total | \$1,155,280,000 |

Smoking-attributable Medicaid expenditures, Washington, fiscal year 1993: \$237,159,000

†Preliminary estimates

Lung Cancer Death Rate*



*1997 deaths per 100,000 population, adjusted to the 1970 total U.S. population.



Scheduled 2001 settlement payment to Washington: \$142,160,616.27

Tobacco Control Funding, 2001

At 54% of the CDC Best Practices recommended lower estimate funding level, Washington ranks 17th (No. 1 is the highest) for tobacco control funding. Washington ranks 26th (No. 1 is the lowest death rate) for average annual deaths related to smoking.

| <i>FUNDING SOURCE</i> | <i>FY01 AMOUNT</i> | <i>FUNDING CYCLE</i> |
|--|--------------------------------------|----------------------|
| State Appropriation—Settlement (Tobacco Only) | \$15,000,000 | 7/00–6/01 |
| State Appropriation—Excise Tax Revenue | \$0 | |
| State Appropriation—Other | \$902,000 | |
| Subtotal: State Appropriation | \$15,902,000 | |
| Federal—CDC Office on Smoking and Health | \$1,424,995 | 6/00–5/01 |
| Federal—SAMHSA | n/a | |
| Non-Government Source—American Legacy Foundation | \$801,290 | 9/00–8/01 |
| Non-Government Source—RWJF/AMA | n/a | |
| Subtotal: Federal/National Sources | \$2,226,285 | |
| Total Per Capita Funding | \$18,128,285 \$3.08 | |

Funding as a Percentage of CDC Best Practices Recommendations

| | <i>RECOMMENDED FOR TOTAL PROGRAM ANNUAL COST</i> | <i>RECOMMENDED PER CAPITA FUNDING LEVEL</i> | <i>PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS</i> |
|----------------|--|---|--|
| Lower Estimate | \$33,341,000 | \$5.94 | 54% |
| Upper Estimate | \$89,381,000 | \$15.93 | 20% |

Excise Tax

| | |
|---|-------------------------------|
| Cigarette tax per pack | 82.5¢ |
| Rank = 5 (No. 1 is highest tax) | |
| Number of packages of cigarettes sold and taxed, per capita, 1999 | .55.1 |
| Rank = 5 (No. 1 is lowest cigarette sales) | |
| Federal and state taxes as a percentage of retail price | 29.6% |
| Annual gross tax revenue from cigarettes | \$253,428,006 |
| Smokeless tobacco tax | .75% of wholesale sales price |