## DRIVING?

## Check the Medicine Label... To Make Sure You're Able

## When Using This Product...

- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Drug Facts the grant construction of pain releases are resized. Acetanian of the pain releases a prescription monoaming of the pain respectively. The construction of constructions of painting of pa one particles of emotional condenses of parking of the prescription of parking of parking of parking of parking of parking of the prescription of the parking of the parkin to depression, psychiatric, or emotional conditions of oxidiant of the depression of the MAOI drug. If you do not know if your prescription of the depression of the product containing acetaminophen. Purpose and any other product containing acetaminophen Julia doctor before use if you nave

Julia docto heard disease I right based as empty sema, asshma, or chronic bronchits ika doctor or pharmacist before use if you are taking sedatives or tranquistion When using this product • do not use more than directed • avoid alcoholic driks Men using this product of the second of the second according to the second of the second according to the second of the second o be careful when driving a motor vehicle or operating machinery seand ask a destockly in chemical distriness, or sleeplessness occurs Simplified for more than 7 days, comes back, or occurs with a lever, rash, or persimplified for 10 days (nain) or for 3 days (fever) symptoms do not improve for 10 days comes back, or occurs warrane in the improve for 10 days (pain) or for 3 days (lever) and occurs with a lever. Signature of the state of the s Regular phase could be signs of a serious condition

The condition of the ach of children, ask a health care professional belove containing asks a health care professional belove contained from a hour of the professional belove contained to hold asks, prompt medical attention is critical for adults and signs or symptoms. Directions

Lake every 4 to 6 hours was to exceed 4 Packets in 24 hours or as designed and over described contents of the state of the water, sip while hot \* Children under 12 years of age: consult a doctor Sweeten to taste if desired.

Microwave heating directions \* add contents of one packet and 6 oz. cool water to a microwave safe out and str. briskly. Microwave on high 1.1/2 minutes or until hot Do no ours or a solid by a doctor.

as of one packet into 6 oz. hot Microwave heating directions \* add contents of one packet and 6 oz. coor water to a aler or overheat, and remember to strilliquid between reheatings



