

1 AHFS Category: 80:08

2 **Tetanus and Diphtheria Toxoids Adsorbed**

3 **For Adult Use**

4 **DECAVAC™**



5

6 **DESCRIPTION**

7 DECAVAC™, Tetanus and Diphtheria Toxoids Adsorbed, For Adult Use (Td), manufactured by
8 Aventis Pasteur Inc. for intramuscular injection, is a sterile suspension of alum-precipitated
9 (aluminum potassium sulfate) toxoids in an isotonic sodium chloride solution. The vaccine, after
10 shaking, is a turbid liquid, whitish-gray in color.

11

12 *Corynebacterium diphtheriae* cultures are grown in a modified Mueller and Miller medium.¹

13 *Clostridium tetani* cultures are grown in a peptone-based medium containing a bovine
14 extract. The meat used in this medium is US sourced. Tetanus and diphtheria toxins

15 produced during the growth of the cultures are detoxified with formaldehyde. The detoxified
16 materials are then separately purified by serial ammonium sulfate fractionation and diafiltration.

17 DECAVAC vaccine is supplied in a unit dose 0.5 mL preservative-free prefilled syringe

18 presentation, but contains a trace amount of thimerosal [(mercury derivative), ($\leq 0.3 \mu\text{g}$

19 mercury/dose)] from the manufacturing process.

20

1 Each 0.5 mL dose is formulated to contain 5 Lf of tetanus toxoid, 2 Lf of diphtheria toxoid, and
2 not more than 0.28 mg of aluminum by assay. The tetanus and diphtheria toxoids induce at least 2
3 units and 0.5 units of antitoxin per mL of serum, respectively, in the guinea pig potency test.
4

5 **CLINICAL PHARMACOLOGY**

6 In the United States (US), immunization against tetanus and diphtheria became widespread in the
7 late 1940s, and resulted in a striking decrease in the incidence of morbidity and mortality from
8 these diseases.
9

10 **TETANUS**

11 Tetanus is an acute and often fatal disease caused by an extremely potent neurotoxin produced by
12 *C tetani*. The toxin causes neuromuscular dysfunction, with rigidity and spasms of skeletal
13 muscles. The muscle spasms usually involve the jaw (lockjaw) and neck and then become
14 generalized.
15

16 The occurrence of tetanus disease in the US decreased steadily from 560 reported cases in 1947 to
17 an average of 43 cases reported annually during 1998-2000. Among patients with known outcome,
18 the case-fatality ratio during 1998-2000 was 18%, 5 times lower than the case-fatality ratio of 91%
19 reported in 1947. In the mid to late 1990s, the age distribution of reported tetanus cases among
20 adults shifted to a younger age group. Among cases reported during 1998-2000, 9% were <20
21 years of age, 55% were 20-59 years of age, and 36% were ≥60 years of age. In previous decades,
22 most cases were among persons ≥60 years of age. Adults ≥60 years of age continue to have the

1 highest rates of tetanus and tetanus-related deaths. The majority of tetanus cases during 1998-2000
2 occurred among persons who were not appropriately vaccinated against tetanus or who had an
3 unknown vaccination history.²

4

5 Neonatal tetanus occurs among infants born under unhygienic conditions to inadequately
6 vaccinated mothers. Vaccinated mothers confer protection to their infants through transplacental
7 transfer of maternal antibody.³ From 1998 through 2000, one case of neonatal tetanus was reported
8 in the US.²

9

10 Spores of *C tetani* are ubiquitous. Serologic tests indicate that naturally acquired immunity to
11 tetanus toxin does not occur in the US. Thus, universal primary vaccination, with subsequent
12 maintenance of adequate antitoxin levels by means of appropriately timed boosters, is necessary to
13 protect all age-groups.³ Following adequate immunization with tetanus toxoid, it is thought that
14 protection persists for at least 10 years. Protection against disease is due to the development of
15 neutralizing antibodies to tetanus toxin. A serum tetanus antitoxin level of at least 0.01 IU/mL,
16 measured by neutralization assays, is considered the minimum protective level.^{4,5} More recently, a
17 level ≥ 0.1 to 0.2 IU/mL has been considered as protective.⁶

18

19 DIPHTHERIA

20 *Corynebacterium diphtheriae* may cause both localized and generalized disease. The systemic
21 intoxication is caused by diphtheria exotoxin, an extracellular protein metabolite of toxigenic
22 strains of *C diphtheriae*. Both toxigenic and nontoxigenic strains of *C diphtheriae* can cause

1 disease, but only strains that produce toxin cause myocarditis and neuritis. Toxigenic strains are
2 more often associated with severe or fatal respiratory infections than with cutaneous infections.³

3

4 Prior to the widespread use of diphtheria toxoid in the late 1940s, diphtheria disease was common
5 in the US. More than 200,000 cases, primarily among children, were reported in 1921.
6 Approximately 5% - 10% of cases were fatal; the highest case-fatality rates were in the very
7 young and the elderly. More recently, reported cases of diphtheria of all types declined from 306
8 in 1975 to 59 in 1979; most were cutaneous diphtheria reported from a single state. After 1979,
9 cutaneous diphtheria was no longer reportable.³ From 1980 through 2000, 51 cases of diphtheria
10 were reported in the US, an average of 2 or 3 cases per year. Only one case was reported each year
11 in 1998, 1999, and 2000. The case-fatality rate for diphtheria has changed very little since the
12 1950s. Of 49 reported cases with known age during the period 1980-2000, 55% were in persons
13 =20 years of age; 43% were among persons =40 years of age.⁷ Most cases have occurred in
14 unimmunized or inadequately immunized persons. Although diphtheria disease is rare in the US, it
15 appears that *C diphtheriae* continues to circulate in areas of the country with previously endemic
16 diphtheria.⁷

17

18 Diphtheria continues to occur in other parts of the world. A major epidemic of diphtheria occurred
19 in the newly independent states of the former Soviet Union beginning in 1990. This epidemic
20 resulted in approximately 150,000 cases and 5,000 deaths during the years 1990-1998.⁸ This
21 outbreak is believed to be due to several factors, including a lack of routine immunization of
22 adults in these countries.⁹

1

2 Complete immunization significantly reduces the risk of developing diphtheria and immunized
3 persons who develop disease have milder illness. Protection against disease is due to the
4 development of neutralizing antibodies to diphtheria toxin. A serum antitoxin level of
5 0.01 IU/mL is the lowest level giving some degree of protection. Antitoxin levels of at least
6 0.1 IU/mL are generally regarded as protective.⁵ Following adequate immunization with diphtheria
7 toxoid, it is thought that protection persists for ≥ 10 years.¹⁰ Immunization with diphtheria toxoid
8 does not, however, eliminate carriage of *C diphtheriae* in the pharynx, nose, or on the skin.

9

10 EFFICACY OF DECAVAC VACCINE

11 The efficacy of tetanus toxoid and diphtheria toxoid used in DECAVAC vaccine was determined
12 on the basis of immunogenicity studies, with a comparison to a serological correlate of protection
13 (0.01 antitoxin units/mL) established by the Panel on Review of Bacterial Vaccines & Toxoids.⁵

14

15 A clinical study to evaluate the serological responses and adverse reactions was performed in 58
16 individuals 6 years of age and older. The results indicated protective levels of antibody were
17 achieved in greater than 90% of the study population after primary immunization with both
18 components. Booster effects were achieved in 100% of the individuals with pre-existing antibody
19 responses.¹¹

20

21 No immunogenicity data are available on concomitant administration of DECAVAC vaccine with
22 other US licensed vaccines.

1

2 INDICATIONS AND USAGE

3 DECAVAC vaccine is indicated for active immunization of persons 7 years of age or older for
4 prevention of tetanus and diphtheria. For immunization of infants and children younger than 7
5 years of age against tetanus and diphtheria, refer to the manufacturers' package inserts for
6 Diphtheria and Tetanus Toxoids and Acellular Pertussis Vaccine Adsorbed (DTaP) and for
7 Diphtheria and Tetanus Toxoids Adsorbed (For Pediatric Use) (DT).

8

9 If passive protection against tetanus is required, Tetanus Immune Globulin (Human) (TIG) may be
10 administered at a separate site with a separate needle and syringe. (See **DOSAGE AND**
11 **ADMINISTRATION** section, and **Tetanus Prophylaxis in Wound Management** subsection.)

12

13 Persons who have had tetanus or diphtheria should still be immunized since these clinical
14 infections do not always confer immunity.

15

16 As with any vaccine, vaccination with DECAVAC vaccine may not protect 100% of individuals.

17

18 CONTRAINDICATIONS

19 Hypersensitivity to any component of the vaccine is a contraindication to receipt of DECAVAC
20 vaccine. (See **DESCRIPTION** section.)

21

1 It is a contraindication to use DECAVAC vaccine after anaphylaxis or other serious allergic
2 reaction following a previous dose of this vaccine, any other tetanus or diphtheria toxoid
3 containing vaccine, or any component of this vaccine. Because of uncertainty as to which
4 component of the vaccine may be responsible, no further vaccination with diphtheria or tetanus
5 components should be carried out. Alternatively, such individuals may be referred to an allergist
6 for evaluation if further immunizations are to be considered.

7

8 **WARNINGS**

9 A booster dose of Td is recommended at 11-12 years of age if at least 5 years have elapsed since
10 the last dose of tetanus and diphtheria-toxoid containing vaccine.¹² Subsequent routine boosters
11 with Td are recommended every 10 years (see **DOSAGE AND ADMINISTRATION**).¹² More
12 frequent administration of Td is not recommended except under circumstances of wound
13 management or diphtheria prophylaxis (see **DOSAGE AND ADMINISTRATION**) since it may
14 be associated with increased incidence and severity of adverse reactions.³

15

16 Persons who experienced Arthus-type hypersensitivity reactions or a temperature of >103°F
17 (>39.4°C) following a prior dose of tetanus toxoid usually have high serum tetanus antitoxin levels
18 and should not be given even emergency doses of DECAVAC vaccine more frequently than every
19 10 years, even if they have a wound that is neither clean nor minor.^{3,13}

20

21 If Guillain-Barré Syndrome occurred within 6 weeks of receipt of a prior vaccine containing
22 tetanus toxoid, the decision to give subsequent doses of DECAVAC vaccine or any vaccine

1 containing tetanus toxoid should be based on careful consideration of the potential benefits and
2 possible risks.⁶

3

4 Because of the risk of hemorrhage, DECAVAC vaccine should not be given to persons with any
5 bleeding disorder, such as hemophilia or thrombocytopenia, or to persons on anticoagulant therapy
6 unless the potential benefit clearly outweighs the risk of administration. If the decision is made to
7 administer DECAVAC vaccine in such persons, it should be given with caution, with steps taken
8 to avoid the risk of bleeding and hematoma formation following injection.^{3,6}

9

10 The Advisory Committee on Immunization Practices (ACIP) has published guidelines for
11 vaccination of persons with recent or acute illness.⁶

12

13 **PRECAUTIONS**

14 **GENERAL**

15 Care is to be taken by the health-care provider for the safe and effective use of DECAVAC
16 vaccine.

17

18 EPINEPHRINE INJECTION (1:1000) AND OTHER APPROPRIATE AGENTS AND
19 EQUIPMENT MUST BE IMMEDIATELY AVAILABLE SHOULD AN ACUTE
20 ANAPHYLACTIC REACTION OCCUR DUE TO ANY COMPONENT OF THE VACCINE.

21

1 Prior to an injection of any vaccine, all known precautions should be taken to prevent adverse
2 reactions. This includes a review of the patient's previous immunization history, the presence of
3 any contraindications to immunization, the current health status, and history concerning possible
4 sensitivity to the vaccine or similar vaccine. (See **CONTRAINDICATIONS** section.)

5

6 Special care should be taken to ensure that the injection does not enter a blood vessel.

7

8 Immunocompromised persons (whether from disease or treatment) may not obtain the expected
9 immune response to DECAVAC vaccine.

10

11 Administration of Td vaccines is not contraindicated in immunocompromised persons.^{6,14}

12

13 A separate, sterile syringe and needle or a sterile disposable unit should be used for each patient to
14 prevent transmission of blood borne infectious agents. Needles should not be recapped and should
15 be disposed of according to biohazard waste guidelines.

16

17 INFORMATION FOR PATIENTS

18 Prior to administration of DECAVAC vaccine, health-care providers should inform the patient,
19 parent, or guardian of the benefits and risks of immunization and of the importance of completing
20 the primary immunization series or receiving recommended booster doses, as appropriate.

21

22

1 The health-care provider should inform the patient, parent, or guardian about the potential for
2 adverse reactions that have been temporally associated with the administration of DECAVAC
3 vaccine or other vaccines containing similar components. Parents or guardians, or patients should
4 be instructed to report any serious adverse reactions to their health-care provider. Adverse events
5 following immunization should be reported by health-care providers to the Vaccine Adverse Event
6 Reporting System (VAERS) (See **ADVERSE REACTIONS**, Reporting of Adverse Events).

7

8 It is extremely important when the patient returns for a subsequent dose, that the parent, guardian,
9 or patient should be questioned concerning any symptoms and/or signs of an adverse reaction after
10 a previous dose. (See **CONTRAINDICATIONS** and **ADVERSE REACTIONS** Sections).

11

12 The health-care provider should provide the Vaccine Information Statements (VISs), which are
13 required by the National Childhood Vaccine Injury Act of 1986 to be given with each
14 immunization.

15

16 **DRUG INTERACTIONS**

17 Immunosuppressive therapies, including irradiation, antimetabolites, alkylating agents, cytotoxic
18 drugs and corticosteroids (used in greater than physiologic doses), may reduce the immune
19 response to vaccines (see **PRECAUTIONS – GENERAL** section).

20

21 No information is available regarding concomitant administration of DECAVAC vaccine with
22 other US licensed vaccines.

1

2 CARCINOGENESIS, MUTAGENESIS, IMPAIRMENT OF FERTILITY No studies have been
3 performed with DECAVAC vaccine to evaluate carcinogenicity, mutagenic potential, or impact on
4 fertility.

5

6 PREGNANCY CATEGORY C

7 Animal reproduction studies have not been conducted with DECAVAC vaccine. It is also not
8 known whether DECAVAC vaccine can cause fetal harm when administered to a pregnant woman
9 or can affect reproduction capacity. DECAVAC vaccine should be given to a pregnant woman
10 only if clearly needed.

11

12 The ACIP has published recommendations for use of Tetanus and Diphtheria Toxoids Adsorbed
13 For Adult Use in pregnant women.⁶

14

15 NURSING MOTHERS

16 It is not known whether DECAVAC vaccine is excreted in human milk. Because many drugs are
17 excreted in human milk, caution should be exercised when DECAVAC vaccine is administered to
18 a nursing woman.

19

20 PEDIATRIC USE

21 DECAVAC vaccine is not indicated for infants and children younger than 7 years of age. For
22 immunization of infants and children younger than 7 years of age against tetanus and diphtheria,

1 refer to the manufacturers' package inserts for Diphtheria and Tetanus Toxoids and Acellular
2 Pertussis Vaccine Adsorbed (DTaP) and for Diphtheria and Tetanus Toxoids Adsorbed (For
3 Pediatric Use) (DT).

4

5 GERIATRIC USE

6 Clinical studies of DECAVAC vaccine did not include subjects aged 59 years and over to
7 determine whether they respond differently than younger subjects.

8

9 **ADVERSE REACTIONS**

10 Because clinical trials are conducted under widely varying conditions, adverse reaction rates
11 observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trials
12 of another vaccine and may not reflect the rates observed in practice. The adverse reaction
13 information from clinical trials does, however, provide a basis for identifying the adverse events
14 that appear to be related to vaccine use and for approximating rates.

15

16 In a clinical study involving 58 individuals 6 years of age and older, 19% of the individuals noted
17 local reactions consisting of erythema, tenderness and induration at the injection site and 2%
18 systemic reactions consisting of headache, malaise and temperature elevations.¹¹

19

20 **ADDITIONAL ADVERSE REACTIONS**

21 Additional adverse reactions, included in this section, have been reported in conjunction with
22 receipt of vaccines containing tetanus toxoid and/or diphtheria toxoid.

1

2 Arthus-type hypersensitivity reactions, characterized by severe local reactions (generally starting
3 2-8 hours after an injection), may follow receipt of tetanus toxoid. Such reactions may be
4 associated with high levels of circulating antitoxin in persons who have had overly frequent
5 injections of tetanus toxoid. (See **WARNINGS**.)¹⁵

6

7 Persistent nodules at the site of injection have been reported following the use of adsorbed
8 products.³

9

10 Cases of allergic or anaphylactic reaction (ie, hives, swelling of the mouth, difficulty breathing,
11 hypotension, or shock) have been reported after receiving some preparations containing diphtheria
12 and/or tetanus toxoid.² Death following vaccine-caused anaphylaxis has been reported.¹⁵

13

14 Certain neurological conditions have been reported in temporal association with some tetanus
15 toxoid-containing vaccines or tetanus and diphtheria toxoid-containing vaccines. A review by the
16 Institute of Medicine (IOM) concluded that the evidence favors acceptance of a causal relation
17 between tetanus toxoid and both brachial neuritis and Guillain-Barré syndrome.¹⁵ Other
18 neurological conditions that have been reported include: demyelinating diseases of the central
19 nervous system, peripheral mononeuropathies, cranial mononeuropathies, and EEG disturbances
20 with encephalopathy (with or without permanent intellectual and/or motor function impairment).
21 The IOM has concluded that the evidence is inadequate to accept or reject a causal relation
22 between these conditions and vaccines containing tetanus and/or diphtheria toxoids.¹⁵ In the

1 differential diagnosis of polyradiculoneuropathies following administration of a vaccine
2 containing tetanus toxoid, tetanus toxoid should be considered as a possible etiology.¹⁵

3

4 POSTMARKETING REPORTS

5 Additional adverse events reported between 1998-2003 during post-approval use of Tetanus and
6 Diphtheria Toxoids Adsorbed, For Adult Use, manufactured by Aventis Pasteur Inc. include local
7 reactions at injection site (ie, swelling, redness, warmth, cellulitis), myalgia, arthralgia, muscle
8 stiffness, nausea, vomiting, paraesthesia, dizziness, convulsions and rash. Events were included in
9 this list because of the seriousness or frequency of reporting. Because these events are reported
10 voluntarily from a population of uncertain size, it is not always possible to reliably estimate their
11 frequencies or to establish a causal relationship to components of Tetanus and Diphtheria Toxoids
12 Adsorbed, For Adult Use, manufactured by Aventis Pasteur Inc.¹⁶

13

14 REPORTING OF ADVERSEEVENTS

15 The National Vaccine Injury Compensation Program, established by the National Childhood
16 Vaccine Injury Act of 1986, requires physicians and other health-care providers who administer
17 vaccines to maintain permanent vaccination records of the manufacturer and lot number of the
18 vaccine administered in the vaccine recipient's permanent medical record, along with the date of
19 administration of the vaccine, and the name, address, and title of the person administering the
20 vaccine. The Act further requires the health-care professional to report to the US Department of
21 Health and Human Services the occurrence following immunization of any event set forth in the
22 Vaccine Injury Table. For Td vaccines, these include anaphylaxis or anaphylactic shock within 7

1 days; brachial neuritis within 28 days; an acute complication or sequelae (including death) of an
2 illness, disability, injury, or condition referred to above; or any events that would contraindicate
3 further doses of vaccine, according to this DECAVAC vaccine package insert.^{17,18,19}

4

5 Reporting by parents, guardians or patients of all adverse events occurring after vaccine
6 administration should be encouraged. Adverse events following immunization should be reported
7 by health-care providers to the US Department of Health and Human Services (DHHS) Vaccine
8 Adverse Event Reporting System (VAERS). Reporting forms and information about reporting
9 requirements or completion of the form can be obtained from VAERS through a toll-free number
10 1-800-822-7967.^{17,18,19}

11

12 Health-care providers should also report these events to the Pharmacovigilance Department,
13 Aventis Pasteur Inc., Discovery Drive, Swiftwater, PA 18370 or call 1-800-822-2463.

14

15 **DOSAGE AND ADMINISTRATION**

16 Parenteral drug products should be inspected visually for extraneous particulate matter and/or
17 discoloration prior to administration, whenever solution and container permit. (See
18 **DESCRIPTION** section.) If these conditions exist, the vaccine should not be administered.

19

20 **SHAKE SYRINGE WELL** *before administering the vaccine.* The vaccine, after shaking, is a
21 turbid liquid, whitish-gray in color. Discard syringe containing vaccine if the vaccine cannot be
22 resuspended.

1

2 Before injection, the skin over the site to be injected should be cleansed with a suitable germicide.

3

4 Inject 0.5 mL intramuscularly in the area of the vastus lateralis (mid-thigh laterally) or deltoid.

5 The vaccine should not be injected into the gluteal area or areas where there may be a major nerve
6 trunk.

7

8 Do not administer this product intravenously or subcutaneously.

9

10 The needle length should be sufficient to deliver the vaccine intramuscularly, but not so long as to
11 involve underlying nerves and blood vessels or bone. The health-care professional should
12 determine the appropriate size and length of the needle for individual patients.

13

14 PRIMARY IMMUNIZATION

15 DECAVAC vaccine is approved for administration in persons 7 years of age and older who have
16 not been immunized previously against tetanus and diphtheria, as a primary immunization series
17 of three 0.5 mL doses. For primary immunization with Td vaccines, the intervals between doses
18 recommended by the Advisory Committee on Immunization Practices (ACIP) are 4 to 8 weeks
19 between the first and second dose, and 6 to 12 months between the second and third dose.³

20

21 DECAVAC vaccine may be used to complete the primary immunization series for tetanus and
22 diphtheria in children 7 years of age or older who have received one or two doses of whole-cell

1 pertussis DTP, DTaP and/or DT vaccine. However, the safety and efficacy of DECAVAC vaccine
2 in such children have not been evaluated.

3

4 Interruption of the recommended schedule with a delay between doses should not interfere with
5 the final immunity achieved with DECAVAC vaccine. There is no need to start the series over
6 again, regardless of the time elapsed between doses.³

7

8 Routine Booster Immunization

9 DECAVAC vaccine is approved for booster immunization in persons 7 years of age and older who
10 have completed primary immunization against tetanus and diphtheria.

11

12 A booster dose of Td is recommended by the ACIP in persons 11-12 years of age if at least 5 years
13 have elapsed since the last dose of tetanus and diphtheria toxoid-containing vaccine.¹² Subsequent
14 routine boosters with Td are recommended every 10 years.^{12,19} If a dose is given sooner than 10
15 years, as part of wound management or on exposure to diphtheria, the next booster is not needed
16 for 10 years thereafter.³ MORE FREQUENT BOOSTER DOSES ARE NOT RECOMMENDED
17 AND MAY BE ASSOCIATED WITH INCREASED INCIDENCE AND SEVERITY OF
18 ADVERSE REACTIONS.^{3,6} (See **WARNINGS** section.)

19

1

2 DIPHtheria PROPHYLAXIS FOR CASE CONTACTS

3 The ACIP has published recommendations on vaccination for diphtheria prophylaxis in
4 individuals who have had contact with a person with confirmed or suspected diphtheria.³

5

6 TETANUS PROPHYLAXIS IN WOUND MANAGEMENT

7 The need for active immunization with a tetanus toxoid-containing preparation, with or without
8 passive immunization with TIG (Human) depends on both the condition of the wound and the
9 patient's vaccination history (Table 1).

10

11 A thorough attempt must be made to determine whether a patient has completed primary
12 immunization. Individuals who have completed primary immunization against tetanus, and who
13 sustain wounds which are minor and uncontaminated, should receive a booster dose of a tetanus
14 toxoid-containing preparation only if they have not received tetanus toxoid within the preceding
15 10 years. For tetanus prone wounds (eg, wounds contaminated with dirt, feces, soil, and saliva;
16 puncture wounds; avulsions; and wounds resulting from missiles, crushing, burns, and frostbite), a
17 booster is appropriate if the patient has not received a tetanus toxoid-containing preparation within
18 the preceding 5 years. If a booster dose is given sooner than 10 years as part of wound
19 management, the next routine booster should not be given for 10 years thereafter.³

20

21 Individuals who have not completed primary immunization against tetanus, or whose
22 immunization history is unknown or uncertain, should be immunized with a tetanus toxoid-

1 containing product. Completion of primary immunization thereafter should be ensured. In
 2 addition, if these individuals have sustained a tetanus-prone wound, the use of TIG (Human) is
 3 recommended. TIG (Human) should be administered at a separate site, with a separate needle and
 4 syringe, according to the manufacturer's package insert. If a contraindication to using tetanus
 5 toxoid-containing preparations exists in a person who has not completed a primary immunizing
 6 course of tetanus toxoid and other than a clean, minor wound is sustained, only passive
 7 immunization with TIG (Human) should be given.³

8

9 Td is the recommended preparation for active tetanus immunization in wound management of
 10 patients =7 years of age.³ In such persons, a preparation containing tetanus and diphtheria toxoids
 11 is preferred instead of single-antigen tetanus toxoid to enhance diphtheria protection. DECAVAC
 12 vaccine is approved for wound management in patients 7 years of age and older.

13

14 TABLE 1³ SUMMARY GUIDE TO TETANUS PROPHYLAXIS IN ROUTINE

15 WOUND MANAGEMENT FOR PERSONS 7 YEARS OF AGE AND OLDER*

History of Adsorbed Tetanus Toxoid (doses)	Clean, Minor Wounds		All Other Wounds**	
	Td§	TIG	Td§	TIG
Unknown or < three = Three [¶]	Yes No†	No No	Yes No‡	Yes No

16 * Important details are in the text of the DOSAGE AND ADMINISTRATION section.

17 **Such as, but not limited to, wounds contaminated with dirt, feces, soil, and saliva; puncture
 18 wounds; avulsions; and wounds resulting from missiles, crushing, burns, and frostbite.

1 ¶ If only three doses of fluid tetanus toxoid have been received, then a fourth dose of toxoid,
2 preferably an adsorbed toxoid should be given.

3 † Yes, if > 10 years since last dose.

4 ‡ Yes, if > 5 years since last dose. (More frequent boosters are not needed and can accentuate
5 side effects.)

6 § DECAVAC vaccine is approved for wound management in persons 7 years of age or older.

7

8 CONCOMITANT VACCINE ADMINISTRATION

9 No safety and immunogenicity data are available on the concomitant administration of
10 DECAVAC vaccine with other US licensed vaccines.

11

12 HOW SUPPLIED

13 Luer-Lok® latex-free syringe, 0.5 mL (10 x 0.5 mL syringes per package) – Product No. 49281-
14 291-10

15 Luer-Lok® is a trademark of Becton Dickinson and Company.

16

17 CPT® Code: 90718

18 CPT is a registered trademark of the American Medical Association.

19

20 STORAGE

21 Store between 2° - 8°C (35° - 46°F). DO NOT FREEZE.

22

1 Do not use vaccine after expiration date.

2

3 **REFERENCES**

4 1. Mueller JH, et al. Production of diphtheria toxin of high potency (100 Lf) on a
5 reproducible medium. *J Immunol* 40: 1941; 21-32.

6 2. Pascual FB, et al. Tetanus Surveillance-United States, 1998-2000. In: *Surveillance*
7 *Summaries*, June 20, 2003. *MMWR* 2003;52(No.SS-3):1-8.

8 3. Recommendations of the Immunization Practices Advisory Committee (ACIP).
9 Diphtheria, Tetanus, and Pertussis: Recommendations for vaccine use and other preventive
10 measures. *MMWR* 40: 1991; No. RR-10.

11 4. Wassilak SGF, et al. Tetanus Toxoid. In: Plotkin SA and Orenstein WA, eds. *Vaccines*. 3rd
12 ed. Philadelphia, PA: W.B. Saunders Company;1999:441-474.

13 5. Department of Health and Human Services, Food and Drug Administration. Biological
14 Products; Bacterial Vaccines and Toxoids; Implementation of Efficacy Review; Proposed
15 Rule. *Federal Register* Vol50 No 240: 1985; 51002-51117.

16 6. CDC. General recommendations on immunization: Recommendations of the Advisory
17 Committee on Immunization Practices (ACIP) and the American Academy of Family
18 Physicians (AAFP). *MMWR* 2002;51(RR-2):1-35.

19 7. CDC. Diphtheria. In: Atkinson W and Wolfe C, eds. *Epidemiology and Prevention of*
20 *Vaccine-Preventable Diseases*. 7th ed. Atlanta, GA: Public Health Foundation. 2002:39-48.

21 8. Dittmann S, et al. Successful control of epidemic diphtheria in the states of the former
22 Union of Soviet Socialist Republics: Lessons Learned. *J Infect Dis* 2000;181:S10-S22.

- 1 9. Hardy IRB, et al. Current situation and control strategies for resurgence of diphtheria in
2 newly independent states of the former Soviet Union. *Lancet* 1996;347:1739-1744.
 - 3 10. Mortimer EA and Wharton M. Diphtheria toxoid. In: Plotkin SA and Orenstein WA, eds.
4 *Vaccines*. 3rd ed. Philadelphia, PA: W.B. Saunders Company,1999:140-157.
 - 5 11. Myers MG, et al. Primary immunization with tetanus and diphtheria toxoids. *JAMA* 248:
6 1982; 2478-2480.
 - 7 12. CDC. Recommended childhood and adolescent immunization schedule-United States,
8 2003. *MMWR* 2003;52(04)Q1-Q4.
 - 9 13. CDC. Update: Vaccine side effects, adverse reactions, contraindications, and precautions
10 recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR*
11 1996;45(RR-12):22-31.
 - 12 14. CDC. Use of vaccines and immune globulins in persons with altered immunocompetence.
13 *MMWR* 1993; 42(RR-04):1-9.
 - 14 15. Institute of Medicine (US). Stratton KR, et al, eds. *Adverse events associated with*
15 *childhood vaccines: evidence bearing on causality*. Washington (DC): National Academy
16 Press. 1994:67-117.
 - 17 16. Data on File, 030104.
 - 18 17. CDC. Vaccine Adverse Event Reporting System – United States. *MMWR* 39: 1990;730-
19 733.
 - 20 18. CDC. National Childhood Vaccine Injury Act: requirements for permanent vaccination
21 records and for reporting of selected events after vaccination. *MMWR* 37: 1988; 197-200.
-

1 19. Food and Drug Administration. New reporting requirements for vaccine adverse events.

2 FDA Drug Bull 18 (2): 1988; 16-18.

3 20.CDC. Recommended adult immunization schedule-United States, 2002-2003. MMWR

4 2002;51(40):904-908.

5

6

7 Manufactured by:

Product information

8 **Aventis Pasteur Inc.**

as of March 2004

9 Swiftwater PA 18370 USA

10 ***Aventis Pasteur***

The logo for Aventis, featuring a stylized plant or star symbol to the left of the word "Aventis" in a bold, italicized sans-serif font.

11

4935/4936