

### Current Status of Smallpox Vaccine

**To the Editor:** The possible use of smallpox virus as a weapon by terrorists has stimulated growing international concern and led to a recent review by the World Health Organization of the global availability of smallpox vaccine. This review found approximately 60 million doses worldwide, with little current vaccine manufacture, although limited vaccine seed remains available (1). Ongoing discussions in the United States suggest that the national stockpile should contain at least 40 million doses to be held in reserve for emergency use, including in case of a terrorist release of smallpox virus (O'Toole, this issue, pp. 540-6).

The current U.S. stockpile contains approximately 15.4 million doses of vaccinia vaccine (Dryvax) made from the New York City Board of Health strain of vaccinia and was produced by Wyeth Laboratories in 13 separate lots. The vaccine is lyophilized in glass vials with rubber stoppers and sealed with a metal band. When rehydrated, each vial contains 100 doses and has a potency of at least  $10^8$  plaque-forming units (pfu)/ml. Some vials of the vaccine stockpile have shown elevated moisture levels and thus failed routine quality control testing; however, the vaccine in these vials remains potent, and the failed lots have not been discarded.

The diluent used to rehydrate the vaccine contains brilliant green, which makes the vaccine easier to visualize when administered with bifurcated needles. Over time, the brilliant green has deteriorated, and most of the available diluent does not pass quality control. Discussions are under way with Wyeth to begin production of sufficient new diluent for the entire stockpile.

The vaccine is administered by superficial inoculation (scarification) with a bifurcated needle. Fewer than 1 million bifurcated needles are held as part of the stockpile. As with the diluent, Wyeth has been requested to produce additional bifurcated needles.

Vaccinia virus produces adverse reactions in a small percentage of vaccinated persons. Adverse reactions are treated with vaccinia immune globulin (VIG), currently only available from Baxter Healthcare Corporation (5,400 vials of VIG in stock). Each vial contains 5 ml of VIG; the recommended dose for postvaccine complica-

tions is 0.6 ml per kg of body weight. This volume is sufficient to treat adverse reactions in approximately 675 adults. Further, the entire stockpile of VIG has been placed on hold while the cause of a slight pink discoloration is investigated. Until the cause of the discoloration is determined or another approved supply of VIG is obtained, no vaccinia vaccine is being released. While unknown, the rate of adverse reactions in today's population is likely to be greater than seen during the global eradication campaign because of recent increases in the number of immunocompromised persons. The Department of Defense has recently contracted the processing of new lots of VIG (to be administered intravenously rather than by the intramuscular route like existing VIG stocks); however, maintaining adequate stocks of VIG will remain a challenge.

In the event of release of smallpox virus, persons at high risk and persons exposed but not yet showing clinical illness would be vaccinated immediately. Intensive case detection and vaccination of contacts and other persons at risk would follow. All vaccine, including lots retained after failed quality control tests, would be made available for emergency use. Previous studies have found that more than 90% of susceptible persons respond to vaccinia virus with a titer of  $10^7$  pocks/ml (2). In an emergency, consideration would be given to diluting the existing vaccine as much as 10-fold, so that each vial could conceivably contain 1,000 doses of vaccine, rather than the current 100 doses. The present vaccine container is sufficiently large to accommodate the added diluent. The absence of sufficient quantities of VIG to protect against adverse reactions during a mass immunization campaign would necessitate careful screening of those receiving the vaccine; some persons with adverse reactions would likely go untreated.

While the intentional release of smallpox virus would represent a global emergency, the existing national stockpile could be effectively used to limit the spread of disease and buy time while the pharmaceutical industry begins emergency vaccine production.

**James W. LeDuc and John Becher**

Centers for Disease Control and Prevention, Atlanta, Georgia, USA

## References

1. World Health Organization. Report of the meeting of the ad hoc committee on Orthopox virus infections. Department of Communicable Disease Surveillance and Response. WHO, 14-15 January 1999.
2. Cockburn WC, Cross RM, Downie AW, Dumbell KR, Kaplan C, Mclean D, et al. Laboratory and vaccination studies with dried smallpox vaccines. *Bull World Health Organ* 1957;16:63-77.

## West Nile Fever in Czechland

**To the Editor:** After heavy rains in July 1997, extensive floods occurred along the Morava River, Czech Republic. Populations of *Aedes* mosquitoes increased rapidly in the flooded areas, prompting surveillance for mosquito-borne virus infections in the Breclav area, South Moravia. We collected 11,334 female mosquitoes (9,100 *Aedes vexans*, 917 *Ae. cinereus*, 11 *Ae. cantans*, 1,074 *Ae. sticticus*, and 232 *Culex p. pipiens*) from July through September 1997 and tested them for virus in 117 monospecific pools by intracranial inoculation of suckling mice. Seven virus isolates were obtained and identified by complement-fixation and neutralization tests. Six isolates (five from *Ae. vexans*, one from *Ae. cinereus*) were identified as the bunyavirus Tahyna, California serogroup, and one (strain 97-103 from 57 *C. p. pipiens* collected at Lanzhot, 48°40'N, 16°56'E, on September 17) was identified as the flavivirus West Nile (1). A crossed comparison of 97-103 and topotype Eg-101 (2) West Nile virus strains and their antisera (prepared in mice by three intraperitoneal doses at weekly intervals) by plaque reduction neutralization (PRN) on XTC-2 cells (3,4) showed their antigenic relationships: reciprocal titers of homologous/heterologous sera were 512/512 in Eg-101 and 512/64 in 97-103. Strain 97-103 has lower virulence than Eg-101 in that it does not kill adult ICR mice and may represent a subtype of West Nile virus.

Blood samples were obtained from 619 persons seeking treatment at hospital and outpatient clinics in the Breclav area from June 23 through September 29, 1997. Sera were inactivated at 56°C for 30 minutes, diluted 1:8, and assayed by PRN for antibodies against *c.* 30 plaque-forming units (PFU) per well of West Nile virus strains Eg-101 and 97-103. All sera causing 90% reduction of PFU at 1:8 dilution were titrated, and the highest serum dilution showing

50% PFU reduction was regarded as the titer. Antibodies neutralizing West Nile virus were detected in 13 (2.1%) persons: 2.8% of 179 male and 1.8% of 440 female. Persons with detectable West Nile virus antibody were questioned about their health history during the previous 5 years, and their medical records were reviewed; none recalled having had tickborne encephalitis (Central-European encephalitis [CEE] virus is the only other flavivirus present in Czechland) or having been vaccinated against CEE or yellow fever virus. Titers of PRN antibodies to CEEV were all below 16. Two of the seropositive persons had traveled abroad during the last 5 years: one to Croatia in 1996, and one to South Australia during 1951 to 1994.

Paired serum samples were obtained from 72 of the 619 persons examined. A significant increase ( $\geq 4$  times) in antibody titer against West Nile virus between the first (acute-phase) and second (convalescent-phase) samples was detected four times: in 2 of 41 young persons ( $\leq 16$  years of age) and in 2 of 31 adults ( $> 16$  years of age). Among the four seroconverting persons, only the two children had clinical symptoms compatible with West Nile fever. A 9-year-old boy had fever (39°C) for 4 days, sore throat, headache, muscle ache, pronounced fatigue, and nausea lasting approximately 6 days, with recovery after 13 days. Neutralizing antibodies to West Nile virus, Eg-101 and 97-103, were 64 and 32 on July 22 and 512 and 256 on August 4, respectively. A 9-year-old girl had fever (38°C-39°C) for 3 days, sore throat, headache, muscle ache, pronounced fatigue, nausea, vomiting, maculopapular rash (including flushed face), and slightly enlarged inguinal lymph nodes. The illness lasted approximately 7 days, with complete recovery after 17 days. Neutralizing antibodies to West Nile virus, Eg-101 and 97-103, were 64 and 32 on August 6 and 256 and 128 on August 20, respectively. Of the remaining nine seropositive persons lacking paired serum samples, one had severe headache, muscle ache, prolonged fatigue, nausea, pain on eye movement, maculopapular rash, and insomnia in summer of 1997. Two other persons had had "summer fever" (sore throat and lymphadenitis; headache with pain on eye movement) in 1997. The other persons who seroconverted did not report any substantial illness. In total, clinical symptoms in five persons are compatible with West Nile fever.

These are the first reported human cases of West Nile fever in Central Europe (5); an extensive outbreak occurred in Romania in 1996, with approximately 500 patients hospitalized and a 4% to 8% fatality rate (6,7). West Nile virus should be viewed as a potential agent of local sporadic cases, clusters, or outbreaks, even in temperate Europe. Environmental factors (including human activities) that enhance vector population densities (heavy rains followed by floods, irrigation, higher than usual temperatures due to global warming) might produce an increased incidence of West Nile fever and other new or reemerging mosquito-borne diseases. Surveillance for West Nile fever should monitor population density and infection rate of principal vectors, antibodies in vertebrates and exposed human groups, and routine diagnosis of human infections.

**Zdenek Hubálek, Jiri Halouzka, and  
Zina Juricová**

Institute of Vertebrate Biology, Academy of Sciences,  
Brno, Czech Republic

### References

1. Hubálek Z, Halouzka J, Juricová Z, Šebesta O. First isolation of mosquito-borne West-Nile virus in the Czech Republic. *Acta Virol* 1998;42:119-20.
2. Melnick JL, Paul JR, Riordan JF, Barnett VHH, Goldblum N, Zabin E. Isolation from human sera in Egypt of a virus apparently identical to West Nile virus. *Proc Soc Exp Biol Med* 1951;77:661-5.
3. de Madrid AT, Porterfield JS. The flaviviruses (group B arboviruses): a cross-neutralization study. *J Gen Virol* 1974;23:91-6.
4. Leake CJ, Varma MGR, Pudney M. Cytopathic effect and plaque formation by arboviruses in a continuous cell line (XTC-2) from the toad *Xenopus laevis*. *J Gen Virol* 1977;35:335-9.
5. Hubálek Z, Halouzka J. Arthropod-borne viruses of vertebrates in Europe. *Acta Scientiarum Naturalium Academiae Scientiarum Bohemicae Brno* 1996;30, no. 4-5:1-95.
6. Le Guenno B, Bougermouh A, Azzam T, Bouakaz R. West Nile: a deadly virus? *Lancet* 1996;348:1315.
7. Tsai TF, Popovici F, Cernescu C, Campbell GL, Nedelcu NI. West Nile encephalitis epidemic in southeastern Romania. *Lancet* 1998;352:767-71.

### Ofloxacin-Resistant *Vibrio cholerae* O139 in Hong Kong

**To the Editor:** Unexpected outbreaks of cholera occurred in many areas of the world in 1997-98, partly because of weather changes associated

with the El Niño phenomenon (1). Outbreaks caused by antibiotic-resistant *Vibrio cholerae* O1 and O139 have been documented in the Indian subcontinent (2-4), Africa (5), and Ukraine (6).

In Hong Kong, nonduplicate bacterial strains of *V. cholerae* O1 and O139 isolated from patients and environmental sources and received in the Public Health Laboratory between January 1, 1993, and June 30, 1998, were identified by conventional biochemical tests (7,8) and API 20E (bioMérieux, France); serotyped by slide agglutination with polyvalent O1 and mono-specific Inaba and Ogawa antisera (Murex, Dartford, United Kingdom); and checked with O139 antiserum (Denka Seiken, Tokyo, Japan). Biotyped and antibiotic susceptibilities were determined by the Kirby-Bauer disk-diffusion assay (8-10). Antibiotics tested included chloramphenicol and tetracycline (from 1993 to 1996) and ofloxacin (added in routine testing from 1997). *V. cholerae* isolates available for further study were tested with the standard broth microdilution method (11) to measure minimum inhibitory concentrations (MICs) of susceptibilities to chloramphenicol, tetracycline, and ofloxacin.

No antibiotic resistance was seen in *V. cholerae* isolates in testing conducted from 1969 to 1995. The first *V. cholerae* isolate with reduced susceptibility to chloramphenicol but sensitive to tetracycline was encountered in Hong Kong in 1996. This O1 El Tor Ogawa strain was imported from Nepal. Since then, more O1 strains were isolated that exhibited reduced antibiotic susceptibilities to chloramphenicol and tetracycline but not to ofloxacin (12). In May 1998, seven *V. cholerae* O139 strains were isolated that displayed patterns of antibiotic susceptibilities strikingly different from those of O1 isolates; the former were all sensitive to tetracycline but showed reduced susceptibilities to chloramphenicol and ofloxacin. All *V. cholerae* O1 strains tested have been susceptible to ofloxacin; O1 isolates falling into intermediate categories for chloramphenicol and tetracycline susceptibilities (31% and 27.6%, respectively) were common.

The first isolate of *V. cholerae* O139 in Hong Kong came from the imported case of a patient who had traveled to other provinces of China (13,14). Isolation of O139 continued sporadically since then, with six cases between 1993 and the

1st quarter of 1998. In May 1998, a cluster of seven imported cases of *V. cholerae* O139 were reported with strains isolated from seven persons who became ill with severe diarrhea after visiting Zhuhai in Guangdong Province, China. Of 13 *V. cholerae* O139 isolates tested, 7 showed intermediate resistance to chloramphenicol and high-level resistance to ofloxacin (MIC 16 µg/ml) but no resistance to tetracycline (MIC 50s and MIC 90s were 0.25 µg/ml). This is the first evidence of a quinolone-resistant strain of *V. cholerae* O139 in Hong Kong. Of the O1 isolates, none were resistant to chloramphenicol and ofloxacin, but six were resistant to tetracycline (MIC 50s and MIC 90s were 0.25 µg/ml and 8 µg/ml, respectively).

Although all O1 isolates were sensitive to chloramphenicol, there was only a twofold difference in MIC<sub>90</sub> to chloramphenicol between O1 and O139 isolates. MIC<sub>90</sub>s of ofloxacin for O139 were nearly 10 times higher than those for O1 strains.

The novel appearance of O139 resistant to ofloxacin with MICs of 16 µg/ml from Guangdong Province, China, was of special concern. Preliminary results using pulsed-field gel electrophoresis analysis of chromosomal DNA showed that these ofloxacin-resistant O139 strains had identical fingerprint patterns and probably belonged to the clone that had caused severe diarrheal disease in the region. Two previous surveys of *V. cholerae* antibiotic susceptibilities had not described any ofloxacin-resistant O139 strains (15,16). The potential for rapid spread of these strains threatens cholera prevention and control efforts that may still rely on chemotherapy.

Different antimicrobial resistance patterns of *V. cholerae* O1 and O139 were noted. Among the resistant O1 isolates, four were local, one was from other provinces of China, and one was from Thailand. All the resistant O139 isolates were imported from Guangdong Province, China. Antibiotic resistance was found in strains from local isolates and from neighboring countries. The unique patterns of antimicrobial resistance for the O1 and O139 isolates suggest different mechanisms of resistance. As quinolones are used heavily in this region to treat cholera and other enteric diseases, selective pressure could encourage emergence of ofloxacin resistance. Prudent use of antibiotics should be exercised during

antimicrobial therapy and prophylaxis for cholera and other enteric diseases to decrease the selection of more resistant clones in our locality.

**Kai Man Kam, Kit Yee Luey, Tze Leung  
Cheung, Kwan Yee Ho, Kwok Hang Mak, and  
Paul Thian Aun Saw**

Department of Health, Hong Kong SAR Government

### References

1. World Health Organization, Geneva. Cholera in 1997. *Wkly Epidemiol Rec* 1998;73:201-8.
2. Gharagozloo RA, Naficy K, Mouin M, Nassirzadeh MH, Yalda R. Comparative trial of tetracycline, chloramphenicol, and trimethoprim/sulphamethoxazole in eradication of *Vibrio cholerae* El Tor. *BMJ* 1970;4:281-2.
3. Glass RI, Huq I, Alim AR, Yunus M. Emergence of multiply antibiotic-resistant *Vibrio cholerae* in Bangladesh. *J Infect Dis* 1980;142:939-42.
4. Ramamurthy T, Garg S, Sharma R, Bhattacharya SK, Nair GB, Shimada T, et al. Emergence of novel strain of *Vibrio cholerae* with epidemic potential in southern and eastern India. *Lancet* 1993;341:703-4.
5. Finch MJ, Morris JG Jr, Kaviti J, Kagwanja W, Levine MM. Epidemiology of antimicrobial resistant cholera in Kenya and east Africa. *Am J Trop Med Hyg* 1988;39:484-90.
6. Clark CG, Kravetz AN, Alekseenko VV, Krendellev YuD, Johnson WM. Microbiological and epidemiological investigation of cholera epidemic in Ukraine during 1994 and 1995. *Epidemiol Infect* 1998;121:1-13.
7. Cowan ST, Steel KJ. Manual for the identification of medical bacteria. 2nd ed. Cambridge: Cambridge University Press; 1974.
8. Bradford AK, Bopp CA, Wells JG. Isolation and identification of *Vibrio cholerae* O1 from fecal specimens. In: Wachsmuth IK, Blake AB, Olsvik Ø, editors. *Vibrio cholerae* and cholera: molecular to global perspectives. Washington: American Society for Microbiology; 1994. p. 3-26.
9. National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial disk susceptibility tests approved standard. NCCLS document M2-A6. Villanova (PA): The Committee; 1997.
10. National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial susceptibility testing; NCCLS document M100-S8, 18(1). Villanova (PA): The Committee; 8th information supplement, 1998.
11. National Committee for Clinical Laboratory Standards. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. 4th ed. NCCLS document M7-A4. Villanova (PA): The Committee; 1997.
12. Wong W, Ho YY. Imported cholera cases among tourists returning from Thailand. *Public Health & Epidemiology Bulletin*, Department of Health, Hong Kong. 1998;7:21-4.
13. Kam KM, Leung TH, Ho PYY, Ho KY, Saw TA. Outbreak of *Vibrio cholerae* O1 in Hong Kong related to contaminated fish tank water. *Public Health* 1995;109:389-95.

14. Lee SH, Lai ST, Lai JY, Leung NK. Resurgence of cholera in Hong Kong. *Epidemiol Infect* 1996;117:43-9.
15. Yamamoto T, Nair GB, Albert MJ, Parodi CC, Takeda Y. Survey of in vitro susceptibilities of *Vibrio cholerae* O1 and O139 to antimicrobial agents. *Antimicrob Agents Chemother* 1995;39:241-4.
16. Sciortino CV, Johnson JA, Hamad A. Vitek system antimicrobial susceptibility testing of O1, O139, and non-O1 *Vibrio cholerae*. *J Clin Microbiol* 1996;34:897-900.

### Plant Pathology and Public Health

The day will come when the sign of the plant pathologist will stand forth in the street alongside that of the physician and surgeon. . . . For what will it profit us if all the ills and diseases of the human race be banished and we then face starvation because of diseases and pests in our food (1).

**To the Editor:** Every year plant diseases affect human society, resulting in inadequate nutrition and economic loss. The potato famine in the mid-1800s is the best-known example of a fungal plant pathogen's effect on history (2-4); *Phytophthora infestans* has recently reemerged in the Americas (5). Among the silent problems that have enormous effects on human society each year are crop infections by geminiviruses and tomato spotted wilt virus (6). These plant viruses are transmitted by whiteflies, leafhoppers, or thrips to hundreds of species of plants. They cause diseases of crops and ornamental plants around the world.

More obvious problems include ergotism, caused by the alkaloids produced by the fungus *Claviceps purpurea*. Ergotism was associated with the growth of rye, particularly in cool climates that cannot support wheat, and was implicated in the aberrant human behavior responsible at least in part for the Salem witch trials and St. Anthony's fire (2,7). In the last 5 years, a new plant disease, sorghum ergot (*Claviceps africana*), has spread north from Brazil into the United States. This fungus also causes disease in Australia, a sudden change from its known occurrence in Africa (8). Sorghum is the fifth most important cereal crop in the world, with approximately 45 million hectares under cultivation for food, beverages, feed, and fodder (8). Ergot alkaloid toxicity has not yet been demonstrated, but potential nutritional and economic losses could have substantial impact on public health.

With our increased awareness of the fragility

of the environment, including the quality of our drinking water, opportunities may exist for physicians to interact with plant pathologists. Concern is growing about the use of *Burkholderia cepacia*, a bacterial phytopathogen, for the biologic control of seedling diseases (9). Although *B. cepacia* is effective for the biologic control of fungal diseases in the agricultural environment (10), this bacterium could contaminate the public water supply and subsequently influence the health of the immunosuppressed or persons with cystic fibrosis (9-11). This risk exemplifies the need to integrate plant health measures with human and veterinary health guidelines.

Plant pathology and public health also intersect with post-harvest fungal infections of seed and grain, particularly *Aspergillus flavus* and *Fusarium moniliforme* (2), which produce aflatoxin and fumonisin, respectively. During the past 2 drought years in Texas, aflatoxin in contaminated corn and peanuts has become a public health problem. In 1998, more than 50 pet dogs died of aflatoxicosis, perhaps by eating aflatoxin B1-contaminated corn used in dog food (12).

Although the veterinary and medical communities are well aware of the risks associated with plant pathogens when they enter the animal or human food supply, more routine interactions with plant pathologists could benefit public health. For example, plant pathologists can often predict impending plant disease outbreaks. This information can be used by epidemiologists to sound a warning about impending food shortages or poor food quality, particularly in developing countries. Plant pathologists are also developing new types of resistance in host plants and alternative strategies for managing plant diseases. These measures should improve food quality and reduce the negative public health impact associated with plant diseases.

**Karen-Beth G. Scholthof**

Texas A&M University, College Station, Texas, USA

### References

1. Whetzel HH. The relation of plant pathology to human affairs. Mayo Foundation Lectures 1926-1927. Philadelphia: W.B. Saunders Co.; 1928. p. 151-78.
2. Hudler GW. Magical mushrooms, mischievous molds. Princeton: Princeton University Press; 1998.
3. Woodham-Smith C. The great hunger: Ireland 1845-1849. New York: Old Town Books; 1962.

- Schumann GL. Plant diseases: their biology and social impact. St. Paul: American Phytopathological Society Press; 1993.
- Goodwin SB, Smart CD, Sandrock RW, Deahl KL, Punja ZK, Fry WE. Genetic change within populations of *Phytophthora infestans* in the United States and Canada during 1994 to 1996—role of migration and recombination. *Phytopathology* 1998;88:939-49.
- Prins M, Goldbach R. The emerging problem of tospovirus infection and nonconventional methods of control. *Trends Microbiol* 1998;6:31-5.
- Matossian MK. Poisons of the past: molds, epidemics, and history. New Haven: Yale University Press; 1989.
- Bandyopadhyay R, Frederickson DE, McLaren NW, Odvody GN, Ryley MJ. Ergot: a new disease threat to sorghum in the Americas and Australia. *Plant Disease* 1998;82:356-67.
- Holmes A, Govan J, Goldstein R. Agricultural use of *Burkholderia (Pseudomonas) cepacia*: a threat to human health? *Emerg Infect Dis* 1998;4:221-7.
- Park JL. *Burkholderia cepacia*: friend or foe? 1998. Available from: URL: <http://www.scisoc.org/feature/BurkholderiaCepacia/>
- King EB, Parke JL. Population density of the biocontrol agent *Burkholderia cepacia* AMMDR1 on four pea cultivars. *Soil Biology and Biochemistry* 1996;28:307-12.
- Texas Veterinary Diagnostic Medical Laboratory. *Epidemiol Bull* 1998. Available from: URL: <http://www.tvmdl.tamu.edu/>

### Pet-Associated Zoonoses

**To the Editor:** We read with interest the article by Grant and Olsen on preventing zoonotic diseases in immunocompromised persons (1). We completely agree with the benefits of communication between physicians and veterinarians. However, we want to emphasize that pet-associated illnesses are not limited to the immunocompromised; pregnant women and young infants should be included in this high-risk category. Our recently published survey (2) reaffirms the need for education of the general public, parents, and—to a lesser extent—pediatricians regarding pet-associated hazards.

**Leslie L. Barton,\* Rodrigo G. Villar,† and Megan Connick‡**

\*University of Arizona Health Sciences Center, Tucson, Arizona, USA; †Indian Health Service, Gallup, New Mexico, USA; and ‡Washington University School of Medicine, St. Louis, Missouri, USA

### References

- Grant S, Olsen CW. Preventing zoonotic diseases in immunocompromised persons: the role of physicians and veterinarians. *Emerg Infect Dis* 1999;5:159-63.
- Villar RG, Connick M, Barton LL. Parent and pediatrician knowledge, attitudes, and practices regarding pet-associated hazards. *Arch Pediatr Adolesc Med* 1998;152:1035-7.