PV 3393 AMP

### **ZYPREXA®** (Olanzapine) Tablets

### ZYPREXA® ZYDIS® (Olanzapine) Orally Disintegrating Tablets

DESCRIPTION

**DESCHIPTION**ZYPREXA (olanzapine) is a psychotropic agent that belongs to the thienobenzodiazepine class. The chemical designation is 2-methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b] [1,5]benzodiazepine. The molecular formula is C<sub>17</sub>H<sub>20</sub>N<sub>4</sub>S, which corresponds to a molecular weight of 312.44. The chemical structure is:

Olanzapine is a vellow crystalline solid, which is practically insoluble in water.

Olanzapine is a yellow crystalline solid, which is practically insoluble in water.

ZYPREXA tablets are intended for oral administration only.

Each tablet contains olanzapine equivalent to 2.5 mg (8 μmol), 5 mg (16 μmol), 7.5 mg (24 μmol), 10 mg (32 μmol), 15 mg (48 μmol), or 20 mg (64 μmol). Inactive ingredients are carnauba wax, crospovidone, hydroxypropyl cellulose, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, and other inactive ingredients. The color coating contains Titanium Dioxide (all strengths), FD&C Blue No. 2 Aluminum Lake (15 mg), or Synthetic Red Iron Oxide (20 mg). The 2.5, 5.0, 7.5, and 10 mg tablets are imprinted with edible ink which contains FD&C Blue No. 2 Aluminum Lake.

ZYPREXA ZYDIS (olanzapine orally disintegrating tablets) is intended for oral administration only. Each orally disintegrating tablet contains olanzapine equivalent to 5 mg (16 µmol), 10 mg (32 µmol), 15 mg (48 µmol) or 20 mg (64 µmol). It begins disintegrating in the mouth within seconds, allowing its contents to be subsequently swallowed with or without liquid. ZYPREXA ZYDIS (olanzapine orally disintegrating tablets) also contains the following inactive ingredients: gelatin, ananitol, asparame, sodium methyl paraben and

### CLINICAL PHARMACOLOGY

Olanzapine is a selective monoaminergic antagonist with high affinity binding to the following receptors: serotonin  $SHT_{2A/2C}$  ( $K_{[=4]}$  and 11 nM, respectively), dopamine  $D_{1-4}$  ( $K_{[=1]}$  1-31 nM), muscarinic  $M_{1-5}$  ( $K_{[=1]}$  5-6 nM), histamine  $H_1$  ( $K_{[=7]}$  nM), and adrenergic  $\alpha_1$  receptors ( $K_{[=19]}$  nM). Olanzapine binds weakly to  $GABA_n$ ,  $B2D_n$ , and  $\beta$  adrenergic receptors ( $K_{[=19]}$  nM).

ouanzapine binds weakly to GABA, BZD, and  $\beta$  adrenergic receptors (K>10  $\mu$ M). The mechanism of action of olanzapine, as with other drugs having efficacy in schizophrenia, is unknown. However, it has been proposed that this drug's efficacy in schizophrenia is mediated through a combination of dopamine and serotonin type 2 (5HT<sub>2</sub>) antagonism. The mechanism of action of olanzapine in the treatment of acute menic episodes associated with Bipolar I Disporter is unknown. Antagonism at receptors other than dopamine and 5HT<sub>2</sub> with similar receptor affinities may explain some of the other therapeutic and side effects of olanzapine. Olanzapine's antagonism of muscarinic  $M_{1.5}$  receptors may explain its anticholinergic effects. Olanzapine's antagonism of histamine H<sub>1</sub> receptors may explain the somnolence observed with this drug. Olanzapine's antagonism of adrenergic  $\alpha_1$  receptors may explain the orthostatic hypotension observed with this drug.

FIGURE 1. Collarapine is well absorbed and reaches peak concentrations in approximately 6 hours following an oral dose. It is eliminated extensively by first pass metabolism, with approximately 40% of the dose metabolized before reaching the systemic circulation. Food does not affect the rate or extent of olanzapine absorption. Pharmacokinetic studies showed that ZYPREXA tablets and ZYPREXA ZYDIS (olanzapine orally disintegrating tablets) dosage forms of olanzapine are bioequivalent.

Osage forms of oilargapine are unequaraterit.

Olanzapine displays linear kinetics over the clinical dosing range. Its half-life ranges from 21 to 54 hours (5th to 95th percentile; mean of 30 hr), nd apparent plasma clearance ranges from 12 to 47 L/hr (5th to 95th percentile; mean of 25 L/hr).

Administration of olanzapine once daily leads to steady-state concentrations in about one week that are approximately twice the concentrations fler single doses. Plasma concentrations, half-life, and clearance of olanzapine may vary between individuals on the basis of smoking status, gender, and age (see Special Populations).

gender, and age (see Special Populations).

Olanzapine is extensively distributed throughout the body, with a volume of distribution of approximately 1000 L. It is 93% bound to plasma proteins over the concentration range of 7 to 1100 ng/mL, binding primarily to albumin and  $\alpha_1$ -acid glycoprotein.

Metabolism and Elimination — Following a single oral dose of  $^{14}$ C labeled olanzapine,  $^{27}$ 6 of the dose of olanzapine was recovered in the urine as unchanged drug, indicating that olanzapine is highly metabolized. Approximately 57% and 30% of the dose was recovered in the urine and feces, respectively. In the plasma, olanzapine accounted for only 12% of the AUC for total radioactivity, indicating significant exposure to metabolites. After multiple dosing, the major circulating metabolites were the 10-N-glucuronide, present at steady state at 44% of the concentration of olanzapine, and  $^{4}$ N-desmethyl olanzapine, present at steady state at 31% of the concentration of olanzapine. Both metabolites lack pharmacological activity at the concentrations observed.

Direct glucuronidation and cytochrome P450 (CYP) mediated oxidation are the primary metabolic pathways for olanzapine. In vitro studies ggest that CYPs 142 and 256, and the flavin-containing monooxygenase system are involved in olanzapine oxidation. CYP2D6 mediated didation appears to be a minor metabolic pathway in vivo, because the clearance of olanzapine in reduced in subjects who are deficient in the enzyme

Special Populations
Renal Impairment — Because olanzapine is highly metabolized before excretion and only 7% of the drug is excreted unchanged, renal dysfunction alone is unlikely to have a major impact on the pharmacokinetics of olanzapine. The pharmacokinetic characteristics of olanzapine were similar in patients with severe renal impairment and normal subjects, indicating that dosage adjustment based upon the degree of renal impairment is not required. In addition, olanzapine is not removed by dialysis. The effect of renal impairment on metabolite elimination has not been studied. Hepatic Impairment — Although the presence of hepatic impairment may be expected to reduce the clearance of olanzapine, a study of the effect impaired liver function in subjects (n=6) with clinically significant (Childs Pugh Classification A and B) cirrhosis revealed little effect on the pharmacokinetics of olanzapine

pharmacokinetics of olanzapine.

Age — In a study involving 24 healthy subjects, the mean elimination half-life of olanzapine was about 1.5 times greater in elderly (>65 years) than in non-elderly subjects (≤65 years). Caution should be used in dosing the elderly, especially if there are other factors that might additively influence drug metabolism and/or pharmacodynamic sensitivity (see DOSAGE AND ADMINISTRATION).

Gender — Clearance of olanzapine is approximately 30% lower in women than in men. There were, however, no apparent differences between men and women in effectiveness or adverse effects. Dosage modifications based on gender should not be needed.

Smoking Status — Olanzapine clearance is about 40% higher in smokers than in nonsmokers, although dosage modifications are not routinely recommended.

recommended.

<u>Race</u> — No specific pharmacokinetic study was conducted to investigate the effects of race. A cross-study comparison between data obtained in Japan and data obtained in the US suggests that exposure to olanzapine may be about 2-fold greater in the Japanese when equivalent doses are administered. Clinical trial safety and efficacy data, however, did not suggest clinically significant differences among Caucasian patients, patients of African descent, and a third pooled category including Asian and Hispanic patients. Dosage modifications for race are, therefore, not recommended.

<u>Combined Effects</u> — The combined effects of age, smoking, and gender could lead to substantial pharmacokinetic differences in populations. The clearance in young smoking males, for example, may be 3 times higher than that in elderly nomoking fermales. Dosing modification may be necessary in patients who exhibit a combination of factors that may result in slower metabolism of olanzapine (see DOSAGE AND ADMINISTRATION).

ADMINISTRATION). For specific information about the pharmacology of lithium or valproate, refer to the CLINICAL PHARMACOLOGY section of the package inserts for these other products.

### **Clinical Efficacy Data**

Schizophrenia

The efficacy of olarzapine in the treatment of schizophrenia was established in 2 short-term (6-week) controlled trials of inpatients who met DSM III-R criteria for schizophrenia. A single haloperidol arm was included as a comparative treatment in one of the two trials, but this trial did not compare these two drugs on the full range of clinically relevant doses for both.

Several instruments were used for assessing psychiatric signs and symptoms in these studies, among them the Brief Psychiatric Rating Scale (BPRS), a multi-titem inventory of general psychopathology traditionally used to evaluate the effects of drug treatment in schizophrenia. The BPRS psychosis cluster (conceptual disorganization, hallucinatory behavior, suspiciousness, and unusual thought content) is considered a particularly useful subset for assessing actively psychotic schizophrenic patients. A second traditional assessment, the Clinical Global Impression (CGI), reflects the impression of a skilled observer, fully familiar with the manifestations of schizophrenia, about the overall clinical state of the patient, in addition, two more recently developed but less well evaluated scales were employed; these included the 30-tem Positive and Negative Symptoms Scale (PANSS), in which is embedded the 18 items of the BPRS, and the Scale for Assessing Negative Symptoms (SANS). The trial summaries below focus on the following outcomes: PANSS total and/or BPRS total; BPRS psychosis cluster; PANSS negative subscale or SANS; and GGI Severity. The results of the trials follow: SANS; and CGI Severity. The results of the trials follow:

(1) In a 6-week, placebo-controlled trial (n=149) involving two fixed olanzapine doses of 1 and 10 mg/day (once daily schedule), olanzapine, at 10 mg/day (but not at 1 mg/day), was superior to placebo on the PANSS total score (also on the extracted BPRS total), on the BPRS psychosis cluster, on the PANSS Negative subscale, and on CGI Severity.

cluster, on the PANSS Negative subscale, and on CGI Severity.

(2) In a 6-week, placebo-controlled trial (n=253) involving 3 fixed dose ranges of olanzapine (5.0 ± 2.5 mg/day, 10.0 ± 2.5 mg/day, and 15.0 ± 2.5 mg/day), and conce daily schedule, the two highest olanzapine dose groups (actual mean doses of 12 and 16 mg/day, respectively) were superior to placebo on BPRS total score, BPRS psychosis cluster, and CGI severity score; the highest olanzapine dose group was superior to placebo on the SANS. There was no clear advantage for the high dose group over the medium dose group.

Examination of population subsets (race and gender) did not reveal any differential responsiveness on the basis of these subgroupings.

In a longer-term trial, adult outpatients (n=326) who predominantly met DSM-IV criteria for schizophrenia and who remained stable on olanzapine doses (ranging from 10 to 20 mg/day) or to placebo. The follow-up period to observe patients for relapse, defined in terms of increases in BPRS positive symptoms or hospitalization, was planned for 12 months, however, criteria were met for stopping the trial early due to an excess of placebo relapses compared to olanzapine relapses, and olanzapine was superior to placebo on time to relapse, the primary outcome for this study. Thus, olanzapine was more effective than placebo at maintaining efficacy in patients stabilized for approximately 8 weeks and followed for an observation period of up to 8 months.

Bioolar Disorder

Bipolar Disorder

Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes was established in 2 short-term (one 3-week and Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes was established in 2 short-term (one 3-week and Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes was established in 2 short-term (one 3-week and Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes was established in 2 short-term (one 3-week and Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes was established in 2 short-term (one 3-week and Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes was established in 2 short-term (one 3-week and Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes was established in 2 short-term (one 3-week and Monotherapy — The efficacy of olanzapine in the treatment of acute manic or mixed episodes. The efficacy of olanzapine in the treatment of acute manic or mixed episodes. one 4-week) placebo-controlled trials in patients who met the DSM-IV criteria for Bipolar I Disorder with manic or mixed episodes. These trials included patients with or without psychotic features and with or without a rapid-cycling course.

The primary rating instrument used for assessing manic symptoms in these trials was the Young Mania Rating Scale (Y-MRS), an 11-tien clinician-rated scale traditionally used to assess the degree of manic symptomatology (irritability, disruptive/aggressive behavior, sleep, elevated mood, speech, increased activity, sexual interest, language/thought disorder, thought content, appearance, and insight) in a range from 0 (no manic features) to 60 (maximum score). The primary outcome in these trials was change from baseline in the Y-MRS total score. The results of the trials follow:

(1) In one 3-week placebo-controlled trial (n=67) which involved a dose range of olanzapine (5-20 mg/day, once daily, starting at 10 m olanzapine was superior to placebo in the reduction of Y-MRS total score. In an identically designed trial conducted simultaneously with the fi olanzapine demonstrated a similar treatment difference, but possibly due to sample size and site variability, was not shown to be superior to proceed to the conduction of the conducti (2) In a 4-week placebo-controlled trial (n=115) which involved a dose range of olanzapine (5-20 mg/day, once daily, starting at 15 mg/day),

rior to placebo in the reducti on of Y-MRS total score. olanzapine was superior to placebo in the reduction of Y-MRS total score.

(3) In another trial, 361 patients meeting DSM-IV criteria for a manic or mixed episode of bipolar disorder who had responded during an initial open-label treatment phase for about two weeks, on average, to olanzapine 5 to 20 mg/day were randomized to either continuation of olanzapine at their same dose (n=225) or to placebo (n=136), for observation of relapse. Approximately 50% of the patients had discontinued from the olanzapine group by day 59 and 50% of the placebo group had discontinued by day 23 of double-blind treatment. Response during the open-label phase was defined by having a decrease of the Y-MRS total score to ≤12 and HAM-D 21 to ≤8. Relapse during the double-blind phase was defined as an increase of the Y-MRS or HAM-D 21 total score to ≤15, or being hospitalized for either mania or depression. In the randomized phase, patients receiving continued olanzapine experienced a significantly longer time to relapse.

Combination Therapy — The efficacy of olanzapine with concomitant lithium or valproate in the treatment of acute manic episodes was established in two controlled trials in patients who met the DSM-IV criteria for Bipolar I Disorder with manic or mixed episodes. These trials included related to the proper of the prope

established in two controlled trials in patients who met the DSM-IV criteria for Bipolar I Disorder with manic or mixed episodes. These trials included patients with or without psychotic features and with or without a rapid-cycling course. The results for trials follow:

(1) In one 6-week placebo-controlled combination trial, 175 outpatients on lithium or valproate therapy with inadequately controlled manic or mixed symptoms (Y-MRS 216) were randomized to receive either olanzapine or placebo, in combination with their original therapy. Olanzapine (in a dose range of 5-20 mg/day, once daily, starting at 10 mg/day) combined with lithium or valproate (in a therapeutic range of 0.6 mEq/L to 1.2 mEq/L or 50 µg/mL to 125 µg/mL, respectively) was superior to lithium or valproate alone in the reduction of Y-MRS total score.

(2) In a second 6-week placebo-controlled combination trial, 169 outpatients on lithium or valproate therapy with inadequately controlled manic or mixed symptoms (Y-MRS ≥16) were randomized to receive either olanzapine or placebo, in combination with their original therapy. Olanzapine (in a dose range of 5-20 mg/day, once daily, starting at 10 mg/day) combined with lithium or valproate (in a therapeutic range of 0.6 mEq/L to 1.2 mEq/L or 50 µg/mL to 125 µg/mL, respectively) was superior to lithium or valproate alone in the reduction of Y-MRS total score. INDICATIONS AND USAGE

# ZYPREXA is indicated for the treatment of schizophrenia.

The efficacy of ZYPREXA was established in short-term (6-week) controlled trials of schizophrenic inpatients (see CLINICAL PHARMACOLOGY) The effectiveness of oral ZYPREXA at maintaining a treatment response in schizophrenic patients who had been stable on ZYPREXA for approximately 8 weeks and were then followed for a period of up to 8 months has been demonstrated in a placebo-controlled trial (see CLINICAL PHARMACOLOGY). Nevertheless, the physician who elects to use ZYPREXA for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient (see DOSAGE AND ADMINISTRATION).

Acute Monotherapy — ZYPREXA is indicated for the treatment of acute mixed or manic episodes associated with Bipolar I Disorde

ACUSE MONOMERS — 2YPHEXA is indicated for the treatment of acuse mixed or maintc episodes associated with sippliar 1 bisorder. The efficiency of ZYPREXA was established in two placebo-controlled trials (one 3-week and one 4-week) with patients meeting DSM-IV criteria for Bipolar I Disorder who currently displayed an acute manic or mixed episode with or without psychotic features (see CLINICAL PHARIMACOLOGY). Maintenance Monotherapy — The benefit of maintaining bipolar patients on monotherapy with ZYPREXA after achieving a responder status for an average duration of two weeks was demonstrated in a controlled trial (see Clinical Efficacy Data *under* CLINICAL PHARIMACOLOGY). The physician who elects to use ZYPREXA for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient (see DOSAGE AND ADMINISTRATION). Combination Therapy — The combination of ZYPREXA with lithium or valproate is indicated for the short-term treatment of acute manic episodes sociated with Bipolar | Disorder.

The efficacy of ZYPREXA in combination with lithium or valproate was established in two placebo-controlled (6-week) trials with patients meeting DSM-IV criteria for Bipolar I Disorder who currently displayed an acute manic or mixed episode with or without psychotic features (see CLINICAL PHARMACOLOGY). CONTRAINDICATIONS

# ZYPREXA is contraindicated in patients with a known hypersensitivity to the product

For specific information about the contraindications of lithium or valproate, refer to the CONTRAINDICATIONS section of the package inserts for

WARNINGS

Hyperglycemia and Diabetes Mellitus — Hyperglycemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics including olanzapine. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with exclusional patients and the increasaing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycemia-related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent hyperglycemia-related adverse events in patients treated with the atypical antipsychotics. Precise risk estimates for hyperglycemia-related adverse events in patients treated with the atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus who are started on applical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g., obesity, family history of diabetes) who are startment Any patient treated with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polypraja, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic was one patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug.

Cerebrovascular Adverse Events, Including Stroke, in Elderly Patients with Dementia — Cerebrovascular adverse event

Cerebrovascular Adverse Events, Including Stroke, in Elderly Patients with Dementia — Cerebrovascular adverse events (e.g., stroke, transic ischemic attack), including fatalities, were reported in patients in trials of olanzapine in elderly patients with dementia-related psychosis, placebo-controlled trials, there was a significantly higher incidence of cerebrovascular adverse events in patients treated with olanzapine compar to patients treated with placebo. Olanzapine is not approved for the treatment of patients with dementia-related psychosis. Neuroleptic Malignant Syndrome (NMS) — A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with administration of antipsychotic drugs, including olanzapine. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status and evidence of autonomic instability (irregular pulse or bloor pressure, tachycardia, diaphoresis and cardiac dysrhythmia). Additional signs may include elevated creatinine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure.

oysrifynma). Additional signs may include elevated creatinine phosphokinase, myoglobinuna (mabdomyojishs), and acute renal failure.

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include: 1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy; 2) intensive symptomatic treatment and medical monitoring; and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, the potential retroduction of drug therapy should be carefully considered. The patient should be carefully monitored, since recurrences of NMS have been reported.

Tardive Dyskinesia — A syndrome of potentially irreversible, involuntary, dyskinetic movements may develop in patients treated with antipsychotic drugs. Although the prevalence of the syndrome appears to be highest among the elderly, especially elderly women, it is impossible to rely upon prevalence estimates to predict, at the inception of antipsychotic traument, which patients are likely to develop the syndrome. Whether antipsychotic drug products differ in their potential to cause tardive dyskinesia is unknown. The risk of developing tardive dyskinesia and the likelihood that it will become irreversible are believed to increase as the duration of trea

and the total cumulative dose of antipsychotic drugs administered to the patient increase. However, the syndrome can develop, although much less commonly, after relatively brief treatment periods at low doses. There is no known treatment for established cases of tardive dyskinesia, although the syndrome may remit, partially or completely, if antipsychotic treatment is withdrawn. Antipsychotic treatment, itself, however, may suppress (or partially suppress) the signs and symptoms of the syndrome and thereby may possibly mask the underlying process. The effect that symptomatic suppression has upon the long-term course of the syndrome is unknown.

syndrome is unknown.

Given these considerations, olanzapine should be prescribed in a manner that is most likely to minimize the occurrence of tardive dyskinesia.

Chronic antipsychotic treatment should generally be reserved for patients (1) who suffer from a chronic illness that is known to respond to antipsychotic drugs, and (2) for whom alternative, equally effective, but potentially less harmful treatments are not available or appropriate. In patients who do require chronic treatment, the smallest dose and the shortest duration of treatment producing a satisfactory clinical responses should be sought. The need for continued treatment should be reassessed periodically.

If signs and symptoms of tardive dyskinesia appear in a patient on olanzapine, drug discontinuation should be considered. However, some patients may require treatment with olanzapine despite the presence of the syndrome. For specific information about the warnings of lithium or valproate, refer to the WARNINGS section of the package inserts for these

PRECAUTIONS

General

Orthostatic Hypotension — Olanzapine may induce orthostatic hypotension associated with dizziness, tachycardia, and in some patients, syncope, especially during the initial dose-litration period, probably reflecting its α<sub>1</sub>-adrenergic antagonistic properties. Syncope was reported in 0.6% (15/2500) of olanzapine-treated patients in phase 2-3 studies. The risk of orthostatic hypotension and syncope may be minimized by initiating therapy with 5 mg QD (see DOSAGE AND ADMINISTRATION). A more gradual titration to the target dose should be considered if hypotension occurs. Olanzapine should be used with particular caution in patients with known cardiovascular disease (history of myocardial infarction or

# $ZYPREXA^{\circledR} \ (Olanzapine) \ Tablets \\ ZYPREXA^{\circledR} \ ZYDIS^{\circledR} \ (Olanzapine) \ Orally \ Disintegrating \ Tablets$

ischemia, heart failure, or conduction abnormalities), cerebrovascular disease, and conditions which would predispose patients to hypotension (dehydration, hypovolemia, and treatment with antihypertensive medications).

Seizures — During premarketing testing, seizures occurred in 0.9% (20/2500) of olanzapine-treated patients. There were confounding factors that may have contributed to the occurrence of seizures in many of these cases. Olanzapine should be used cautiously in patients with a history of seizures or with conditions that potentially lower the seizure threshold, e.g., Alzheimer's dementia. Conditions that lower the seizure threshold may be more prevalent in a population of 65 years or older.

be more prevalent in a population of 65 years or older. 

Hyperprolactinemia — As with other drugs that antagonize dopamine D<sub>2</sub> receptors, olanzapine elevates prolactin levels, and a modest elevation persists during chronic administration. Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin dependent in vitro, a factor of potential importance if the prescription of these drugs is contemplated in a patient with previously detected breast cancer of this type. Although disturbances such as galactorrhea, amenorrhea, gynecomastia, and impotence have been reported with prolactin-elevating compounds, the clinical significance of elevated serum prolactin levels is unknown for most patients. As is common with compounds which increase prolactin release, an increase in mammary gland neoplasia was observed in the olanzapine carcinogenicity studies conducted in mice and rats (see Carcinogenesis). However, neither clinical studies nor epidemiologic studies have shown an association between chronic administration of this class of drugs and tumorigenesis in humans; the available evidence is considered too limited to be conclusive.

Transaminase Elevations — In placeho-controlled studies, clinically significant ALT (SQPT) elevations (S) times the unper limit of the normal

conducted in mice and rats (see Carcinogenesis). However, neither clinical studies nor epidemiologic studies have shown an association between chronic administration of this class of drugs and tumorigenesis in humans; the available evidence is considered too limited to be conclusive.

Transaminase Elevations — In placebo-controlled studies, clinically significant ALT (SGPT) elevations (28 times the upper limit of the normal range) were observed in 2% (6/243) of patients exposed to olanzapine compared to none (0/115) of the placebo patients. None of these patients experienced jaundice. In two of these patients, liver enzymes decreased toward normal despite continued treatment and in two others, enzymes decreased upon discontinuation of olanzapine. In the remaining two patients, one, seropositive for hepatilis C, had persistent enzyme elevation for four months after discontinuation, and the other had insufficient follow-up to determine if enzymes normalized.

Within the larger premarketing database of about 2400 patients with baseline SGPT 590 IU/L, the incidence of SGPT elevation to >200 IU/L was \$\circ\$ (50/2381). Again, none of these patients experienced jaundice or other symptoms attributable to liver impairment and most had transient changes that tended to normalize while olanzapine treatment was continued.

Among all 2500 patients in clinical trials, about 1% (23/2500) discontinued treatment due to transaminase increases.

Caution should be exercised in patients with signs and symptoms of hepatic impairment, in patients with pre-existing conditions associated with limited hepatic functional reserve, and in patients with significant hepatic disease (see Laboratory Tests).

Potential for Cognitive and Motor Impairment — Somnolence was a commonly reported adverse event associated with olanzapine treatment, occurring at an incidence of 26% in olanzapine patients compared to 15% in placebo patients. This adverse event was also dose related. Somnolence led to discontinuation in 0.4% (9/2500) of patients in the pr

being subject to dehydration.

<u>Dysphagia</u> — Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Two olanzapine-treated patients (2/407) in two studies in patients with Alzheimer's disease died from aspiration pneumonia during or within 30 days of the termination of the double-blind portion of their respective studies; there were no deaths in the placebo-treated patients. One of these patients had experienced dysphagia prior to the development of aspiration pneumonia. Aspiration pneumonia is a common cause of morbidity and mortality in patients with advanced Alzheimer's disease. Olanzapine and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia.

<u>Suicide</u> — The possibility of a suicide attempt is inherent in schizophrenia and in bipolar disorder, and close supervision of high-risk patients should accompany drug therapy. Prescriptions for olanzapine should be written for the smallest quantity of tablets consistent with good patient management, in order to reduce the risk of overdose.

Use in Patients with Concomitant Illness — Clinical experience with olanzapine in patients with certain concomitant systemic illnesses (see Rena pairment and Hepatic Impairment under CLINICAL PHARMACOLOGY, Special Populations) is limited.

Olanzapine exhibits in vitro muscarinic receptor affinity. In premarketing clinical trials with olanzapine, olanzapine was associated with nestipation, dry mouth, and tachycardia, all adverse events possibly related to cholinergic antagonism. Such adverse events were not often the basis for discontinuations from olanzapine, but olanzapine should be used with caution in patients with clinically significant prostatic hypertrophy,

narrow angle glaucoma, or a history of paralytic ileus.

In a fixed-dose study of olanzapine (olanzapine at doses of 5, 10, and 15 mg/day) and placebo in nursing home patients (mean age: 83 years, range: 61-97; median Mini-Mental State Examination (MMSE): 5, range: 0-22) having various psychiatric symptoms in association with Alzheimer's disease, the following treatment-emergent adverse events were reported in all (each and every) olanzapine-treated groups at an incidence of either (1) two-fold or more in excess of the placebo-treated group, where at least 1 placebo-treated patient was reported to have experienced the event; or (2) at least 2 cases if no placebo-treated patient was reported to have experienced the event: somolence, abnormal gait, fever, dehydration, and back pain. The rate of discontinuation in this study for olanzapine was 12% vs 4% with placebo. Discontinuations due to abnormal gait (1% for olanzapine vs 0% for placebo), accidental injury (1% for olanzapine vs 0% for placebo), and somnolence (3% to olanzapine vs 0% for placebo) were considered to be drug related. As with other CNS-active drugs, olanzapine should be used with caution in elderly patients with dementia (see PRECAUTIONS). narrow angle glaucoma, or a history of paralytic ileus.

Olanzapine has not been evaluated or used to any appreciable extent in patients with a recent history of myocardial infarction or unstable heart disease. Patients with these diagnoses were excluded from premarketing clinical studies. Because of the risk of orthostatic hypotension with olanzapine, caution should be observed in cardiac patients (see Orthostatic Hypotension).

For specific information about the precautions of lithium or valproate, refer to the PRECAUTIONS section of the package inserts for these

Information for Patients

Physicians are advised to discuss the following issues with patients for whom they prescribe olanzapine:

Orthostatic Hypotension — Patients should be advised of the risk of orthostatic hypotension, especially during the period of initial dose titration and in association with the use of concomitant drugs that may potentiate the orthostatic effect of olanzapine, e.g., diazepam or alcohol (see Drug Interactions).

Interference with Cognitive and Motor Performance — Because olanzapine has the potential to impair judgment, thinking, or motor skills, patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that olanzapine therapy does not affect them adversely.

Pregnancy — Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during therapy with olanzapine. Nursing — Patients should be advised not to breast-feed an infant if they are taking olanzapine.

Concomitant Medication — Patients should be advised to inform their physicians if they are taking, or plan to take, any prescription or over-the-counter drugs, since there is a potential for interactions. Alcohol — Patients should be advised to avoid alcohol while taking olanzapine

Heat Exposure and Dehydration — Patients should be advised regarding appropriate care in avoiding overheating and dehydration.

Phenylketonurics — ZYPREXA ZYDIS (olanzapine orally disintegrating tablets) contains phenylalanine (0.34, 0.45, 0.67, or 0.90 mg per 5, 10, 15, or 20 mg tablet, respectively).

Laboratory Tests
Periodic assessment of transaminases is recommended in patients with significant hepatic disease (see Transaminase Elevations).

The risks of using olanzapine in combination with other drugs have not been extensively evaluated in systematic studies. Given the primary CNS effects of olanzapine, caution should be used when olanzapine is taken in combination with other centrally acting drugs and alcohol.

Because of its potential for inducing hypotension, olanzapine may enhance the effects of certain antihypertensive agents.

Olanzapine may antagonize the effects of levodopa and dopamine agonists. Olanzapine may antagonize the effects of levodopa and dopamine agonists.

The Effect of Other Drugs on Olanzapine — Agents that induce CYP1A2 or glucuronyl transferase enzymes, such as omeprazole and rifampin, may cause an increase in olanzapine clearance. Inhibitors of CYP1A2 could potentially inhibit olanzapine clearance. Although olanzapine is metabolized by multiple enzyme systems, induction or inhibition of a single enzyme may appreciably alter olanzapine clearance. Therefore, a dosage increase (for induction) or a dosage decrease (for inhibition) may need to be considered with specific drugs.

Charcoal — The administration of activated charcoal (1 g) reduced the Cmax and AUC of olanzapine by about 60%. As peak olanzapine levels are not typically obtained until about 6 hours after dosing, charcoal may be a useful treatment for olanzapine overdose.

Cimetidine and Antacids — Single doses of cimetidine (800 mg) or aluminum- and magnesium-containing antacids did not affect the oral bioavailability of olanzapine.

Carbamazeoine — Carbamazeoine therapy (200 mg bid) causes an approximately 50% increase in the clearance of olanzapine. This increase is

increase in olanzapine clearance.

Ethanol — Ethanol (45 mg/70 kg single dose) did not have an effect on olanzapine pharmacokinetics.

Fluoxetine — Fluoxetine (60 mg single dose or 60 mg daily for 8 days) causes a small (mean 16%) increase in the maximum concentration of olanzapine and a small (mean 16%) decrease in olanzapine clearance. The magnitude of the impact of this factor is small in comparison to the overall variability between individuals, and therefore dose modification is not routinely recommended.

Fluoxamine — Fluoxamine a CYP1A2 inhibitor, decreases the clearance of olanzapine. This results in a mean increase in olanzapine Cmax following fluovamine of 54% in female nonsmokers and 77% in male smokers. The mean increase in olanzapine AUC is 52% and 108%, respectively. Lower doses of olanzapine should be considered in patients receiving concomitant treatment with fluovamine.

Warfarin — Warfarin (20 mg single dose) did not affect olanzapine pharmacokinetics.

Fiffect of Olanzapine on Other Druos. — In vitro studies utilizing human liver microsomes suggest that olanzapine has little potential to inhibit

Effect of Olanzapine on Other Drugs — In vitro studies utilizing human liver microson CYP1A2, CYP2C9, CYP2C19, CYP2D6, and CYP3A. Thus, olanzapine is unlikely to c

<u>Lithium</u> — Multiple doses of olanzapine (10 mg for 8 days) did not influence the kinetics of lithium. Therefore, concomitant ola administration does not require dosage adjustment of lithium. <u>Valproate</u> — Studies in vitro using human liver microsomes determined that olanzapine has little potential to inhibit the major metabolic path glucuronidation, of alproate. Further, valproate has little effect on the metabolism of olanzapine in vitro. In vivo administration of olanzapine (1) daily for 2 weeks) did not affect the steady state plasma concentrations of valproate. Therefore, concomitant olanzapine administration does

require dosage adjustment of valproate Single doses of olanzapine did not affect the pharmacokinetics of imipramine or its active metabolite desipramine, and warfarin. Multiple doses of olanzapine did not influence the kinetics of diazepam and its active metabolite N-desmethyldiazepam, ethanol, or biperiden. However, the co-administration of either diazepam or ethanol with olanzapine potentiated the orthostatic hypotension observed with olanzapine. Multiple doses of olanzapine did not affect the pharmacokinetics of theophylline or its metabolites.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis — Oral carcinogenicity studies were conducted in mice and rats. Olanzapine was administered to mice in two 78-week studies at doses of 3, 10, 30/20 mg/kg/day (equivalent to 0.8-5 times the maximum recommended human daily dose on a mg/m² basis) and 0.25, 2, 8 mg/kg/day (equivalent to 0.6-2 times the maximum recommended human daily dose on a mg/m² basis). Atts were dosed for 2 years at doses of 0.25, 1, 2.5, 4 mg/kg/day (males) and 0.25, 1, 4, 8 mg/kg/day (females) (equivalent to 0.13-2 and 0.13-4 times the maximum recommended human daily dose on a mg/m² basis). The incidence of liver hemangiomas and hemangiosarcomas was significantly increased in one mouse study in female mice dosed at 8 mg/kg/day (2 times the maximum recommended human daily dose on a mg/m² basis). These tumors were not increased in another mouse study in females dosed at 10 or 30/20 mg/kg/day (2-5 times the maximum recommended human daily dose on a mg/m² basis). The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group. The incidence of early mortalities in males of the 30/20 mg/kg/day group increased in anome mouse study in remains occessed at 100 x0x20 migrygray (225 times the instantion recommended interial dualy dose on a migrim-basis); in this study, there was a high incidence of early mortalities in males of the 50/20 mg/kg/day group. The incidence of mammary gland adenomas and adenocarcinomas was significantly increased in female mice dosed at 22 mg/kg/day and in female rats dosed at 24 mg/kg/day (0.5 and 2 times the maximum recommended human daily dose on a mg/m² basis, respectively). Antipsychotic drugs have been shown to chronically elevate prolactin levels in rodents. Serum prolactin levels were not measured during the olanzapine carcinogenicity studies; however, measurements during subchronic toxicity studies showed that olanzapine elevated serum prolactin levels up to 4-10d in rats at the same doses used in the carcinogenicity study. An increase in mammary gland neoplasms has been found in rodents after chronic administration of other antipsychotic drugs and is considered to be prolactin mediated. The relevance for human risk of the finding of prolactin mediated endocrine tumors in rodents is unknown (see Hyperprolactinemia under PRECAUTIONS, General).

unknown (see ryperprolactinemia under PHECAUTIONS, General).

Mutagenesis — No evidence of mutagenic potential for olanzapine was found in the Ames reverse mutation test, in vivo micronucleus test in mice, the chromosomal aberration test in Chinese hamster ovary cells, unscheduled DNA synthesis test in rat hepatocytes, induction of forward mutation test in mouse lymphoma cells, or in vivo sister chromatid exchange test in bone marrow of Chinese hamsters.

Impairment of Fertility — In a fertility and reproductive performance study in rats, male mating performance, but not fertility, was impaired at a dose of 22.4 mg/kg/day and female fertility was decreased at a dose of 3 mg/kg/day (11 and 1.5 times the maximum recommended human daily dose on a mg/m² basis, respectively). Discontinuance of olanzapine treatment reversed the effects on male mating performance. In female rats, the precoilal period was increased and the mating index reduced at 5 mg/kg/day (2.5 times the maximum recommended human daily dose on a mg/m² basis). Diestrous was prolonged and estrous delayed at 1.1 mg/kg/day (0.6 times the maximum recommended human daily dose on a mg/m² basis); therefore olanzapine may produce a delayed at 1.1 mg/kg/day (0.6 times the maximum recommended human daily dose on a mg/m² basis); therefore olanzapine may produce a delayed at 1.1 mg/kg/day (0.6 times the maximum recommended human daily dose on a mg/m² basis); therefore olanzapine may produce a delayed at 1.1 mg/kg/day (0.6 times the maximum recommended human daily dose on a mg/m² basis); therefore olanzapine may produce a delay in ovulation.

Pregnancy
Pregnancy
Pregnancy
Category C — In reproduction studies in rats at doses up to 18 mg/kg/day and in rabbits at doses up to 30 mg/kg/day (9 and 30 times the maximum recommended human daily dose on a mg/m² basis, respectively) no evidence of teratogenicity was observed. In a rat teratology study, early resorptions and increased numbers of nonviable fetuses were observed at a dose of 18 mg/kg/day (9 times the maximum recommended human daily dose on a mg/m² basis). Gestation was prolonged at 10 mg/kg/day (6 times the maximum recommended human daily dose on a mg/m² basis). In a rabbit teratology study, fetal toxicity (manifested as increased resorptions and decreased fetal weight) occurred at a maternally toxic dose of 30 mg/kg/day (30 times the maximum recommended human daily dose on a mg/m² basis).
Placental transfer of olanzapine occurs in rat pups.
There are no adequate and well-controlled trials with olanzapine in pregnant females. Seven pregnancies were observed during clinical trials with olanzapine, including 2 resulting in normal births, 1 resulting in neonatal death due to a cardiovascular defect, 3 therapeutic abortions, and 1 spontaneous abortion. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Labor and Delivery
Parturition in rats was not affected by olanzapine. The effect of olanzapine on labor and delivery in humans is unknown.

Olanzapine was excreted in milk of treated rats during lactation. It is not known if olanzapine is excreted in human milk. It is recommended that romen receiving olanzapine should not breast-feed.

Pediatric Use Safety and effectiveness in pediatric patients have not been established.

Genatire Use
Of the 2500 patients in premarketing clinical studies with olanzapine, 11% (263) were 65 years of age or over. In patients with schizophrenia,
there was no indication of any different tolerability of olanzapine in the elderly compared to younger patients. Studies in patients with various
psychiatric symptoms in association with Alzheimer's disease have suggested that there may be a different tolerability profile in this population
compared to younger patients with schizophrenia. As with other CNS-active drugs, olanzapine should be used with caution in elderly patients with
dementia. Also, the presence of factors that might decrease pharmacokinetic clearance or increase the pharmacodynamic response to olanzapine
should lead to consideration of a lower starting dose for any geriatric patient (see PRECAUTIONS and DOSAGE AND ADMINISTRATION). ADVERSE REACTIONS

The information below is derived from a clinical trial database for olanzapine consisting of 8661 patients with approximately 4165 patient-years of exposure. This database includes: (1) 2500 patients who participated in multiple-dose premarketing trials in schizophrenia and Alzheimer's disease representing approximately 1122 patient-years of exposure as of February 14, 1995; (2) 182 patients who participated in premarketing bipolar maint trials representing approximately (6) patient-years of exposure; (3) 191 patients who participated in a trial of patients having various psychiatric symptoms in association with Alzheimer's disease representing approximately (29 patient-years of exposure; and (4) 5788 patients from 88 additional clinical trials as of December 31, 2001. In addition, information from the premarketing 6-week clinical study database for olanzapine in combination with lithium or valproate, consisting of 224 patients who participated in bipolar mania trials with approximately 22 patient-years of exposure, is included below.

The conditions and duration of treatment with olanzapine varied greatly and included (in overlapping categories) open-label and double-blind nases of studies, inpatients and outpatients, fixed-dose and dose-titration studies, and short-term or longer-term exposure. Adverse reactions were issessed by collecting adverse events, results of physical examinations, vital signs, weights, laboratory analytes, ECGs, chest x-rays, and results of withalmologic examinations.

ophthalmologic examinations.

Certain portions of the discussion below relating to objective or numeric safety parameters, namely, dose-dependent adverse events, vital sign changes, weight gain, laboratory changes, and ECG changes are derived from studies in patients with schizophrenia and have not been duplicated for bipolar mania. However, this information is also generally applicable to bipolar mania.

Adverse events during exposure were obtained by spontaneous report and recorded by clinical investigators using terminology of their own choosing. Consequently, it is not possible to provide a meaningful estimate of the proportion of individuals experiencing adverse events without first grouping similar types of events into a smaller number of standardized event categories. In the tables and tabulations that follow, standard COCTADT distinguary terminology of the proportion of individuals experiencing adverse events without first grouping similar types of events into a smaller number of standardized event categories. In the tables and tabulations that follow, standard

grouping similar types of events into a smaller number of standardized event categories COSTART dictionary terminology has been used initially to classify reported adverse events. The stated frequencies of adverse events represent the proportion of individuals who experienced, at least once, a treatment-emergent adverse

The stated frequencies of adverse events represent the proportion of individuals who experienced, at least once, a treatment-emergent adverse event of the type listed. An event was considered treatment emergent if it occurred for the first time or worsened while receiving therapy following baseline evaluation. The reported events do not include those event terms which were so general as to be uninformative. Events listed elsewhere in labeling may not be repeated below. It is important to emphasize that, although the events occurred during treatment with olanzapine, they were not necessarily caused by it. The entire label should be read to gain a complete understanding of the safety profile of olanzapine. The prescriber should be awarent that the figures in the tables and tabulations cannot be used to predict the incidence of side effects in the course of usual medical practice where patient characteristics and other factors differ from those that prevailed in the clinical trials. Similarly, the cited frequencies cannot be compared with figures obtained from other clinical investigations involving different treatments, uses, and investigators. The cited figures, however, do provide the prescribing physician with some basis for estimating the relative contribution of drug and nondrug factors to the adverse event incidence in the population studied.

Incidence of Adverse Events in Short-Term, Placebo-Controlled and Combination Trials
The following findings are based on premarketing trials for schizophrenia, bipolar mania, a subsequent trial of patients having various psychiatric symptoms in association with Alzheimer's disease, and premarketing combination trials. symptoms in association with Alzheimer's disease, and premarked your continuous account of the symptoms of the

Bipolar Mania Monotherapy — Overall, there was no difference in the incidence of discontinuation due to adverse events (2% for olanzapine vs 2% for placebo). PRECAUTIONS

Adverse Events Associated with Discontinuation of Treatment in Short-Term Combination Trials Adverse Events Asia Scotated with Discontinuation of Treatment in Stort-Term Combination and Treatment in Stort-Term Combination and Treatment in Stort-Term Combination and Treatment in Stort-Term Stort Institute or valproate as monotherapy, discontinuation rates due to adverse events were 11% for the combination of clanzapier of whitlinitium or valproate compared to 2% for predict more remained on lithium or valproate monotherapy. Discontinuations with the combination of clanzapier and lithium or valproate that occurred in more

than 1 patient were: somnolence (3%), weight gain (1%), and peripheral edema (1%). Commonly Observed Adverse Events in Short-Term, Placebo-Controlled Trials
The most commonly observed adverse events associated with the use of olanzapine (incidence of 5% or greater) and not observed at an equivalent incidence among placebo-treated patients (olanzapine incidence at least twice that for placebo) were:

	reatment-Emergent Adverse Events Associated Olanzapine in 6-Week Trials — SCHIZOPHRE	NIA
	Percentage of Patie	nts Reporting Event
Adverse Event	Olanzapine (N=248)	Placebo (N=118)
Postural hypotension	5	2
Constipation	9	3
Weight gain	6	1
Dizziness	11	4
Personality disorder <sup>1</sup>	8	4
Akathisia	5	1

## ZYPREXA® (Olanzapine) Tablets ZYPREXA® ZYDIS® (Olanzapine) Orally Disintegrating Tablets

Common Treatment-Emergent Adverse Events Associated with the Use of Olanzapine in 3-Week and 4-Week Trials — BIPOLAR MANIA						
	Percentage of Patients Reporting Event					
Adverse Event	Olanzapine (N=125)	Placebo (N=129)				
Asthenia	15	6				
Dry mouth	22	7				
Constipation	11	5				
Dyspepsia	11	5				
Increased appetite	6	3				
Somnolence	35	13				
Dizziness	18	6				
Tremor	6	3				

erse Events Occurring at an Incidence of 2% or More Among Olanzapine-Treated Patients in Short-Term, Placebo-Controlled Trials able 1 enumerates the incidence, rounded to the nearest percent, of treatment-emergent adverse events that occurred in 2% or more of patients ed with olanzapine (doses ≥2.5 mg/day) and with incidence greater than placebo who participated in the acute phase of placebo-controlled trials.

# Table 1 Treatment-Emergent Adverse Events: Incidence in Short-Term, Placebo-Controlled Clinical Trials<sup>1</sup>

	Percentage of Patients Reporting Event		
-	Olanzapine	Placebo	
Body System/Adverse Event	(N=532)	(N=294)	
Body as a Whole	12	8	
Accidental injury Asthenia	12	9	
Fever	6	2	
Back pain	5	2	
Chest pain	3	1	
Cardiovascular System			
Postural hypotension	3	1	
Tachycardia	3	1	
Hypertension	2	1	
Digestive System		_	
Dry mouth	9	5	
Constipation	9	4	
Dyspepsia	7	5	
Vomiting	4	3	
Increased appetite	3	2	
Hemic and Lymphatic System			
Ecchymosis	5	3	
Metabolic and Nutritional Disorders			
Weight gain	5	3	
Peripheral edema	3	1	
Musculoskeletal System			
Extremity pain (other than joint)	5	3	
Joint pain	5	3	
Nervous System			
Somnolence	29	13	
Insomnia	12	11	
Dizziness	11	4	
Abnormal gait	6	1	
Tremor	4	3	
Akathisia	3	2	
Hypertonia	3	2	
Articulation impairment	2	1	
Respiratory System			
Rhinitis	7	6	
Cough increased	6	3	
Pharyngitis	4	3	
Special Senses			
Amblyopia	3	2	
Urogenital System			
Urinary incontinence	2	1	
Urinary tract infection	2	1	

Events reported by at least 2% of patients treated with olanzapine, except the following ev pain, agilation, anorexia, anxiety, apathy, contusion, depression, diarrhea, dysmenorrhe nervousness, paranioti reaction, personality disorder<sup>3</sup>, rash, thinking athorama, weight loss Denominator used was for females only (olanzapine, N=201; placebo, N=114). Personality disorder is the COSTART term for designating non-aggressive objectionable be ving events which had an incidence equal to or less than placebo: abdomina norrhea<sup>2</sup>, hallucinations, headache, hostility, hyperkinesia, myalgia, nausea

Commonly Observed Adverse Events in Short-Term Combination Trials

In the bipolar mania combination placebo-controlled trials, the most commonly observed adverse events associated with the combination of olanzapine and lithium or valproate (incidence of ≥5% and at least twice placebo) were:

	on Treatment-Emergent Adverse Events Associate Ilanzapine in 6-Week Combination Trials — BIPOL	
	Percentage of Patie	ents Reporting Event
Adverse Event	Olanzapine with lithium or valproate (N=229)	Placebo with lithium or valproate (N=115)
Dry mouth	32	9
Weight gain	26	7
Increased appetite	24	8
Dizziness	14	7
Back pain	8	4
Constipation	8	4
Speech disorder	7	1
Increased salivation	6	2
Amnesia	5	2
Paresthesia	5	2

Adverse Events Occurring at an Incidence of 2% or More Among Olanzapine-Treated Patients in Short-Term Combination Trials

Table 2 enumerates the incidence, rounded to the nearest percent, of treatment-emergent adverse events that occurred in 2% or more of patients treated with the combination of olanzapine (doses ≥5 mg/day) and lithium or valproate and with incidence greater than lithium or valproate alone who participated in the acute phase of placebo-controlled combination trials.

# Table 2

	Percentage of Patients Reporting Event		
Body System/Adverse Event	Olanzapine with lithium or valproate (N=229)	Placebo with lithium or valproate (N=115)	
Body as a Whole	(N=229)	(N=115)	
Asthenia	18	13	
Back pain	8	4	
Accidental injury	4	2	
Chest pain	3	2	
Cardiovascular System	3		
	0		
Hypertension	2	1	
Digestive System			
Ory mouth	32	9	
ncreased appetite	24	8	
Thirst	10	6	
Constipation	8	4	
ncreased salivation	6	2	
Metabolic and Nutritional Disorders			
Veight gain	26	7	
Peripheral edema	6	4	
Edema	2	1	
Vervous System			
comnolence	52	27	
remor	23	13	
Depression	18	17	
vizziness	14	7	
Speech disorder	7	1	
mnesia	5	2	
Paresthesia	5	2	
Apathy	4	3	
Confusion	4	1	
Euphoria	3	2	
ncoordination	2	0	
Respiratory System			
Pharyngitis	4	1	
Dyspnea	3	1	
Skin and Appendages			
Sweating	3	1	
· ·	2	0	
Acne			
Ory skin	2	0	
Special Senses	•	-	
Amblyopia	9	5	
Abnormal vision	2	0	
Jrogenital System			
Dysmenorrhea <sup>2</sup>	2	0	
Vaginitis <sup>2</sup>	2	0	

pain, abnormal dreams, abnormal ejaculation, agitation, akathisia, anorexia, anxiety, arthralgia, cough increased, diarrhea, dyspepsia, emotional lability, fever, latulence, flu syndrome, headache, hostility, insomnia, libido decreased, libido increased, menstrual disorder², myalgia, nausea, nervousness, pain, paranoid reaction, personality disorder, rash, rhinitis, sleep disorder, thinking abnormal, vomiting.

<sup>2</sup> Denominator used was for females only (olanzapine, N=128; placebo, N=51). For specific information about the adverse reactions observed with lithium or valproate, refer to the ADVERSE REACTIONS section of the

package inserts for these other products.

Additional Findings Observed in Clinical Trials
The following findings are based on clinical trials.

Dose Dependency of Adverse Events in Short-Term, Placebo-Controlled Trials

<u>Extraoyramidal Symptoms</u> — The following table enumerates the percentage of patients with treatment-emergent extrapyramidal symptoms as assessed by categorical analyses of formal rating scales during acute therapy in a controlled clinical trial comparing olanzapine at 3 fixed doses with placebo in the treatment of schizophrenia.

### TREATMENT-EMERGENT EXTRAPYRAMIDAL SYMPTOMS ASSESSED BY RATING SCALES INCIDENCE IN A FIXED DOSAGE RANGE, PLACEBO-CONTROLLED

	CLII	NICAL THIAL — ACUTE FHA	3E	
		Percentage of Patie	nts Reporting Event	
	Placebo	Olanzapine 5 ± 2.5 mg/day	Olanzapine 10 ± 2.5 mg/day	Olanzapine 15 ± 2.5 mg/day
Parkinsonism1	15	14	12	14
Akathisia <sup>2</sup>	23	16	19	27
* No statistically significant differ	rences		•	•

Percentage of patients with a Simpson-Angus Scale total score
 Percentage of patients with a Barnes Akathisia Scale global sco

The following table enumerates the percentage of patients with treatment-emergent extrapyramidal symptoms as assessed by spontaneously ported adverse events during acute therapy in the same controlled clinical trial comparing olanzapine at 3 fixed doses with placebo in the treatment of schizophrenia.



TREATMENT-EMERGENT EXTRAPYRAMIDAL SYMPTOMS ASSESSED BY ADVERSE EVENTS INCIDENCE IN A FIXED DOSAGE RANGE, PLACEBO-CONTROLLED

	CLINICA	AL TRIAL — ACUTE PHASE				
		Percentage of Patients Reporting Event				
	Placebo (N=68)	Olanzapine 5 ± 2.5 mg/day (N=65)	Olanzapine 10 ± 2.5 mg/day (N=64)	Olanzapine 15 ± 2.5 mg/day (N=69)		
Dystonic events <sup>1</sup>	1	3	2	3		
Parkinsonism events <sup>2</sup>	10	8	14	20		
Akathisia events <sup>3</sup>	1	5	11*	10*		
Dyskinetic events <sup>4</sup>	4	0	2	1		
Residual events <sup>5</sup>	1	2	5	1		
Any extrapyramidal event	16	15	25	32*		
* Statistically significantly different from	nlaceho		•			

Statistical systimicating unterests into placebox. Patients with the following COSTART terms were counted in this category: dystonia, generalized spasm, neck rigidity, oculogyric crisis, opisthotonos, torticollis Patients with the following COSTART terms were counted in this category: akinesia, cogwheel rigidity, extrapyramidal syndrome, hypertonia, hypokinesia, ma review themselves the results of the control of the review themselves.

facies, tremor.

Patients with the following COSTART terms were counted in this category: akathisia, hyperkinesia.

Patients with the following COSTART terms were counted in this category: buccoglossal syndrome, choreoathetosis, dyskinesia, tardive dyskinesia

Patients with the following COSTART terms were counted in this category: movement disorder, myoclonus, twitching.

Other Adverse Events — The following table addresses dose relatedness for other adverse events using data from a schizophrenia trial involving fixed dosage ranges. It enumerates the percentage of patients with treatment-emergent adverse events for the three fixed-dose range groups and placebo. The data were analyzed using the Cochran-Armitage test, excluding the placebo group, and the table includes only those adverse events for which there was a statistically significant trend.

	Percentage of Patients Reporting Event					
Adverse Event	Placebo (N=68)	Olanzapine 5 ± 2.5 mg/day (N=65)	Olanzapine 10 ± 2.5 mg/day (N=64)	Olanzapine 15 ± 2.5 mg/day (N=69)		
Asthenia	15	8	9	20		
Dry mouth	4	3	5	13		
Nausea	9	0	2	9		
Somnolence	16	20	30	39		
Tremor	3	0	5	7		

Vital Sign Changes — Olanzapine is associated with orthostatic hypotension and tachycardia (see PRECAUTIONS).

Vital Sign Changes — Olanzapine is associated with orthostatic hypotension and tachycardia (see PRECAUTIONS).

Weight Gain — In placebo-controlled, 6-week studies, weight gain was reported in 5.6% of olanzapine patients compared to 0.8% of placebo patients. Olanzapine patients gained an average of 2.8 kg, compared to an average 0.4 kg weight loss in placebo patients; 29% of olanzapine patients ugained an average of 2.8 kg, compared to 3% of placebo patients. A categorization of patients at baseline on the basis of body mass index (BMI) revaeled a significantly greater effect in patients with low BMI compared no normal or overweight patients; nevertheless, weight gain was greater in all 3 olanzapine groups compared to the placebo group. During long-term continuation therapy with olanzapine (238 median days of exposure), 56% of olanzapine patients met the criterion for having gained greater than 7% of their baseline weight. Average weight gain during long-term therapy was 5.4 kg.

Laboratory Changes — An assessment of the premarketing experience for olanzapine revealed an association with asymptomatic increases in SGPT, SGOT, and GGT (see PRECAUTIONS). Olanzapine administration was also associated with increases in serum prolactin (see PRECAUTIONS), with an asymptomatic elevation of the eosinophil count in 0.3% of patients, and with an increase in CPK.

Given the concern about neutropenia associated with other psychotropic compounds and finding of elukopenia associated with the administration of olanzapine in several animal models (see ANIMAL TOXICOLOGY), careful attention was given to examination of hematologic parameters in premarketing studies with olanzapine. There was no indication of a risk of clinically significant neutropenia associated with olanzapine premarketing database for this drug.

ECG Changes — Between-group comparisons for pooled placebo-controlled trials revealed no statistically significant olanzapine/placebo differences in the proportions of patients experiencing potentially important changes

slight tendency to tachycardia may be related to olanzapine's potential for inducing orthostatic changes (see PRECAUTIONS).

Other Adverse Events Observed During the Clinical Trial Evaluation of Olanzapine
Following is a list of terms that reflect treatment-emergent adverse events reported by patients treated with olanzapine (at multiple doses ≥1 mg/day) in clinical trials (8661 patients, 4165 patient-years of exposure). This listing does not include those events already listed in previous tables or elsewhere in labeling, those events for which a drug cause was remote, those event terms which were so general as to be uninformative, and those events reported only once or twice which did not have a substantial probability of being acutely life-threatening.

Events are further categorized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring in at least 1/100 patients (only those not already listed in the tabulated results from placebo-controlled trials appear in this listing); infrequent adverse events are those occurring in 1/100 to 1/1000 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patients; rare events are those occurring in fewer than 1/1/100 patie

Cardiovascular System — Frequent: hypotension; Infrequent: atrial fibrillation, bradycardia, cerebrovascular accident, congestive heart failure, heart arrest, hemorrhage, migraine, pallor, palpitation, vasodilatation, and ventricular extrasystoles; Rare: arteritis, heart failure, and

Digestive System — Frequent: flatulence, increased salivation, and thirst; Infrequent: dysphagia, esophagitis, fecal impaction, fecal incontinence, gastritis, gastroenteritis, gingivitis, hepatitis, melena, mouth ulceration, nausea and vomiting, oral moniliasis, periodontal abscess, rectal hemorrhage, stomatifis, tongue edema, and tooth caries; Rare: aphthous stomatitis, enteritis, eructation, esophageal ulcer, glossitis, ileus, intestinal obstruction, liver fatty deposit, and tongue discoloration.

Endocrine System — Infrequent: diabetes mellitus; Rare: diabetic acidosis and goiter

Hemic and Lymphatic System — Infrequent: anemia, cyanosis, leukocytosis, leukopenia, lymphadenopathy, and thrombocytopenia; emia and thrombocythemia. Metabolic and Nutritional Disorders — Infrequent: acidosis, alkaline phosphatase increased, bilirubinemia, dehydration, hypercholesteremia, hyperalycemia, hyperuricemia, hypocytoemia, hypocytoemia, hyporattemia, lower extremity edema, and upper extremity edema; Rare; gout, hyperattemia, hyporattemia, hyporatt

Musculoskeletal System — Frequent: joint stiffness and twitching; Infrequent: arthritis, arthrosis, leg cramps, and myasthenia; Rare: bone pain, bursitis, myopathy, osteoporosis, and rheumatoid arthritis.

bursitis, myopathy, osteoporosis, and rheumatoid arthritis.

Nervous System — Frequent: abnormal dreams, amnesia, delusions, emotional lability, euphoria, manic reaction, paresthesia, and schizophrenic reaction; Infrequent: akinesia, alcohol misuse, antisocial reaction, ataxia, CNS stimulation, cogwheel rigidity, delirium, dementia, depersonalization, dysarthria, facial paralysis, hypesthesia, hypotonia, incoordination, libido decreased, libido increased, obsessive compulsive symptoms, phobias, somatization, stimulant misuse, stupor, stuttering, tardive dyskinesia, vertigo, and withdrawal syndrome; Rare: circumoral paresthesia, coma, encephalopathy, neuralgia, neuropathy, nystagmus, paralysis, subarachnoid hemorrhage, and tobacco misuse.

Respiratory System — Frequent: Syspena; Infrequent: apnea, asthma, epistaxis, hemoptysis, hypovantilation, hypoxia, laryngitis, and voice alteration; Rare: atelectasis, hiccup, hypoventilation, lung edema, and stridor.

Skin and Appendages — Frequent: sweating; Infrequent: alopecia, contact dermatitis, dry skin, eczema, maculopapular rash, pruritus, seborrhea, skin discoloration, skin ulcer, urticaria, and vesiculobullous rash; Rare: hirsutism and pustular rash.

sebormea, skin discoloration, skin ulcer, urticana, and vesiculobullous rash; Hare: hirsutism and pustular rash.

Special Senses — Frequent: conjunctivitis; Infrequent: abnormality, dascommodation, blephatis, cataract, deafness, diplopia, dry eyes, ear pain, eye hemorrhage, eye inflammation, eye pain, ocular muscle abnormality, taste perversion, and tinnitus; Rare: corneal lesion, glaucoma, keratoconjunctivitis, macular hypopigmentation, miosis, mydriasis, and pigment deposits lens.

Urogenital System — Frequent: vaginitis\*; Infrequent: abnormal ejaculation\*, amenorrhea\*, breast pain, cystitis, decreased menstruation\*, dysuria, female lactation\*, glycosuria, gynecomastia, hematuria, impotence\*, increased menstruation\*, menorrhagia\*, metrorrhagia\*, polyuria, premenstrual syndrome\*, pyuria, urinary referedinor, urinary retetrition, urinary urgency, urination impaired, uterine fibroids enlarged\*, and vaginal hemorrhage\*; Rare: albuminuria, breast enlargement, mastitis, and oliguria.

### Postintroduction Reports

Adverse events reported since market introduction which were temporally (but not necessarily causally) related to ZYPREXA therapy include the llowing: allergic reaction (e.g., anaphylactoid reaction, angioedema, pruritus or urticaria), diabetic coma, pancreatitis, and priapism.

DRUG ABUSE AND DEPENDENCE

### Controlled Substance Class Olanzapine is not a controlled substance.

Physical and Psychological Dependence
In studies prospectively designed to assess abuse and dependence potential, olanzapine was shown to have acute depressive CNS effectitle or no potential of abuse or physical dependence in rats administered oral doses up to 15 times the maximum recommended human daily (20 mg) and rhesus monkeys administered oral doses up to 8 times the maximum recommended human daily dose on a mg/m² basis.

(20 mg) and rhesus monkeys administered oral doses up to 8 times the maximum recommended human daily dose on a mg/m² basis.

Olanzapine has not been systematically studied in humans for its potential for abuse, tolerance, or physical dependence. While the clinical trials did not reveal any tendency for any drug-seeking behavior, these observations were not systematic, and it is not possible to predict on the basis of this limited experience the extent to which a CNS-active drug will be misused, diverted, and/or abused once marketed. Consequently, patients should be evaluated carefully for a history of drug abuse, and such patients should be observed closely for signs of misuse or abuse of olanzapine (e.g., development of tolerance, increases in dose, drug-seeking behavior). OVERDOSAGE

Human Experience
In premarketing trials involving more than 3100 patients and/or normal subjects, accidental or intentional acute overdosage of olanzapine was identified in 67 patients. In the patient taking the largest identified amount, 300 mg, the only symptoms reported were drowsiness and slurred speech. In the limited number of patients who were evaluated in hospitals, including the patient taking 300 mg, there were no observations indicating an adverse change in laboratory analytes or ECG. Vital signs were usually within normal limits following overdoses.

In postmarketing reports of overdose with olanzapine alone, symptoms have been reported in the majority of cases. In symptomatic patients, symptoms with ≥10% incidence included agitation/aggressiveness, dysarthria, tachycardia, various extrapyramidal symptoms, and reduced level of consciousness ranging from sedation to coma. Among less commonly reported symptoms were tollowing potentially medically serious events: aspiration, cardiopulmonary arrest, cardiac arrhythmias (such as supraventricular tachycardia and one patient experiencing sinus pause with spontaneous resumption of normal rhythm), delirium, possible neuroleptic malignant syndrom, respiratory depression/arrest, convulsion, hypertension, and hypotension. Eli Lilly and Company has received reports of fatality in association with overdose of olanzapine alone. In one case of drant, the amount of cardiobul inscription are recorded to he operable, salve as 600 cate however in a control was recorded to he operable. of death, the amount of acutely ingested olanzapine was reported to be possibly as low as 450 mg; however, in another case, a patient was reported to survive an acute olanzapine ingestion of 1500 mg.

to survive an acute olanzapine ingestion or 1500 mg.

Overdosage, Management

The possibility of multiple drug involvement should be considered. In case of acute overdosage, establish and maintain an airway and ensure adequate oxygenation and ventilation, which may include intubation. Gastric lavage (after intubation, if patient is unconscious) and administration of activated charcoal together with a laxative should be considered. The possibility of obtundation, seizures, or dystonic reaction of the head and neck following overdose may create a risk of aspiration with induced emesis. Cardiovascular monitoring should commence immediately and should include continuous electrocardiographic monitoring to detect possible arrhythmias.

There is no specific antidote to clanzapine. Therefore, appropriate supportive measures should be initiated. Hypotension and circulatory collapse should be treated with appropriate measures such as intravenous fluids and/or sympathomimetic agents. (Do not use epinephrine, dopamine, or other sympathomimetics with beta-agonist activity, since beta stimulation may worsen hypotension in the setting of clanzapine-induced alpha blockade.) Close medical supervision and monitoring should continue until the patient recovers.

DOSAGE AND ADMINISTRATION

# Schizophrenia

Schizophrenia 
<u>Usual Dose</u> — Olanzapine should be administered on a once-a-day schedule without regard to meals, generally beginning with 5 to 10 mg 
initially, with a target dose of 10 mg/day within several days. Further dosage adjustments, if indicated, should generally occur at intervals of not less 
than 1 week, since steady state for olanzapine would not be achieved for approximately 1 week in the typical patient. When dosage adjustments are 
necessary, dose increments/decrements of 5 mg QD are recommended.

Efficacy in schizophrenia was demonstrated in a dose range of 10 to 15 mg/day in clinical trials. However, doses above 10 mg/day were no demonstrated to be more efficacious than the 10 mg/day dose. An increase to a dose greater than the target dose of 10 to mg/day (i.e., to a dose o Dosing in Special Populations — The recommended starting dose is 5 mg in patients who are debilitated, who have a predisposition to hypotensive reactions, who otherwise exhibit a combination of factors that may result in slower metabolism of olanzapine (e.g., nonsmoking female patients ≥65 years of age), or who may be more pharmacodynamically sensitive to olanzapine (see CLINICAL PHARMACOLOGY; also see Use in Patients with Concomitant Illness and Drug Interactions under PRECAUTIONS). When indicated, dose escalation should be performed with caution in these patients.

Maintenance Treatment — While there is no body of evidence available to answer the question of how long the patient treated with olanzapine should remain on it, the effectiveness of oral olanzapine, 10 mg/day to 20 mg/day, in maintaining treatment response in schizophrenic patients who had been stable on ZYPREXA for approximately 8 weeks and were then followed for a period of up to 8 months has been demonstrated in a placebo-controlled trial (see CLINICAL PHARMACOLOGY). Patients should be periodically reassessed to determine the need for maintenance treatment with appropriate dose.

<u>Usual Monotherapy Dose</u> — Olanzapine should be administered on a once-a-day schedule without regard to meals, generally beginning with 10 or 15 mg. Dosage adjustments, if indicated, should generally occur at intervals of not less than 24 hours, reflecting the procedures in the placebo-controlled trails. When dosage adjustments are necessary, dose increments/decrements of 5 mg QD are recommended.

Short-term (3-4 weeks) antimanic efficacy was demonstrated in a dose range of 5 mg to 20 mg/day in clinical trials. The safety of doses above 20 mg/day has not been evaluated in clinical trials. Maintenance Monotherapy — The benefit of maintaining bipolar patients on monotherapy with ZYPREXA at a dose of 5 to 20 mg/day, after achieving a responder status for an average duration of two weeks, was demonstrated in a controlled trial (see Clinical Efficacy Data under CLINICAL PHARMACOLOGY). The physician who elects to use ZYPREXA for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient.

Bipolar Mania Usual Dose in Combination with Lithium or Valproate — When administered in combination with lithium or valproate, olanzapine osing should generally begin with 10 mg once-a-day without regard to meals. Short-term (6 weeks) antimanic efficacy was demonstrated in a dose range of 5 mg to 20 mg/day in clinical trials. The safety of doses above 20 mg/day has not been evaluated in clinical trials.

Dosing in Special Populations — See Dosing in Special Populations under DOSAGE AND ADMINISTRATION, Schizophrenia Administration of ZYPREXA ZYDIS (olanzapine orally disintegrating tablets)

After opening sachet, peel back foil on blister. Do not push tablet through foil. Immediately upon opening the blister, using dry hands, remove tablet and place entire ZYPREXA ZYDIS in the mouth. Tablet disintegration occurs rapidly in saliva so it can be easily swallowed with

HOW SUPPLIED

The ZYPREXA 2.5 mg, 5 mg, 7.5 mg, and 10 mg tablets are white, round, and imprinted in blue ink with LILLY and tablet number. The 15 mg tablets are elliptical, blue, and debossed with LILLY and tablet number. The 20 mg tablets are elliptical, pink, and debossed with LILLY and tablet number. The tablets are available as follows:

	TABLET STRENGTH					
	2.5 mg	5 mg	7.5 mg	10 mg	15 mg	20 mg
Tablet No.	4112	4115	4116	4117	4415	4420
Identification	LILLY	LILLY	LILLY	LILLY	LILLY	LILLY
	4112	4115	4116	4117	4415	4420
NDC Codes:						
Bottles 60	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-
	4112-60	4115-60	4116-60	4117-60	4415-60	4420-60
Blisters -	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-
ID* 100	4112-33	4115-33	4116-33	4117-33	4415-33	4420-33
Bottles 1000	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-
	4112-04	4115-04	4116-04	4117-04	4415-04	4420-04

ZYPREXA ZYDIS (clanzapine orally disintegrating tablets) are yellow, round, and debossed with the tablet strength. The tablets are

ZYPREXA ZYDIS	TABLET STRENGTH			
Tablets*	5 mg	10 mg	15 mg	20 mg
Tablet No.	4453	4454	4455	4456
Debossed	5	10	15	20
NDC Codes:				
Dose Pack 30	NDC 0002-	NDC 0002-	NDC 0002-	NDC 0002-
(Child-Resistant)	4453-85	4454-85	4455-85	4456-85

ZYPREXA is a registered trademark of Eli Lilly and Company.

ZYDIS is a registered trademark of R. P. Scherer Corporation.

"ZYPREXA ZYDIS (olanzapine orally disintegrating tablets) is manufactured for Eli Lilly and Company by Scherer DDS Limited, United Kingdom,
SNB SRII.

Store at controlled room temperature, 20° to 25°C (68° to 77°F) [see USP]. The USP defines controlled room temperature as a temperature maintained thermostatically that encompasses the usual and customary working environment of 20° to 25°C (68° to 77°F); that results in a mean kinetic temperature calculated to be not more than 25°C; and that allows for excursions between 15° and 30°C (59° and 86°F) that are experienced in pharmacies, hospitals, and warehouses. Protect from light and moisture

ANIMAL TOXICOLOGY

In animal studies with olanzapine, the principal hematologic findings were reversible peripheral cytopenias in individual dogs dosed at 10 mg/kg (17 times the maximum recommended human daily dose on a mg/m² basis), dose-related decreases in lymphocytes and neutrophils in mice, and lymphopenia in rats. A few dogs treated with 10 mg/kg developed reversible neutropenia and/or reversible hemolytic anemia between 1 and 10 months of treatment. Dose-related decreases in lymphocytes and neutrophils were seen in mice given doses of 10 mg/kg (equal to 2 times the maximum recommended human daily dose on a mg/m² basis in studies of 3 months' duration. Nonseprefic hymphopania consistent with decreased maximum recommended human daily dose on a mg/m² basis) in studies of 3 months' duration. Nonspecific lymphopenia, consistent with decreased body weight gain, occurred in rats receiving 22.5 mg/kg (11 times the maximum recommended human daily dose on a mg/m² basis) for 3 months or 16 mg/kg (8 times the maximum recommended human daily dose on a mg/m² basis) for 6 or 12 months. No evidence of bone marrow cytotoxicity was found in any of the species examined. Bone marrows were normocellular or hypercellular, indicating that the reductions in circulating blood cells were probably due to peripheral (non-marrow) factors.

Literature revised January 14, 2004

Eli Lilly and Company Indianapolis, IN 46285, USA

www.ZYPREXA.com

PRINTED IN USA

