
State Highlights 2002


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men	7,412 4,933	State Lung Cancer All States	90.2/100,000	Overall Projected Death	111,273
Women Death Rate	2,479 326.6/100,000	State CHD* All States	60.6/100,000 59.7/100,000	Rate Projected All	9,905/100,000
All States	295.5/100,000	State COPD† All States	73.6/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6–8			GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AL†	19.1%	26.5%	30.2%	37.6%
Boys†	22.1%	32.0%	32.5%	44.3%
Girls†	15.7%	20.4%	27.8%	30.7%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Alabama Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Alabama is investing 8% of CDC's *Best Practices* lower estimated recommended funding and 3% of the upper estimated recommended funding. Alabama spent \$269 per capita on smoking-attributable direct medical expenditures. In 1998, about 9% (\$186,000,000 or \$352.36 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$433,000,000 Hospital \$300,000,000 Nursing Home \$241,000,000 Prescription Drugs \$110,000,000 Other \$86,000,000 Annual Total \$1,170,000,000 Annual Per Capita \$269

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,777,000,000 \$407

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$96,961,493 Gross cigarette tax revenue collected in 2000 \$70,665,000 Cigarette tax per pack was \$0.165 in 2001

Cigarette sales were 96.2 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$600,000	10/01-9/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$600,000	
Federal—CDC Office on Smoking and Health	\$1,340,047	6/01-5/02
Federal—SAMHSA	\$78,720	10/01-9/02
Non-Government Source—American Legacy Foundation	\$208,156	10/00–9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$1,626,923	
FY02 Total Investment in Tobacco Control		\$2,226,923
CDC Best Practices Recommended Annual Total (Lower F	\$26,740,000	
CDC Best Practices Recommended Annual Total (Upper E	\$71,235,000	
FY02 Per Capita Investment in Tobacco Control	\$0.50	


^{*}An additional \$2,000,000 was appropriated in FY02 to the Department of Education for drug, alcohol, or tobacco education and prevention programs.




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	448 264 184 289.6/100,000	State Lung Cancer All States State CHD* All States	88.5/100,000 90.2/100,000 47.1/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All	19,516 1 10,233/100,000
All States	295.5/100,000	State COPD† All States	86.4/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8			GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AK†	n/a	n/a	36.5%	n/a
Boys†	n/a	n/a	31.1%	n/a
Girls†	n/a	n/a	35.8%	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Alaska Youth Risk Behavior Survey, 1995.

Disparities Among Adult Population Groups, 2000*





Alaska is investing 61% of CDC's Best Practices lower estimated recommended funding and 30% of the upper estimated recommended funding. Alaska spent \$215 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$60,000,000 or \$804.31 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$129,000,000 \$208

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$21,176,139 Gross cigarette tax revenue collected in 2000 \$43,092,000

Cigarette tax per pack was \$1.00 in 2001


Cigarette sales were 66.0 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$3,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$270,000	7/01–6/02
Subtotal: State Appropriation	\$3,270,000	
Federal—CDC Office on Smoking and Health	\$1,145,052	6/01–5/02
Federal—SAMHSA	\$50,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$434,752	3/01–2/02
Subtotal: Federal/National Sources	\$1,629,804	
FY02 Total Investment in Tobacco Control		\$4,899,804
CDC Best Practices Recommended Annual Total (Lower E	\$8,088,000	
CDC Best Practices Recommended Annual Total (Upper E	\$16,512,000	
FY02 Per Capita Investment in Tobacco Control	\$7.72	




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall 6,890 Men 4,382 Women 2,508 Death Rate 283.5/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	81.2/100,000 90.2/100,000 58.5/100,000 59.7/100,000 76.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	77,724 5,686/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8			GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AZ†	11.4%	17.1%	n/a	n/a
Boys†	11.5%	18.0%	n/a	n/a
Girls†	11.3%	16.1%	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Arizona Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*



As of November 30, 2001, the state legislature had not completed its FY2002 budget bill; therefore, investment in tobacco control figures were not available for this report. Arizona spent \$215 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$247,000,000 or \$485.57 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$435,000,000 \$233,000,000 \$175,000,000 \$94,000,000 \$68,000,000 \$1,005,000,000
Annual Per Capita	\$215

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,309,000,000 \$274


State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$87,669,615 Gross cigarette tax revenue collected in 2000 \$162,795,000

Cigarette tax per pack was \$0.58 in 2001


Cigarette sales were 58.5 packs per capita in 2000

Funding Source	FY02 Amount	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	not available	
State Appropriation—Excise Tax Revenue	not available	
State Appropriation—Other	not available	
Subtotal: State Appropriation	not available	
Federal—CDC Office on Smoking and Health	\$256,630	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$91,500	10/00–9/01
Non-Government Source—RWJF/AMA	\$300,000	6/01-5/02
Subtotal: Federal/National Sources	\$648,130	
FY02 Total Investment in Tobacco Control		not available
CDC Best Practices Recommended Annual Total (Lower	\$27,788,000	
CDC Best Practices Recommended Annual Total (Upper	\$71,102,000	
FY02 Per Capita Investment in Tobacco Control	not available	




Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBU 1999 DISEASE-SI)	TO DIE FROM SMOK	
Overall Men	4,901 3,113	State Lung Cancer All States	113.2/100,000 90.2/100.000	Overall Projected Death	66,957
Women	1,788	State CHD*	70.2/100,000	Řate	9,841/100,000
Death Rate	343.3/100,000	All States	59.7/100,000	Projected All	
All States	295.5/100,000	State COPD† All States	71.8/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6-8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AR†	15.8%	22.4%	35.8%	43.8%
Boys†	15.7%	25.4%	37.5%	49.9%
Girls†	15.7%	19.1%	33.7%	37.0%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Arkansas Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Arkansas is investing 79% of CDC's *Best Practices* lower estimated recommended funding and 30% of the upper estimated recommended funding. Arkansas spent \$250 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$189,000,000 or \$445.13 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$208,000,000 \$157,000,000 \$168,000,000 \$59,000,000 \$42,000,000 \$634,000,000
Annual Per Capita	\$250

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,122,000,000 \$440

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.315 in 2001

Cigarette sales were 99.4 packs per capita in 2000


\$51,364,510 \$82,409,000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$11,931,942	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$376,256	7/01–6/02
Subtotal: State Appropriation	\$12,308,198	
Federal—CDC Office on Smoking and Health	\$1,275,017	6/01–5/02
Federal—SAMHSA	\$153,346	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$381,867	6/01–5/02
Subtotal: Federal/National Sources	\$1,810,230	
FY02 Total Investment in Tobacco Control		\$14,118,428
CDC Best Practices Recommended Annual Total (Lower I	\$17,906,000	
CDC Best Practices Recommended Annual Total (Upper I	\$46,445,000	
FY02 Per Capita Investment in Tobacco Control	\$5.30	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUT 1999 DISEASE-SF	9 TO DIE FROM SMO		SMOKING,
Overall	38,233	State Lung Cancer	74.1/100,000	Overall	637,270
Men Women	22,642 15,591	All States State CHD*	90.2/100,000 60.8/100,000	Projected Death Rate	6,890/100,000
Death Rate All States	261.8/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 68.8/100,000 59.7/100,000	Projected All States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
CA [†]	6.7%	10.0%	21.6%	27.8%
Boys [†]	7.0%	10.6%	22.7%	31.0%
Girls [†]	6.4%	9.4%	20.4%	24.3%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † California Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*



California is investing 83% of CDC's Best Practices lower estimated recommended funding and 31% of the upper estimated recommended funding. California spent \$218 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$2,310,000,000 or \$326.14 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$3,703,000,000 Hospital \$1,363,000,000 Nursing Home \$1,140,000,000 Prescription Drugs \$507,000,000 Other \$424,000,000 Annual Total \$7,137,000,000 Annual Per Capita \$218

Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$7,515,000,000 \$227

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$759,225,833 Gross cigarette tax revenue collected in 2000 \$1,176,859,000

Cigarette tax per pack was \$0.87 in 2001

Cigarette sales were 41.6 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$20,000,000	7/01-6/02
State Appropriation—Excise Tax Revenue	\$114,536,000	7/01–6/02
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$134,536,000	
Federal—CDC Office on Smoking and Health	\$335,610	6/01–5/02
Federal—SAMHSA	\$2,000,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$645,679	10/00-9/01
Non-Government Source—RWJF/AMA	\$299,176	6/01-5/02
Subtotal: Federal/National Sources	\$3,280,465	
FY02 Total Investment in Tobacco Control		\$137,816,465
CDC Best Practices Recommended Annual Total (Lower F	\$165,098,000	
CDC Best Practices Recommended Annual Total (Upper B	\$442,403,000	
FY02 Per Capita Investment in Tobacco Control	\$4.02	

^{*}This amount represents one state's portion of the tobacco settlement revenue.


SMOKING-ATTRIBUTABLE DEATHS, 1999	Smoking-Attribut 1999 Disease-Sh	·	Youth Pr to Die from 1999–	A SMOKING,
Overall 4,276 Men 2,537 Women 1,739 Death Rate 248.1/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	62.0/100,000 90.2/100,000 41.5/100,000 59.7/100,000 85.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	92,246 8,380/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
CO [†]	8.8%	13.6%	25.3%	34.4%
Boys [†]	9.3%	15.9%	23.9%	37.1%
Girls [†]	8.4%	11.1%	26.9%	31.6%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Colorado Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Colorado is investing 61% of CDC's *Best Practices* lower estimated recommended funding and 24% of the upper estimated recommended funding. Colorado spent \$259 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$249,000,000 or \$722.84 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$419,000,000 \$239,000,000 \$213,000,000 \$73,000,000 \$82,000,000 \$1,026,000,000
Annual Per Capita	\$259

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$856,000,000 \$211

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$85,026,250 Gross cigarette tax revenue collected in 2000 \$60,818,000

Cigarette tax per pack was \$0.20 in 2001

Cigarette sales were 73.0 packs per capita in 2000

Investment in Tobacco Control

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$12,891,329	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$12,891,329	
Federal—CDC Office on Smoking and Health	\$1,350,347	6/01–5/02
Federal—SAMHSA	\$99,940	10/01-9/02
Non-Government Source—American Legacy Foundation [†]	10/00-9/01	
Non-Government Source—RWJF/AMA	\$383,176	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$14,924,792
CDC Best Practices Recommended Annual Total (Lower E	\$24,546,000	
CDC Best Practices Recommended Annual Total (Upper E	\$63,255,000	
FY02 Per Capita Investment in Tobacco Control	\$3.39	

^{*}An additional \$6,875,375 was appropriated to the Office of the President at the University of Colorado for a tobacco-related research grant program.


†On October 1, 2001, Colorado was awarded an additional grant from the American Legacy Foundation.




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall 4,820 Men 2,639 Women 2,181 Death Rate 255.9/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	78.6/100,000 90.2/100,000 54.7/100,000 59.7/100,000 55.6/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	83,859 9,963/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current Cigarette		Current Any	Current Cigarette	Current Any
Smoking		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
CT [†]	9.8%	13.1%	25.6%	32.4%
Boys [†]	9.7%	14.1%	24.9%	35.3%
Girls [†]	9.8%	11.9%	26.0%	29.2%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Connecticut Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Connecticut is investing 12% of CDC's *Best Practices* lower estimated recommended funding and 5% of the upper estimated recommended funding. Connecticut spent \$389 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$336,000,000 or \$880.85 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$383,000,000 Hospital \$211,000,000 Nursing Home \$496,000,000 Prescription Drugs \$95,000,000 Other \$88,000,000 Annual Total \$1,273,000,000 Annual Per Capita \$389

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$870,000,000 \$265

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$110,430,429 Gross cigarette tax revenue collected in 2000 \$117,215,000

Cigarette tax per pack was \$0.50 in 2001

Cigarette sales were 71.4 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$1,046,550	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$200,000	7/01–6/02
Subtotal: State Appropriation	\$1,246,550	
Federal—CDC Office on Smoking and Health	\$1,010,252	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$316,727	3/01–6/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$2,573,529
CDC Best Practices Recommended Annual Total (Lower I	\$21,240,000	
CDC Best Practices Recommended Annual Total (Upper F	\$53,895,000	
FY02 Per Capita Investment in Tobacco Control	\$0.75	

^{*}An additional \$13,500,000 was appropriated to the Tobacco & Health Trust Fund for tobacco control and other health programs.


Smoking-Attributable Deati 1999	Smoking-Attributable 1999 Disease-Specifi	TO DIE FROM SMOKING,
Overall 1,1 Men 7 Women 4 Death Rate 313.9/100,0 All States 295.5/100,0	All States 90.2 State CHD* 67.5 All States 59.7	/100,000 Overall 19,584 /100,000 Projected Death Rate 10,064/100,000 /100,000 Projected All States 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current Cigarette		Current Any	Current Cigarette	Current Any
Smoking		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
DE†	15.2%	17.8%	27.1%	31.2%
Boys†	14.8%	18.9%	28.0%	34.6%
Girls†	15.6%	16.6%	26.1%	27.8%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Delaware Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Delaware is investing 72% of CDC's *Best Practices* lower estimated recommended funding and 34% of the upper estimated recommended funding. Delaware spent \$298 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$62,000,000 or \$609.71 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$73,000,000 Hospital \$46,000,000 Nursing Home \$62,000,000 Prescription Drugs \$20,000,000 Other \$21,000,000 Annual Total \$222,000,000 Annual Per Capita \$298

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$258,000,000 \$342

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$23,523,288 Gross cigarette tax revenue collected in 2000 \$25,957,000

Cigarette tax per pack was \$0.24 in 2001

Cigarette sales were 140.7 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$5,454,900	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,454,900	
Federal—CDC Office on Smoking and Health	\$782,761	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$6,237,661
CDC Best Practices Recommended Annual Total (Lower E	\$8,631,000	
CDC Best Practices Recommended Annual Total (Upper E	\$18,464,000	
FY02 Per Capita Investment in Tobacco Control	\$7.85	




SMOKING-ATTRIBUTABLE DEATH 1999	199	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall 72 Men 42 Women 20 Death Rate 250.4/100,00 All States 295.5/100,00	6 All States 5 State CHD* 0 All States	80.0/100,000 90.2/100,000 50.4/100,000 59.7/100,000 38.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	7,786 6,771/100,000 8,830/100,000	

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
DC†	9.4%	14.2%	14.7%	21.0%
Boys†	8.7%	15.2%	16.9%	24.9%
Girls†	10.1%	13.4%	12.3%	16.5%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †District of Columbia Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





The District of Columbia is investing 8% of CDC's Best Practices lower estimated recommended funding and 4% of the upper estimated recommended funding. The District of Columbia spent \$364 per capita on smoking-attributable direct medical expenditures. In 1998, about 9% (\$61,000,000 or \$367.58 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$212,000,000 \$408

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

not available Gross cigarette tax revenue collected in 2000 \$17,050,000

Cigarette tax per pack was \$0.65 in 2001


Cigarette sales were 50.2 packs per capita in 2000

FUNDING SOURCE	FUNDING CYCLE	
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$0	
Federal—CDC Office on Smoking and Health	\$448,157	6/01-5/02
Federal—SAMHSA	\$36,292	10/01-9/02
Non-Government Source—American Legacy Foundation	\$100,000	10/00-9/02
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$584,449
CDC Best Practices Recommended Annual Total (Lower E	\$7,479,000	
CDC Best Practices Recommended Annual Total (Upper E	\$14,571,000	
FY02 Per Capita Investment in Tobacco Control	\$1.03	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
	28,713 17,458 11,255 283.1/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD†	91.9/100,000 90.2/100,000 64.2/100,000 59.7/100,000 65.6/100,000	Overall Projected Death Rate Projected All States	306,724 8,412/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6-8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
FL†	9.8%	13.5%	19.0%	25.8%
Boys†	10.0%	15.6%	19.1%	29.6%
Girls†	9.5%	11.2%	19.0%	22.0%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Florida Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Florida is investing 39% of CDC's *Best Practices* lower estimated recommended funding and 14% of the upper estimated recommended funding. Florida spent \$331 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$976,000,000 or \$512.30 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$1,916,000,000 Hospital \$948,000,000 Nursing Home \$1,194,000,000 Prescription Drugs \$478,000,000 Other \$398,000,000 Annual Total \$4,934,000,000 Annual Per Capita \$331

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$5,441,000,000 \$360

State Revenue from Tobacco Sales and Settlement


Tobacco settlement revenue received in 2001 Gross cigarette tax revenue collected in 2000 Cigarette tax per pack was \$0.339 in 2001 Cigarette sales were 82.5 packs per capita in 2000	\$731,300,000 \$426,269,000
--	--------------------------------

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$29,803,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$29,803,000	
Federal—CDC Office on Smoking and Health	\$750,000	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$50,004	10/00-9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$30,603,004
CDC Best Practices Recommended Annual Total (Lower E	\$78,383,000	
CDC Best Practices Recommended Annual Total (Upper E	\$221,260,000	
FY02 Per Capita Investment in Tobacco Control	\$1.88	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	10,728 6,911 3,817 333.9/100,000	State Lung Cancer All States State CHD* All States	99.3/100,000 90.2/100,000 62.5/100,000 59.7/100,000	Overall Projected Death Rate Projected All	187,782 8,657/100,000
All States	295.5/100,000	State COPD† All States	76.2/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
GA	13.8%†	18.8%†	24.3%††	n/a
Boys	13.9%†	20.6%†	24.7%††	n/a
Girls	13.6%†	16.5%†	24.0%††	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Georgia Youth Tobacco Survey, 1999; †Georgia Youth Risk Behavior Survey, 1993.

Disparities Among Adult Population Groups, 2000*





Georgia is investing 56% of CDC's *Best Practices* lower estimated recommended funding and 21% of the upper estimated recommended funding. Georgia spent \$230 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$419,000,000 or \$343.28 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other	\$742,000,000 \$421,000,000 \$311,000,000 \$149,000,000
Prescription Drugs Other Annual Total Annual Per Capita	\$149,000,000 \$135,000,000 \$1,758,000,000 \$230
ī	•

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,734,000,000 \$351

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

\$152,235,170 \$67,064,000

Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.12 in 2001


Cigarette sales were 70.9 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$20,765,890	7/01-6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other \$0		
Subtotal: State Appropriation	\$20,765,890	
Federal—CDC Office on Smoking and Health	\$1,608,025	6/01-5/02
Federal—SAMHSA	\$446,371	10/01-9/02
Non-Government Source—American Legacy Foundation	\$767,286	10/00–9/01
Non-Government Source—RWJF/AMA	\$316,022	6/01-5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$23,903,594
CDC Best Practices Recommended Annual Total (Lower E	\$42,591,000	
CDC Best Practices Recommended Annual Total (Upper E	\$114,314,000	
FY02 Per Capita Investment in Tobacco Control	\$2.86	




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall 1,095 Men 770 Women 325 Death Rate 174.3/100,000 All States 295.5/100,000	All States State CHD* All States	53.7/100,000 90.2/100,000 35.2/100,000 59.7/100,000 30.1/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	23,858 8,067/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
HI [†]	n/a	n/a	24.5%	27.1%
Boys [†]	n/a	n/a	23.3%	26.2%
Girls [†]	n/a	n/a	25.5%	27.7%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; [†]Hawaii Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*



Hawaii is investing 217% of CDC's Best Practices lower estimated recommended funding and 100% of the upper estimated recommended funding. Hawaii spent \$220 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$91,000,000 or \$493.42 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Smoking-Attributable Productivity Costs, 1999

Annual Total	\$263,000,000
Annual Per Capita	\$222

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$35,800,091 Gross cigarette tax revenue collected in 2000 \$40,050,000

Cigarette tax per pack was \$1.00 in 2001

Cigarette sales were 31.9 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$22,171,571	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$22,171,571	
Federal—CDC Office on Smoking and Health	\$874,172	6/01-5/02
Federal—SAMHSA	\$68,989	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$283,057	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$23,397,789	
CDC Best Practices Recommended Annual Total (Lower E	\$10,778,000	
CDC Best Practices Recommended Annual Total (Upper E	\$23,448,000	
FY02 Per Capita Investment in Tobacco Control	\$19.16	


^{*}The amount reflected above is disbursed to a trust fund. Up to an additional \$9,581,399 was appropriated to the Department of Health for promotion and disease prevention programs, including but not limited to maternal and child health and child development programs, promotion of healthy lifestyles and prevention-oriented public health programs.




SMOKING-ATTRIBUTABLE DEATHS, 1999 SMOKING-ATTRIBUTABLE DEAT 1999 DISEASE-SPECIFIC		·	YOUTH PR TO DIE FROM 1999–	SMOKING,	
Overall Men	1,505 940	State Lung Cancer All States	68.1/100,000 90.2/100,000	Overall Projected Death	32,127
Women Death Rate	565 254.6/100,000	State CHD* All States	49.5/100,000 59.7/100,000	Rate Projected All	8,706/100,000
All States	295.5/100,000	State COPD† All States	73.7/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
ID	9.6%†	13.4%†	27.3%††	n/a
Boys	9.7%†	15.4%†	29.3%††	n/a
Girls	9.4%†	11.1%†	25.5%††	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Idaho Youth Tobacco Survey, 2001; ††Idaho Youth Risk Behavior Survey, 1993.

Disparities Among Adult Population Groups, 2000*





Idaho is investing 24% of CDC's Best Practices lower estimated recommended funding and 11% of the upper estimated recommended funding. Idaho spent \$202 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$65,000,000 or \$527.33 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total Annual Per Capita	\$84,000,000 \$57,000,000 \$57,000,000 \$22,000,000 \$29,000,000 \$249,000,000
Annual Per Capita	\$202

Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$271,000,000 \$217

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$22,531,022 Gross cigarette tax revenue collected in 2000 \$25,250,000

Cigarette tax per pack was \$0.28 in 2001

Cigarette sales were 66.9 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$1,006,500	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$300,000	7/01–6/02
Subtotal: State Appropriation	\$1,306,500	
Federal—CDC Office on Smoking and Health	\$1,113,388	6/01–5/02
Federal—SAMHSA	\$50,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$2,669,888
CDC Best Practices Recommended Annual Total (Lower I	\$11,044,000	
CDC Best Practices Recommended Annual Total (Upper I	\$24,087,000	
FY02 Per Capita Investment in Tobacco Control	\$2.02	

^{*}An additional \$50,000 was appropriated to the Buhl School District Substance Abuse Pilot Program to address tobacco and/or substance abuse. An additional \$120,000 was appropriated to the Idaho Supreme Court to address tobacco and/or substance abuse.




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	18,476 10,947 7,529 303.1/100,000	State Lung Cancer All States State CHD* All States	91.0/100,000 90.2/100,000 69.6/100,000 59.7/100,000	Overall Projected Death Rate Projected All	297,384 9,163/100,000
All States	295.5/100,000	State COPD† All States	63.7/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
IL†	n/a	n/a	35.7%	n/a
Boys†	n/a	n/a	35.8%	n/a
Girls†	n/a	n/a	35.6%	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Illinois Youth Risk Behavior Survey, 1995.

Disparities Among Adult Population Groups, 2000*





Illinois is investing 78% of CDC's Best Practices lower estimated recommended funding and 28% of the upper estimated recommended funding. Illinois spent \$266 per capita on smoking-attributable direct medical expenditures. In 1998, about 18% (\$1,226,000,000 or \$899.06 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$1,097,000,000 Hospital \$801,000,000 Nursing Home \$846,000,000 Prescription Drugs \$268,000,000 Other \$194,000,000 Annual Total \$3,206,000,000 Annual Per Capita \$266

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$3,909,000,000 \$322

State Revenue from Tobacco Sales and Settlement


Tobacco settlement revenue received in 2001 \$288,674,835 Gross cigarette tax revenue collected in 2000 \$485,041,000

Cigarette tax per pack was \$0.58 in 2001

Cigarette sales were 70.0 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$48,900,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$48,900,000	
Federal—CDC Office on Smoking and Health	\$1,655,389	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$265,960	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$50,821,349
CDC Best Practices Recommended Annual Total (Lower I	\$64,909,000	
CDC Best Practices Recommended Annual Total (Upper F	\$179,048,000	
FY02 Per Capita Investment in Tobacco Control	\$4.06	

^{*}An additional \$2,250,000 was appropriated for school-based clinics.




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	10,271 6,146 4,125 341.4/100,000	State Lung Cancer All States State CHD* All States	107.1/100,000 90.2/100,000 71.0/100,000 59.7/100,000	Overall Projected Death Rate Projected All	167,604 1 10,646/100,000
All States	295.5/100,000	State COPD† All States	78.1/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
IN [†]	9.8%	15.3%	31.6%	36.9%
Boys [†]	9.3%	16.3%	32.8%	41.1%
Girls [†]	10.4%	14.2%	30.1%	32.3%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Indiana Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Indiana is investing 19% of CDC's *Best Practices* lower estimated recommended funding and 7% of the upper estimated recommended funding. Indiana spent \$275 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$380,000,000 or \$625.31 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$501,000,000 Hospital \$419,000,000 Nursing Home \$500,000,000 Prescription Drugs \$134,000,000 Other \$73,000,000 Annual Total \$1,627,000,000 Annual Per Capita \$275

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,164,000,000 \$364

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$106,515,675 Gross cigarette tax revenue collected in 2000 \$117,604,000

Cigarette tax per pack was \$0.155 in 2001

Cigarette sales were 125.5 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$5,000,000	7/01-6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,000,000	
Federal—CDC Office on Smoking and Health	\$1,492,125	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01-5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$6,692,125
CDC Best Practices Recommended Annual Total (Lower E	\$34,784,000	
CDC Best Practices Recommended Annual Total (Upper E	\$95,804,000	
FY02 Per Capita Investment in Tobacco Control	\$1.09	

^{*}Funding dedicated to tobacco control from settlement revenue is deposited into a trust fund. In FY01, \$35,000,000 was appropriated to tobacco control. The FY01 appropriation will be rolled over into the FY02 budget.




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall 4,615 Men 2,936 Women 1,679 Death Rate 265.6/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	79.7/100,000 90.2/100,000 57.7/100,000 59.7/100,000 68.6/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All States	80,319 h 10,948/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
IA [†]	11.8%	16.4%	32.7%	39.0%
Boys [†]	11.3%	17.5%	34.2%	45.6%
Girls [†]	12.3%	15.1%	31.1%	31.9%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Iowa Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Iowa is investing 58% of CDC's Best Practices lower estimated recommended funding and 23% of the upper estimated recommended funding. Iowa spent \$278 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$235,000,000 or \$745.39 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total Annual Per Capita	\$220,000,000 \$190,000,000 \$264,000,000 \$68,000,000 \$52,000,000 \$794,000,000 \$278
Annual Per Capita	\$278

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$824,000,000 \$287

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$53,940,181 Gross cigarette tax revenue collected in 2000 \$92,817,000

Cigarette tax per pack was \$0.36 in 2001


Cigarette sales were 88.9 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$9,420,394	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other		
Subtotal: State Appropriation	\$9,420,394	
Federal—CDC Office on Smoking and Health	\$1,043,669	6/01–5/02
Federal—SAMHSA	\$663,250	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$11,127,313	
CDC Best Practices Recommended Annual Total (Lower E	\$19,347,000	
CDC Best Practices Recommended Annual Total (Upper E	\$48,713,000	
FY02 Per Capita Investment in Tobacco Control	\$3.78	




	ING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000		
Overall Men Women Death Rate	3,864 2,510 1,354 269.8/100,000	State Lung Cancer All States State CHD* All States	83.6/100,000 90.2/100,000 48.1/100,000 59.7/100,000	Overall Projected Death Rate Projected All	57,358 8,045/100,000
All States	295.5/100,000	State COPD† All States	70.1/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
KS†	8.1%	12.0%	26.1%	33.6%
Boys†	8.0%	13.2%	24.7%	37.2%
Girls†	7.9%	10.3%	27.5%	29.7%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Kansas Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Kansas is investing 15% of CDC's Best Practices lower estimated recommended funding and 6% of the upper estimated recommended funding. Kansas spent \$274 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$153,000,000 or \$632.56 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$741,000,000 \$279

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$51,707,586 Gross cigarette tax revenue collected in 2000 \$50,887,000

Cigarette tax per pack was \$0.24 in 2001


Cigarette sales were 79.8 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE	
State Appropriation—Settlement (Tobacco Only)	\$500,000	7/01–6/02	
State Appropriation—Excise Tax Revenue	State Appropriation—Excise Tax Revenue \$0		
State Appropriation—Other			
Subtotal: State Appropriation	\$500,000		
Federal—CDC Office on Smoking and Health	\$1,229,700	6/01-5/02	
Federal—SAMHSA	10/01-9/02		
Non-Government Source—American Legacy Foundation	10/00-9/01		
Non-Government Source—RWJF/AMA	6/01-5/02		
Subtotal: Federal/National Sources	\$2,271,608		
FY02 Total Investment in Tobacco Control		\$2,771,608	
CDC Best Practices Recommended Annual Total (Lower E	\$18,052,000		
CDC Best Practices Recommended Annual Total (Upper E	\$44,689,000		
FY02 Per Capita Investment in Tobacco Control	\$1.02		


Smoking-Attributable Deaths, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall 7,791 Men 4,836 Women 2,955 Death Rate 387.1/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	121.4/100,000 90.2/100,000 78.6/100,000 59.7/100,000 91.6/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	114,995 11,559/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
KY†	21.5%	28.3%	37.4%	46.2%
Boys†	21.9%	33.0%	35.6%	50.2%
Girls†	21.1%	23.2%	39.0%	41.8%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; \dagger Kentucky Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Kentucky is investing 20% of CDC's *Best Practices* lower estimated recommended funding and 7% of the upper estimated recommended funding. Kentucky spent \$298 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$380,000,000 or \$589.46 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$371,000,000 Hospital \$313,000,000 Nursing Home \$299,000,000 Prescription Drugs \$108,000,000 Other \$80,000,000 Annual Total \$1,171,000,000 Annual Per Capita \$298

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,841,000,000 \$465

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$104,957,133 Gross cigarette tax revenue collected in 2000 \$18,724,000

Cigarette tax per pack was \$0.03 in 2001

Cigarette sales were 156.2 packs per capita in 2000

FUNDING SOURCE	FUNDING CYCLE	
State Appropriation—Settlement (Tobacco Only)*	\$3,040,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other		
Subtotal: State Appropriation	\$3,040,000	
Federal—CDC Office on Smoking and Health	\$1,128,413	6/01–5/02
Federal—SAMHSA	\$182,208	10/01-9/02
Non-Government Source—American Legacy Foundation	\$99,261	10/00-9/01
Non-Government Source—RWJF/AMA	3/01–2/02	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$4,912,696
CDC Best Practices Recommended Annual Total (Lower E	\$25,090,000	
CDC Best Practices Recommended Annual Total (Upper E	\$69,895,000	
FY02 Per Capita Investment in Tobacco Control	\$1.21	


^{*}An additional \$2,500,000 was appropriated to the Kentucky Agency for Substance Abuse for smoking cessation and other substance abuse programs.




Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBUTE 1999 DISEASE-ST)	Youth Pr to Die from 1999–	SMOKING,
Overall	6,421	State Lung Cancer		Overall	107,968
Men	3,896	All States	90.2/100,000	Projected Death	
Women	2,525	State CHD*	65.8/100,000	Rate	8,851/100,000
Death Rate	314.1/100,000	All States	59.7/100,000	Projected All	
All States	295.5/100,000	State COPD† All States	60.1/100,000 59.7/100,000	Štates	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
LA	17.1%†	26.3% [†]	36.4%††	n/a
Boys	18.1%†	30.6% [†]	38.2%††	n/a
Girls	15.4%†	20.8% [†]	34.6%††	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Louisiana Youth Tobacco Survey, 2001; ††Louisiana Youth Risk Behavior Survey, 1997.

Disparities Among Adult Population Groups, 2000*





Louisiana is investing 9% of CDC's *Best Practices* lower estimated recommended funding and 3% of the upper estimated recommended funding. Louisiana spent \$264 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$518,000,000 or \$719.08 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$392,000,000 Hospital \$308,000,000 Nursing Home \$268,000,000 Prescription Drugs \$101,000,000 Other \$82,000,000 Annual Total \$1,151,000,000 Annual Per Capita \$264

Smoking-Attributable Productivity Costs, 1999

Annual Total	\$1,662,000,000
Annual Per Capita	\$380

State Fiscal Resources Available for Tobacco Control

Tobacco settlement revenue received in 2001 \$139,997,812 Gross cigarette tax revenue collected in 2000 \$92,299,000

Cigarette tax per pack was \$0.24 in 2001

Cigarette sales were 104.3 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$500,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other		
Subtotal: State Appropriation	\$500,000	
Federal—CDC Office on Smoking and Health	\$1,140,677	6/01-5/02
Federal—SAMHSA	\$400,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$299,174	6/01-5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$2,339,851	
CDC Best Practices Recommended Annual Total (Lower E	\$27,132,000	
CDC Best Practices Recommended Annual Total (Upper E	\$71,431,000	
FY02 Per Capita Investment in Tobacco Control	\$0.52	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men	2,132 1,216	State Lung Cancer All States	95.7/100,000 90.2/100,000	Overall Projected Deatl	33,292
Women Death Rate	916 305.2/100,000	State CHD* All States	56.7/100,000 59.7/100,000	Rate Projected All	11,052/100,000
All States	295.5/100,000	State COPD† All States	82.5/100,000 59.7/100,000	Štates	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8			GRADES	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
ME†	11.7%	13.7%	28.6%	32.6%
Boys†	11.3%	14.4%	28.6%	35.2%
Girls†	12.3%	12.9%	28.5%	29.9%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Maine Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Maine is investing 125% of CDC's Best Practices lower estimated recommended funding and 55% of the upper estimated recommended funding. Maine spent \$377 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$169,000,000 or \$996.71 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$134,000,000 \$110,000,000 Hospital Nursing Home \$134,000,000 Prescription Drugs \$43,000,000 Other \$49,000,000 Annual Total \$470,000,000 Annual Per Capita \$377

Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$406,000,000 \$324

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$47,268,164 \$77,235,000 Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.74 in 2001

Cigarette sales were 82.9 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE			
State Appropriation—Settlement (Tobacco Only)*	\$12,655,488	7/01–6/02			
State Appropriation—Excise Tax Revenue	\$0				
State Appropriation—Other	\$0				
Subtotal: State Appropriation	\$12,655,488				
Federal—CDC Office on Smoking and Health	\$1,001,691	6/01–5/02			
Federal—SAMHSA	\$30,000	10/01-9/02			
Non-Government Source—American Legacy Foundation	\$0				
Non-Government Source—RWJF/AMA	\$331,493	6/01–5/02			
Subtotal: Federal/National Sources					
FY02 Total Investment in Tobacco Control	FY02 Total Investment in Tobacco Control				
CDC Best Practices Recommended Annual Total (Lower I	\$11,189,000				
CDC Best Practices Recommended Annual Total (Upper F	\$25,353,000				
FY02 Per Capita Investment in Tobacco Control	\$10.96				

^{*}An additional \$1,100,000 was appropriated to the Department of Health Services, Medical Care for payments to Medicaid health care providers for services that include smoking cessation and other health services.


SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall	6,831	State Lung Cancer	93.2/100,000	Overall	104,889
Men	3,987	All States	90.2/100,000	Projected Death	
Women	2,844	State CHD*	59.8/100,000	Řate	7,734/100,000
Death Rate	283.3/100,000	All States	59.7/100,000	Projected All	
All States	295.5/100,000	State COPD†	61.2/100,000	Štates	8,830/100,000
		All States	59.7/100,000		

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MD†	7.3%	11.8%	23.7%	29.9%
Boys†	7.3%	12.5%	23.3%	32.1%
Girls†	7.3%	10.8%	23.8%	27.4%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Maryland Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Maryland is investing 107% of CDC's Best Practices lower estimated recommended funding and 41% of the upper estimated recommended funding. Maryland spent \$299 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$372,000,000 or \$663.31 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$588,000,000 Hospital \$309,000,000 Nursing Home \$400,000,000 Prescription Drugs \$129,000,000 Other \$107,000,000 Annual Total \$1,533,000,000 Annual Per Capita \$299

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,555,000,000 \$301

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$140,202,664 Gross cigarette tax revenue collected in 2000 \$200,760,000

Cigarette tax per pack was \$0.66 in 2001

Cigarette sales were 57.7 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE		
State Appropriation—Settlement (Tobacco Only)*	\$30,671,765	7/01–6/02		
State Appropriation—Excise Tax Revenue	\$0			
State Appropriation—Other	\$0			
Subtotal: State Appropriation	\$ 30,671,765			
Federal—CDC Office on Smoking and Health	\$1,370,605	6/01-5/02		
Federal—SAMHSA	\$0			
Non-Government Source—American Legacy Foundation	\$0			
Non-Government Source—RWJF/AMA	\$257,036	6/01-5/02		
Subtotal: Federal/National Sources				
FY02 Total Investment in Tobacco Control		\$32,299,406		
CDC Best Practices Recommended Annual Total (Lower	\$30,301,000			
CDC Best Practices Recommended Annual Total (Upper	\$78,601,000			
FY02 Per Capita Investment in Tobacco Control	FY02 Per Capita Investment in Tobacco Control			

^{*}A total of \$10,674,000 of the \$30,671,756 may not be expended until the Secretary of Budget and Management determines that sufficient tobacco settlement funds are available. An additional \$1,177,408 is budgeted by the Department of Health from general fund revenue for tobacco control.




Smoking-Attributable Deaths, 1999		1999	Smoking-Attributable Deaths, 1999 Disease-Specific		cojected A Smoking, 2000
	9,008 5,064 3,944 2/100,000 5/100,000	State Lung Cancer All States State CHD* All States State COPD†	85.1/100,000 90.2/100,000 47.2/100,000 59.7/100,000 60.9/100,000	Overall Projected Death Rate Projected All States	117,753 7,850/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MA†	n/a	n/a	30.3%	35.1%
Boys†	n/a	n/a	29.9%	37.8%
Girls†	n/a	n/a	30.7%	32.3%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Massachusetts Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





As of November 30, 2001, the state legislature had not completed its FY2002 budget bill; therefore, investment in tobacco control figures were not available for this report. Massachusetts spent \$450 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$817,000,000 or \$899.73 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$861,000,000 Hospital \$570,000,000 Nursing Home \$934,000,000 Prescription Drugs \$194,000,000 Other \$207,000,000 Annual Total \$2,766,000,000 Annual Per Capita \$450

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,588,000,000 \$257

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$240,246,288 Gross cigarette tax revenue collected in 2000 \$274,467,000

Cigarette tax per pack was \$0.76 in 2001

Cigarette sales were 58.3 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	not available	
State Appropriation—Excise Tax Revenue	not available	
State Appropriation—Other	not available	
Subtotal: State Appropriation	not available	
Federal—CDC Office on Smoking and Health	\$1,571,990	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$449,370	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		not available
CDC Best Practices Recommended Annual Total (Lower I	\$35,244,000	
CDC Best Practices Recommended Annual Total (Upper F	\$92,758,000	
FY02 Per Capita Investment in Tobacco Control	not available	




	BUTABLE DEATHS, 999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Pr to Die from 1999–	SMOKING,
Overall Men Women Death Rate All States	14,741 8,905 5,836 299.0/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	88.6/100,000 90.2/100,000 71.3/100,000 59.7/100,000 66.6/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	255,356 9,837/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MI [†]	9.3%	14.2%	27.6%	34.1%
Boys [†]	9.8%	15.8%	25.8%	36.1%
Girls [†]	8.7%	12.4%	27.5%	32.0%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Michigan Youth Tobacco Survey, 2001.

Disparities Among Adult Population Groups, 2000*





Michigan is investing 12% of CDC's Best Practices lower estimated recommended funding and 4% of the upper estimated recommended funding. Michigan spent \$270 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$881,000,000 or \$646.11 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$897,000,000 \$734,000,000 \$575,000,000 \$286,000,000 \$163,000,000
Annual Total Annual Per Capita	\$2,655,000,000

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$3,405,000,000 \$345

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.75 in 2001

Cigarette sales were 83.7 packs per capita in 2000

\$258,862,265 \$601,780,000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE		
State Appropriation—Settlement (Tobacco Only)	\$0			
State Appropriation—Excise Tax Revenue	\$0			
State Appropriation—Other*	\$3,900,000	10/01–9/02		
Subtotal: State Appropriation	\$3,900,000			
Federal—CDC Office on Smoking and Health	\$1,775,695	6/01-5/02		
Federal—SAMHSA	\$0			
Non-Government Source—American Legacy Foundation	\$318,530	10/00-9/01		
Non-Government Source—RWJF/AMA	\$340,380	6/01-5/02		
Subtotal: Federal/National Sources				
FY02 Total Investment in Tobacco Control	FY02 Total Investment in Tobacco Control			
CDC Best Practices Recommended Annual Total (Lower F	\$54,804,000			
CDC Best Practices Recommended Annual Total (Upper F	\$154,558,000			
FY02 Per Capita Investment in Tobacco Control	FY02 Per Capita Investment in Tobacco Control			


^{*}Funding amounts not verified by state budget office.




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attribut 1999 Disease-Sp	Ź	YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall Men Women Death Rate All States	5,618 3,410 2,208 231.1/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	73.8/100,000 90.2/100,000 37.9/100,000 59.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	112,085 8,710/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MN†	9.1%	12.6%	32.4%	38.7%
Boys†	8.7%	12.9%	32.0%	42.7%
Girls†	9.5%	12.3%	32.6%	34.1%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Minnesota Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Minnesota is investing 105% of CDC's Best Practices lower estimated recommended funding and 41% of the upper estimated recommended funding. Minnesota spent \$341 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$363,000,000 or \$675.57 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$679,000,000 \$296,000,000 \$434,000,000 \$105,000,000 \$97,000,000 \$1,611,000,000
Annual Total	\$1,611,000,000
Annual Per Capita	\$341

Smoking-Attributable Productivity Costs, 1999

\$336,997,000

\$177,262,000

Annual Total Annual Per Capita	\$1,027,000,000 \$215

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.48 in 2001


Cigarette sales were 76.0 packs per capita in 2000

FUNDING SOURCE	FY02 Amount	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$17,610,935	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Non-Governmental Funding—MN Partnership for Action Against Tobacco	\$10,900,000	7/01–6/02
Subtotal: State Sources	\$28,510,935	
Federal—CDC Office on Smoking and Health	\$1,376,165	6/01–5/02
Federal—SAMHSA	\$71,710	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$232,748	03/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$30,191,558	
CDC Best Practices Recommended Annual Total (Lower I	\$28,624,000	
CDC Best Practices Recommended Annual Total (Upper F	\$74,013,000	
FY02 Per Capita Investment in Tobacco Control	\$6.07	


Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall Men	4,961 3,306	State Lung Cancer All States	115.0/100,000 90.2/100.000	Overall Projected Death	58,515
Women	1,655	State CHD*	80.2/100,000	Řate	7,549/100,000
Death Rate All States	367.8/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 71.5/100,000 59.7/100,000	Projected All States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MS [†]	17.8%	25.3%	30.5%	40.6%
Boys [†]	19.2%	29.4%	33.3%	47.8%
Girls [†]	16.3%	20.8%	27.6%	33.3%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Mississippi Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Mississippi is investing 120% of CDC's Best Practices lower estimated recommended funding and 48% of the upper estimated recommended funding. Mississippi spent \$204 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$206,000,000 or \$424.29 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$184,000,000 \$154,000,000 \$130,000,000 \$55,000,000 \$38,000,000 \$561,000,000
	. , ,
Annual Per Capita	\$204

Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$1,298,000,000 \$469

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

\$211,149,286

Gross cigarette tax revenue collected in 2000 Cigarette tax per pack was \$0.18 in 2001

\$49,247,000


Cigarette sales were 97.2 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Non-Governmental Source—Partnership for a Healthier	\$21,782,613	6/01-5/02
Mississippi		
Subtotal: State Sources	\$21,782,613	
Federal—CDC Office on Smoking and Health	\$380,796	6/01–5/02
Federal—SAMHSA	\$131,834	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$299,196	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$22,594,439	
CDC Best Practices Recommended Annual Total (Lower F	\$18,788,000	
CDC Best Practices Recommended Annual Total (Upper F	\$46,804,000	
FY02 Per Capita Investment in Tobacco Control	\$7.87	


Overall 10,271 State Lung Cancer 101.6/100,000 Overall	
Men 6,238 Women All States 90.2/100,000 Projected Death Women 4,033 State CHD* 79.7/100,000 Rate 9,770/Projected All Death Rate 345.3/100,000 All States 59.7/100,000 Projected All Projected Death	139,484 100,000 100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MO	14.9%†	19.5%†	32.8% ††	39.0%††
Boys	13.7%†	n/a	35.6% ††	44.0%††
Girls	16.3%†	n/a	30.1% ††	33.9%††

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Missouri Youth Tobacco Survey, 1999; ††Missouri Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





Missouri is investing 72% of CDC's *Best Practices* lower estimated recommended funding and 26% of the upper estimated recommended funding. Missouri spent \$307 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$415,000,000 or \$565.59 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$507,000,000 Hospital \$452,000,000 Nursing Home \$466,000,000 Prescription Drugs \$133,000,000 Other \$110,000,000 Annual Total \$1,668,000,000 Annual Per Capita \$307

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,173,000,000 \$397

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$142,110,326 Gross cigarette tax revenue collected in 2000 \$107,169,000

Cigarette tax per pack was \$0.17 in 2001

Cigarette sales were 113.8 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$22,176,700	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$22,176,700	
Federal—CDC Office on Smoking and Health	\$1,166,052	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources	\$1,366,052	
FY02 Total Investment in Tobacco Control		\$23,542,752
CDC Best Practices Recommended Annual Total (Lower I	\$32,767,000	
CDC Best Practices Recommended Annual Total (Upper F	\$91,359,000	
FY02 Per Capita Investment in Tobacco Control	\$4.17	


^{*}Due to a budget shortfall, \$3,500,000 of this money is withheld.




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUT 1999 DISEASE-SF		YOUTH PROJECTED TO DIE FROM SMOKING 1999–2000	
Overall Men Women	1,434 858 576	State Lung Cancer All States State CHD*	86.3/100,000 90.2/100,000 44.5/100,000	Overall Projected Death Rate	17,143 7,451/100,000
Death Rate All States	294.7/100,000 295.5/100,000	All States State COPD [†] All States	59.7/100,000 90.3/100,000 59.7/100,000	Projected All States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADES	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MT [†]	n/a	n/a	35.0%	45.0%
Boys [†]	n/a	n/a	35.4%	52.3%
Girls [†]	n/a	n/a	34.6%	37.6%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Montana Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





Montana is investing 20% of CDC's *Best Practices* lower estimated recommended funding and 10% of the upper estimated recommended funding. Montana spent \$246 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$52,000,000 or \$514.32 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$67,000,000 \$62,000,000 \$52,000,000 \$18,000,000 \$17,000,000 \$216,000,000
Annual Per Capita	\$246

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$247,000,000 \$280

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$26,248,117 Gross cigarette tax revenue collected in 2000 \$12,442,000

Cigarette tax per pack was \$0.18 in 2001


Cigarette sales were 75.5 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$502,165	7/01–6/02
Subtotal: State Appropriation	\$502,165	
Federal—CDC Office on Smoking and Health	\$875,000	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$178,282	10/00-9/01
Non-Government Source—RWJF/AMA	\$349,174	6/01-5/02
Subtotal: Federal/National Sources	\$1,402,456	
FY02 Total Investment in Tobacco Control		\$1,904,621
CDC Best Practices Recommended Annual Total (Lower E	\$9,355,000	
CDC Best Practices Recommended Annual Total (Upper E	\$19,679,000	
FY02 Per Capita Investment in Tobacco Control	\$2.09	


	FRIBUTABLE DEATHS, 1999 SMOKING-ATTRIBUTABLE DEA 1999 DISEASE-SPECIFIC		·	TO DIE FRO	Projected om Smoking, 9–2000
Overall Men Women Death Rate	2,449 1,552 897 264.2/100,000	State Lung Cancer All States State CHD* All States	81.2/100,000 90.2/100,000 43.5/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All	10,153/100,000
All States	295.5/100,000	State COPD† All States	72.7/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8			GRADES 9–12		
Current C	_	Current Any	Current Cigarette	Current Any	
Smok		Tobacco Use	Smoking	Tobacco Use	
National*	11.0%	15.1%	28.0%	34.5%	
NE	10.0%†	13.8% [†]	29.0%††	35.8%††	
Boys	10.2%†	15.6% [†]	27.1%††	38.9%††	
Girls	9.8%†	11.8% [†]	30.8%††	32.8%††	

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Nebraska Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Nebraska is investing 65% of CDC's *Best Practices* lower estimated recommended funding and 28% of the upper estimated recommended funding. Nebraska spent \$252 per capita on smoking-attributable direct medical expenditures. In 1998, about 12% (\$105,000,000 or \$499.41 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$116,000,000
Hospital	\$104,000,000
Nursing Home	\$139,000,000
Prescription Drugs	\$39,000,000
Other	\$21,000,000
Annual Total	\$419,000,000
Annual Per Capita	\$252

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$439,000,000 \$264

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$36,903,219 Gross cigarette tax revenue collected in 2000 \$44,994,000

Cigarette tax per pack was \$0.34 in 2001

Cigarette sales were 77.6 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$7,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$7,000,000	
Federal—CDC Office on Smoking and Health	\$1,347,138	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$296,098	6/01–5/02
Subtotal: Federal/National Sources	\$1,643,236	
FY02 Total Investment in Tobacco Control		\$8,643,236
CDC Best Practices Recommended Annual Total (Lower E	\$13,308,000	
CDC Best Practices Recommended Annual Total (Upper E	\$31,041,000	
FY02 Per Capita Investment in Tobacco Control	\$5.01	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBU 1999 DISEASE-S	1999 TO DIE FROM SMOI		OM SMOKING,
Overall Men Women Death Rate	3,359 1,910 1,449 414.3/100,000	State Lung Cancer All States State CHD* All States	90.2/100,000 82.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All	11,084/100,000
All States	295.5/100,000	State COPD [†] All States	107.4/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

	GRADE	ES 6-8	GRADES	9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NV†	n/a	n/a	32.6%	39.9%
Boys†	n/a	n/a	32.4%	44.6%
Girls†	n/a	n/a	32.5%	34.5%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; [†]Nevada Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





Nevada is investing 38% of CDC's *Best Practices* lower estimated recommended funding and 16% of the upper estimated recommended funding. Nevada spent \$253 per capita on smoking-attributable direct medical expenditures. In 1998, about 19% (\$96,000,000 or \$748.56 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$199,000,000
Hospital	\$113,000,000
Nursing Home	\$43,000,000
Prescription Drugs	\$41,000,000
Other	\$45,000,000
Annual Total	\$441,000,000
Annual Per Capita	\$253
_	

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$762,000,000 \$421

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$37,830,589 Gross cigarette tax revenue collected in 2000 \$61,017,000

Cigarette tax per pack was \$0.35 in 2001


Cigarette sales were 93.2 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$4,093,565	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$4,093,565	
Federal—CDC Office on Smoking and Health	\$748,437	6/01–5/02
Federal—SAMHSA	\$70,500	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA \$204,062		6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$5,116,564	
CDC Best Practices Recommended Annual Total (Lower E	\$13,477,000	
CDC Best Practices Recommended Annual Total (Upper E	\$32,993,000	
FY02 Per Capita Investment in Tobacco Control	\$2.46	


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	1,688 989 699 291.9/100,000	State Lung Cancer All States State CHD* All States	92.5/100,000 90.2/100,000 58.4/100,000 59.7/100,000	Overall Projected Death Rate Projected All	11,209/100,000
All States	295.5/100,000	State COPD† All States	78.7/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smol		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NH	12.0%†	15.2%†	36.0%††	n/a
Boys	11.1%†	16.0%†	32.0%††	n/a
Girls	12.8%†	14.2%†	39.9%††	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †New Hampshire Youth Tobacco Survey, 2000; *New Hampshire Youth Tobacco Survey, 2000; *New Hampshire Youth Tobacco Survey, 2000; *New Hampshire Youth Tobacco Survey,

Disparities Among Adult Population Groups, 2000*





New Hampshire is investing 40% of CDC's Best Practices lower estimated recommended funding and 18% of the upper estimated recommended funding. New Hampshire spent \$371 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$90,000,000 or \$958.66 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$153,000,000 \$90,000,000 \$119,000,000 \$37,000,000 \$41,000,000 \$440,000,000
Annual Per Capita	\$371

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$339,000,000 \$282

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$43,303,862 Gross cigarette tax revenue collected in 2000 \$93,588,000

Cigarette tax per pack was \$0.52 in 2001


Cigarette sales were 147.3 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$3,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$3,000,000	
Federal—CDC Office on Smoking and Health	\$991,588	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	6/01–5/02	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$4,406,545	
CDC Best Practices Recommended Annual Total (Lower E	\$10,888,000	
CDC Best Practices Recommended Annual Total (Upper E	\$24,766,000	
FY02 Per Capita Investment in Tobacco Control	\$3.53	


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women	10,756 6,332 4,424	State Lung Cancer All States State CHD*	80.7/100,000 90.2/100,000 53.5/100,000	Overall Projected Death Rate	168,828 8,087/100,000
Death Rate All States	243.7/100,000 295.5/100,000	All States State COPD [†] All States	59.7/100,000 51.1/100,000 59.7/100,000	Projected All States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NJ [†]	10.5%	18.9%	27.6%	38.9%
Boys [†]	9.9%	20.4%	26.4%	42.3%
Girls [†]	11.0%	17.3%	28.4%	35.0%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; [†]New Jersey Youth Tobacco Survey, 1999.

Disparities Among Adult Population Groups, 2000*





New Jersey is investing 72% of CDC's Best Practices lower estimated recommended funding and 27% of the upper estimated recommended funding. New Jersey spent \$306 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$755,000,000 or \$928.76 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$854,000,000
Hospital	\$497,000,000
Nursing Home	\$712,000,000
Prescription Drugs	\$255,000,000
Other	\$163,000,000
Annual Total	\$2,481,000,000
Annual Per Capita	\$306

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,226,000,000 \$273

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$239,846,695 Gross cigarette tax revenue collected in 2000 \$393,179,000

Cigarette tax per pack was \$0.80 in 2001


Cigarette sales were 60.1 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$30,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$30,000,000	
Federal—CDC Office on Smoking and Health	\$1,257,351	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$757,118	10/00-9/01
Non-Government Source—RWJF/AMA	\$447,197	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$32,461,666
CDC Best Practices Recommended Annual Total (Lower E	\$45,073,000	
CDC Best Practices Recommended Annual Total (Upper E	\$121,328,000	
FY02 Per Capita Investment in Tobacco Control	\$3.83	


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
	2,109 1,302 807 7/100,000 5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	62.8/100,000 90.2/100,000 55.2/100,000 59.7/100,000 82.3/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	44,564 8,763/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NM [†]	n/a	n/a	30.1%	n/a
Boys [†]	n/a	n/a	n/a	n/a
Girls [†]	n/a	n/a	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; [†]New Mexico Youth Risk Behavior Survey, 1991.

Disparities Among Adult Population Groups, 2000*





New Mexico is investing 51% of CDC's *Best Practices* lower estimated recommended funding and 22% of the upper estimated recommended funding. New Mexico spent \$208 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$144,000,000 or \$436.51 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$135,000,000 \$103,000,000 \$58,000,000 \$29,000,000 \$35,000,000 \$360,000,000
Annual Per Capita	\$208

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$397,000,000 \$228

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$36,989,850 Gross cigarette tax revenue collected in 2000 \$21,007,000

Cigarette tax per pack was \$0.21 in 2001


Cigarette sales were 53.8 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$5,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,000,000	
Federal—CDC Office on Smoking and Health	\$1,181,300	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$460,229	10/00-9/01
Non-Government Source—RWJF/AMA	\$293,480	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$6,935,009
CDC Best Practices Recommended Annual Total (Lower E	\$13,711,000	
CDC Best Practices Recommended Annual Total (Upper E	\$31,947,000	
FY02 Per Capita Investment in Tobacco Control	\$3.75	


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men	24,572 13,439	State Lung Cancer All States	76.7/100,000 90.2/100,000	Overall Projected Death	438,769
Women Death Rate All States	11,133 252.1/100,000 295.5/100,000	State CHD* All States State COPD† All States	66.7/100,000 59.7/100,000 51.5/100,000 59.7/100,000	Rate Projected All States	9,355/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6-8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NY†	9.3%	11.8%	26.8%	32.8%
Boys†	9.8%	13.5%	24.8%	34.2%
Girls†	8.8%	9.9%	29.2%	30.9%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; \dagger New York Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





New York is investing 49% of CDC's Best Practices lower estimated recommended funding and 17% of the upper estimated recommended funding. New York spent \$351 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$4,271,000,000 or \$1,389.77 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$1,767,000,000 \$1,296,000,000 \$2,230,000,000 \$494,000,000 \$592,000,000 \$6,379,000,000
Annual Per Capita	\$351

Smoking-Attributable Productivity Costs, 1999

Annual Total	\$5,303,000,000
Annual Per Capita	\$291

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$754,339,700 Gross cigarette tax revenue collected in 2000 \$738,115,000

Cigarette tax per pack was \$1.11 in 2001 and increased to \$1.50 effective April 2002

Cigarette sales were 57.8 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$42,500,000	4/01–3/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$42,500,000	
Federal—CDC Office on Smoking and Health	\$1,999,998	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$1,529,984	10/00–9/01
Non-Government Source—RWJF/AMA	\$450,016	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$46,479,998
CDC Best Practices Recommended Annual Total (Lower F	\$95,830,000	
CDC Best Practices Recommended Annual Total (Upper F	\$269,296,000	
FY02 Per Capita Investment in Tobacco Control	\$2.44	


^{*}An additional \$2,000,000 is appropriated to the Department of Health for administrative costs for cancer and tobacco control programs. Also, an additional \$2,000,000 was appropriated for services and expenses related to tobacco control and cancer programs. In New York State, approximately one-half of all Tobacco Settlement funds are shared with counties. These figures represent the New York State share of Tobacco Settlement appropriations and do not include settlement funds appropriated by counties for tobacco prevention and control.




SMOKING-ATTRIBUTABLE DEAT 1999	1999	Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall 11,5 Men 7,3 Women 4,1 Death Rate 301.1/100,0 All States 295.5/100,0	All States State CHD* All States	90.2/100,00 0* 62.4/100,00 59.7/100,00 PD† 69.8/100,00	Projected Dear Rate Projected All States	210,718 th 10,718/100,000 8,830/100,000	

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NC†	15.0%	18.4%	31.6%	38.3%
Boys†	16.0%	21.0%	33.4%	44.0%
Girls†	14.0%	15.7%	29.7%	32.4%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †North Carolina Youth Tobacco Survey, 1999.

Disparities Among Adult Population Groups, 2000*



North Carolina is investing 9% of CDC's Best Practices lower estimated recommended funding and 3% of the upper estimated recommended funding. North Carolina spent \$255 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$600,000,000 or \$513.30 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$646,000,000 \$478,000,000 \$491,000,000 \$161,000,000 \$147,000,000 \$1,923,000,000
Annual Total	\$1,923,000,000
Annual Per Capita	\$255

Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$2,829,000,000 \$370

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$142,728,818 Gross cigarette tax revenue collected in 2000

\$42,378,000

Cigarette tax per pack was \$0.05 in 2001


Cigarette sales were 109.0 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$650,000	7/01–6/02
Subtotal: State Appropriation	\$650,000	
Federal—CDC Office on Smoking and Health	\$1,848,922	6/01–5/02
Federal—SAMHSA	\$208,961	10/01-9/02
Non-Government Source—American Legacy Foundation	\$1,141,057	10/00-9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$3,198,940	
FY02 Total Investment in Tobacco Control		\$3,848,940
CDC Best Practices Recommended Annual Total (Lower E	\$42,591,000	
CDC Best Practices Recommended Annual Total (Upper E	\$118,626,000	
FY02 Per Capita Investment in Tobacco Control		\$0.47


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUT 1999 DISEASE-SF	99 TO DIE FROM SMOKI		SMOKING,
Overall Men	855 593	State Lung Cancer All States State CHD*	64.1/100,000 90.2/100,000	Overall Projected Death	15,638
Women Death Rate All States	262 227.7/100,000 295.5/100,000	All States State COPD† All States	53.1/100,000 59.7/100,000 50.9/100,000 59.7/100,000	Rate Projected All States	9,722/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6-8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
ND†	n/a	n/a	40.6%	n/a
Boys†	n/a	n/a	40.2%	n/a
Girls†	n/a	n/a	41.0%	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †North Dakota Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





North Dakota is investing 41% of CDC's Best Practices lower estimated recommended funding and 20% of the upper estimated recommended funding. North Dakota spent \$303 per capita on smoking-attributable direct medical expenditures. In 1998, about 11% (\$37,000,000 or \$604.68 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$54,000,000 \$56,000,000 \$61,000,000 \$13,000,000 \$9,000,000 \$193,000,000
Annual Per Capita	\$303

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$158,000,000 \$249

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$22,701,724 Gross cigarette tax revenue collected in 2000 \$21,132,000

Cigarette tax per pack was \$0.44 in 2001

Cigarette sales were 72.5 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$2,055,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$2,055,000	
Federal—CDC Office on Smoking and Health	\$1,099,997	6/01–5/02
Federal—SAMHSA	\$30,537	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources	\$1,330,534	
FY02 Total Investment in Tobacco Control		\$3,385,534
CDC Best Practices Recommended Annual Total (Lower E	\$8,161,000	
CDC Best Practices Recommended Annual Total (Upper E	\$16,547,000	
FY02 Per Capita Investment in Tobacco Control		\$5.27




SMOKING-ATTRIBUTABLE DEATH	1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall 18,9 Men 11,0 Women 7,8 Death Rate 317.6/100,0 All States 295.5/100,0	All States State CHD* All States	r 97.6/100,000 90.2/100,000 70.4/100,000 59.7/100,000 74.2/100,000 59.7/100,000	Overall Projected Deat Rate Projected All States	314,717 h 10,896/100,000 8,830/100,000	

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6-8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
OH†	13.7%	18.7%	33.4%	41.1%
Boys†	14.7%	22.0%	33.0%	45.6%
Girls†	12.7%	15.2%	33.7%	35.9%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Ohio Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Ohio is investing 197% of CDC's Best Practices lower estimated recommended funding and 70% the upper estimated recommended funding. Ohio spent \$304 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$1,113,000,000 or \$862.47 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$1,030,000,000 Hospital \$808,000,000 Nursing Home \$1,113,000,000 Prescription Drugs \$274,000,000 Other \$191,000,000 Annual Total \$3,416,000,000 Annual Per Capita \$304

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$4,146,000,000 \$368

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$312,446,299 Gross cigarette tax revenue collected in 2000 \$271,260,000

Cigarette tax per pack was \$0.24 in 2001

Cigarette sales were 99.9 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE		
State Appropriation—Settlement (Tobacco Only)*	\$119,578,148	7/01–6/02		
State Appropriation—Excise Tax Revenue	\$0			
State Appropriation—Other	\$0			
Subtotal: State Appropriation	\$119,578,148			
Federal—CDC Office on Smoking and Health	\$1,525,232	6/01–5/02		
Federal—SAMHSA	\$82,950	10/01-9/02		
Non-Government Source—American Legacy Foundation	\$0			
Non-Government Source—RWJF/AMA	\$444,658	3/01–2/02		
Subtotal: Federal/National Sources				
FY02 Total Investment in Tobacco Control	FY02 Total Investment in Tobacco Control			
CDC Best Practices Recommended Annual Total (Lower)	\$61,735,000			
CDC Best Practices Recommended Annual Total (Upper l	\$173,676,000			
FY02 Per Capita Investment in Tobacco Control	\$10.67			

^{*}The amount reflected above is disbursed to an endowment fund where the funds will be used by the Foundation for tobacco use reduction program. The Foundation may, but is not required to, treat these funds as an endowment. An additional \$636,000 will be expended by the Department of Public Safety to enforce underage-use laws in establishments with liquor stores. Also, an additional \$579,323 will go to the Department of Health to provide emergency medications and oxygen to low-income seniors adversely affected by tobacco.


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		HS, YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall	5,781	State Lung Cancer	98.4/100,000	Overall	77,943
Men	3,589	All States	90.2/100,000	Projected Death	
Women	2,192	State CHD*	75.3/100,000	Rate	8,734/100,000
Death Rate	320.1/100,000	All States	59.7/100,000	Projected All	
All States	295.5/100,000	State COPD† All States	71.3/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6-8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
OK†	16.9%	21.0%	33.0%	42.0%
Boys†	17.9%	23.8%	31.8%	47.3%
Girls†	15.8%	17.8%	34.0%	36.4%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Oklahoma Youth Tobacco Survey, 1999.

Disparities Among Adult Population Groups, 2000*



Oklahoma is investing 17% of CDC's *Best Practices* lower estimated recommended funding and 7% of the upper estimated recommended funding. Oklahoma spent \$272 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$170,000,000 or \$498.01 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$290,000,000
Hospital	\$230,000,000
Nursing Home	\$232,000,000
Prescription Drugs	\$80,000,000
Other	\$75,000,000
Annual Total	\$907,000,000
Annual Per Capita	\$272

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,334,000,000 \$397

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.23 in 2001

Cigarette sales were 108.9 packs per capita in 2000

Investment in Tobacco Control

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE			
State Appropriation—Settlement (Tobacco Only)	\$1,750,000	7/01–6/02			
State Appropriation—Excise Tax Revenue	\$0				
State Appropriation—Other	\$0				
Subtotal: State Appropriation	\$1,750,000				
Federal—CDC Office on Smoking and Health	\$1,299,907	6/01–5/02			
Federal—SAMHSA	\$0				
Non-Government Source—American Legacy Foundation	\$750,000	10/00-9/01			
Non-Government Source—RWJF/AMA	\$0				
Subtotal: Federal/National Sources	\$2,049,907				
FY02 Total Investment in Tobacco Control	FY02 Total Investment in Tobacco Control				
CDC Best Practices Recommended Annual Total (Lower I	\$21,825,000				
CDC Best Practices Recommended Annual Total (Upper I	\$56,310,000				
FY02 Per Capita Investment in Tobacco Control	\$1.09				

\$64,265,818


\$63,926,000




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUT 1999 DISEASE-SF	·	TO DIE FROM SM	
Overall Men Women	4,986 2,991 1,995	State Lung Cancer All States State CHD*	85.7/100,000 90.2/100,000 47.2/100,000	Overall Projected Death Rate	73,132 8,639/100,000
Death Rate All States	278.6/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 74.3/100,000 59.7/100,000	Projected All States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
OR	n/a	n/a	n/a	n/a
Boys	n/a	n/a	n/a	n/a
Girls	n/a	n/a	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: *National Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Oregon is investing 62% of CDC's *Best Practices* lower estimated recommended funding and 25% of the upper estimated recommended funding. Oregon spent \$265 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$224,000,000 or \$438.97 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$328,000,000 Hospital \$182,000,000 Nursing Home \$208,000,000 Prescription Drugs \$74,000,000 Other \$79,000,000 Annual Total \$871,000,000 Annual Per Capita \$265

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$908,000,000 \$274

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$68,264,932 Gross cigarette tax revenue collected in 2000 \$166,359,000

Cigarette tax per pack was \$0.68 in 2001

Cigarette sales were 72.0 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$2,500,000	7/01-6/02
State Appropriation—Excise Tax Revenue	\$9,086,500	7/01–6/02
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$11,586,500	
Federal—CDC Office on Smoking and Health	\$1,112,463	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$353,453	6/01–5/02
Subtotal: Federal/National Sources	\$1,465,916	
FY02 Total Investment in Tobacco Control		\$13,052,416
CDC Best Practices Recommended Annual Total (Lower	\$21,131,000	
CDC Best Practices Recommended Annual Total (Upper	\$52,840,000	
FY02 Per Capita Investment in Tobacco Control	\$3.75	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		S, YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall Men Women Death Rate All States	19,798 12,045 7,753 270.2/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	85.3/100,000 90.2/100,000 59.1/100,000 59.7/100,000 59.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	311,074 1 10,645/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
PA†	13.1%	17.0%	27.6%	34.0%
Boys†	13.4%	19.3%	26.9%	36.8%
Girls†	12.7%	14.4%	28.3%	31.0%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Pennsylvania Youth Tobacco Survey, 2001.

Disparities Among Adult Population Groups, 2000*





Pennsylvania is investing 66% of CDC's Best Practices lower estimated recommended funding and 23% of the upper estimated recommended funding. Pennsylvania spent \$338 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$1,335,000,000 or \$876.33 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$1,231,000,000
Hospital	\$918,000,000
Nursing Home	\$1,311,000,000
Prescription Drugs	\$363,000,000
Other	\$231,000,000
Annual Total	\$4,054,000,000
Annual Per Capita	\$338

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$3,941,000,000 \$329

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$341,834,697 Gross cigarette tax revenue collected in 2000 \$333,665,000

Cigarette tax per pack was \$0.31 in 2001

Cigarette sales were 87.9 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$41,402,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$41,402,000	
Federal—CDC Office on Smoking and Health	\$1,260,000	6/01–5/02
Federal—SAMHSA	\$300,000	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$43,162,000
CDC Best Practices Recommended Annual Total (Lower E	\$65,568,000	
CDC Best Practices Recommended Annual Total (Upper E	\$184,759,000	
FY02 Per Capita Investment in Tobacco Control	\$3.50	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men	1,704 942	State Lung Cancer All States	97.2/100,000 90.2/100,000	Overall Projected Death	23,875
Women Death Rate	763 283.2/100,000	State CHD* All States	69.2/100,000 59.7/100,000	Rate Projected All	9,634/100,000
All States	295.5/100,000	State COPD [†] All States	60.6/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

GRADES 6–8			Grades 9–12		
Current Cigarette		Current Any	Current Cigarette Current A		
Smoking		Tobacco Use	Smoking Tobacco		
National*	11.0%	15.1%	28.0%	34.5%	
RI [†]	9.1%	13.2%	26.0%	32.1%	
Boys [†]	7.3%	12.4%	26.9%	36.0%	
Girls [†]	10.8%	13.6%	25.2%	27.8%	

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Rhode Island Youth Tobacco Survey, 2001.

Disparities Among Adult Population Groups, 2000*





Rhode Island is investing 46% of CDC's Best Practices lower estimated recommended funding and 21% of the upper estimated recommended funding. Rhode Island spent \$400 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$140,000,000 or \$912.01 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$111,000,000 \$90,000,000 \$119,000,000 \$33,000,000 \$42,000,000 \$395,000,000
Annual Per Capita	\$400

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$283,000,000 \$286

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$44,589,359 Gross cigarette tax revenue collected in 2000 \$58,788,000

Cigarette tax per pack was \$1.00 in 2001

Cigarette sales were 83.1 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$3,000,000	7/01–6/02
Subtotal: State Appropriation	\$3,000,000	
Federal—CDC Office on Smoking and Health	\$1,100,000	6/01-5/02
Federal—SAMHSA	\$200,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$199,819	6/01-5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$4,499,819
CDC Best Practices Recommended Annual Total (Lower E	\$9,888,000	
CDC Best Practices Recommended Annual Total (Upper E	\$21,908,000	
FY02 Per Capita Investment in Tobacco Control	\$4.27	




SMOKING-ATTRIBUTABLE DEATH	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall 5,9 Men 3,9 Women 2,0 Death Rate 315.6/100,0 All States 295.5/100,0	All States State CHD* All States	96.5/100,000 90.2/100,000 64.1/100,000 59.7/100,000 69.5/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	90,715 8,985/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8			GRADES 9–12	
Current Cigarette Smoking				
National* SC [†] Boys [†] Girls [†]	11.0% n/a n/a n/a	15.1% n/a n/a n/a	28.0% 36.0% 37.7% 34.2%	34.5% 41.5% 46.4% 36.8%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †South Carolina Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





South Carolina is investing 14% of CDC's *Best Practices* lower estimated recommended funding and 5% of the upper estimated recommended funding. South Carolina spent \$223 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$307,000,000 or \$516.11 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,672,000,000 \$430

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$72,961,122 Gross cigarette tax revenue collected in 2000 \$28,077,000

Cigarette tax per pack was \$0.07 in 2001


Cigarette sales were 103.9 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$1,620,470	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$62,809	7/01–6/02
Subtotal: State Appropriation	\$1,683,279	
Federal—CDC Office on Smoking and Health	\$1,200,000	6/01-5/02
Federal—SAMHSA	10/01-9/02	
Non-Government Source—American Legacy Foundation	10/00-9/01	
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$3,248,862
CDC Best Practices Recommended Annual Total (Lower E	\$23,905,000	
CDC Best Practices Recommended Annual Total (Upper E	\$62,013,000	
FY02 Per Capita Investment in Tobacco Control	\$0.80	


SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUT 1999 DISEASE-SH)	Youth Pr to Die from 1999–	A SMOKING,
Overall 1,066 Men 728 Women 338 Death Rate 230.2/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	70.9/100,000 90.2/100,000 49.1/100,000 59.7/100,000 54.3/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	20,033 9,886/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99



Youth Tobacco Use


GRADES 6-8		GRADES 9–12		
Current C	_	Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
SD	12.4%†	16.3%†	32.6%††	39.2%††
Boys	14.5%†	20.4%†	30.4%††	41.5%††
Girls	10.0%†	11.7%†	34.9%††	36.9%††

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: *National Youth Tobacco Survey, 2000; †South Dakota Youth Tobacco Survey, 1999; ††South Dakota Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*

*Calculations for racial/ethnic groups are based on combined 1999 and 2000 data.



Percent of Population Groups Who Smoke



South Dakota is investing 52% of CDC's Best Practices lower estimated recommended funding and 25% of the upper estimated recommended funding. South Dakota spent \$293 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$45,000,000 or \$504.20 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$68,000,000 \$58,000,000 \$63,000,000 \$14,000,000 \$11,000,000 \$214,000,000
Annual Per Capita	\$293

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$189,000,000 \$258

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$21,642,999 Gross cigarette tax revenue collected in 2000 \$19,251,000

Cigarette tax per pack was \$0.33 in 2001


Cigarette sales were 75.1 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$1,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$2,269,532	7/01–6/02
Subtotal: State Appropriation	\$3,269,532	
Federal—CDC Office on Smoking and Health	\$886,491	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$331,533	6/01-5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$4,487,556
CDC Best Practices Recommended Annual Total (Lower I	\$8,688,000	
CDC Best Practices Recommended Annual Total (Upper I	\$18,214,000	
FY02 Per Capita Investment in Tobacco Control	\$5.90	




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DE 1999 DISEASE-SPECIFIC	THS, YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000
Overall 9,624 Men 6,099 Women 3,525 Death Rate 346.0/100,000 All States 295.5/100,000	State Lung Cancer 110.3/10 All States 90.2/10 State CHD* 77.3/10 All States 59.7/10 State COPD† 76.9/10 All States 59.7/10	,000 Projected Death ,000 Rate 8,890/100,000 ,000 Projected All ,000 States 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
TN†	16.6%	23.2%	32.4%	41.3%
Boys†	17.5%	27.0%	33.4%	47.1%
Girls†	15.6%	19.1%	31.3%	35.1%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; \dagger Tennessee Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Tennessee is investing 6% of CDC's Best Practices lower estimated recommended funding and 2% of the upper estimated recommended funding. Tennessee spent \$311 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$531,000,000 or \$288.17 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$615,000,000 Hospital \$406,000,000 Nursing Home \$435,000,000 Prescription Drugs \$139,000,000 Other \$96,000,000 Annual Total \$1,691,000,000 Annual Per Capita \$311

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,440,000,000 \$445

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 $\,$

\$151,394,114 \$79,981,000

Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.13 in 2001


Cigarette sales were 108.7 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$0	
Federal—CDC Office on Smoking and Health	\$1,439,207	6/01-5/02
Federal—SAMHSA	\$125,469	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$312,060	6/01-5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$1,876,736	
CDC Best Practices Recommended Annual Total (Lower E	\$32,233,000	
CDC Best Practices Recommended Annual Total (Upper E	\$89,079,000	
FY02 Per Capita Investment in Tobacco Control	\$0.33	




Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
	24,158 15,105 9,071 288.8/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	87.1/100,000 90.2/100,000 64.1/100,000 59.7/100,000 69.9/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	486,631 8,267/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
TX†	13.9%	17.6%	28.1%	34.6%
Boys†	14.6%	19.8%	32.0%	41.8%
Girls†	13.2%	15.6%	23.6%	26.3%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; [†]Texas Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Texas is investing 16% of CDC's *Best Practices* lower estimated recommended funding and 6% of the upper estimated recommended funding. Texas spent \$231 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$1,265,000,000 or \$543.87 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$1,778,000,000 Hospital \$1,046,000,000 Nursing Home \$910,000,000 Prescription Drugs \$397,000,000 Other \$421,000,000 Annual Total \$4,552,000,000 Annual Per Capita \$231

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$5,540,000,000 \$276

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$974,220,834 Gross cigarette tax revenue collected in 2000 \$536,877,000

Cigarette tax per pack was \$0.41 in 2001


Cigarette sales were 69.3 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$12,500,000	9/01-8/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$1,207,016	9/01-8/02
Subtotal: State Appropriation	\$13,707,016	
Federal—CDC Office on Smoking and Health	\$969,828	6/01-5/02
Federal—SAMHSA	\$1,560,117	10/01-9/02
Non-Government Source—American Legacy Foundation	\$172,500	10/00-9/01
Non-Government Source—RWJF/AMA	\$337,060	6/01-5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$16,746,521	
CDC Best Practices Recommended Annual Total (Lower I	\$103,288,000	
CDC Best Practices Recommended Annual Total (Upper I	\$284,735,000	
FY02 Per Capita Investment in Tobacco Control	\$0.79	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall Men Women Death Rate	1,218 828 390 159.8/100,000	State Lung Cancer All States State CHD* All States	38.7/100,000 90.2/100,000 27.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All	38,325 5,333/100,000
All States	295.5/100,000	State COPD [†] All States	50.4/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
UT†	n/a	n/a	11.9%	14.5%
Boys†	n/a	n/a	11.7%	16.2%
Girls†	n/a	n/a	11.8%	12.2%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; \dagger Utah Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





Utah is investing 11% of CDC's Best Practices lower estimated recommended funding and 5% of the upper estimated recommended funding. Utah spent \$130 per capita on smoking-attributable direct medical expenditures. In 1998, about 12% (\$81,000,000 or \$373.05 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$244,000,000 \$115

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$27,593,646 \$46,235,000 Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.515 in 2001

Cigarette sales were 40.7 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$0	
State Appropriation—Excise Tax Revenue	\$250,000	7/01–6/02
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$250,000	
Federal—CDC Office on Smoking and Health	\$1,268,944	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation†	\$0	
Non-Government Source—RWJF/AMA	\$199,885	4/01–3/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$1,718,829	
CDC Best Practices Recommended Annual Total (Lower I	\$15,230,000	
CDC Best Practices Recommended Annual Total (Upper I	\$33,383,000	
FY02 Per Capita Investment in Tobacco Control	\$0.75	

^{*}A total of \$4,039,800 was appropriated to the Department of Health Community and Family Services for tobacco, alcohol, and drug prevention programs. An additional \$2,000,000 may be appropriated as a supplemental for FY02 per Senate Bill 15 in the upcoming 2002 general session. †On October 1, 2001, Utah received \$467,635 from the American Legacy Foundation.


SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall 824 Men 517 Women 309 Death Rate 274.3/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	90.2/100,000 90.2/100,000 49.4/100,000 59.7/100,000 75.6/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	15,801 10,711/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
VT	11.9%†	15.3% [†]	33.4%††	38.6%††
Boys	9.6%†	14.8% [†]	33.3%††	42.1%††
Girls	14.2%†	15.7% [†]	33.6%††	34.8%††

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Vermont Youth Tobacco Survey, 2000; †Vermont Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





Vermont is investing 98% of CDC's Best Practices lower estimated recommended funding and 49% of the upper estimated recommended funding. Vermont spent \$308 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$56,000,000 or \$451.52 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$60,000,000 \$39,000,000 \$47,000,000 \$16,000,000 \$20,000,000 \$182,000,000
Annual Per Capita	\$308

Smoking-Attributable Productivity Costs, 1999

Annual Total	\$172,000,000
Annual Per Capita	\$290

State Revenue from Tobacco Sales and Settlement


Tobacco settlement revenue received in 2001 \$24,458,081 Gross cigarette tax revenue collected in 2000 \$24,144,000

Cigarette tax per pack was \$0.44 in 2001

Cigarette sales were 88.9 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$5,824,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,824,000	
Federal—CDC Office on Smoking and Health	\$1,146,500	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$500,000	10/00-9/01
Non-Government Source—RWJF/AMA	\$301,584	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$7,772,084
CDC Best Practices Recommended Annual Total (Lower E	\$7,905,000	
CDC Best Practices Recommended Annual Total (Upper E	\$15,938,000	
FY02 Per Capita Investment in Tobacco Control	\$12.67	

^{*}An additional \$1,028,500 was appropriated for to bacco and other health programs.




	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Pr to Die from 1999–	A SMOKING,	
Overall Men Women Death Rate	9,157 5,604 3,553 290.8/100,000	State Lung Cancer All States State CHD* All States	94.9/100,000 90.2/100,000 54.0/100,000 59.7/100,000	Overall Projected Death Rate Projected All	134,529 7,739/100,000
All States	295.5/100,000	State COPD† All States	65.6/100,000 59.7/100,000	States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

Grades 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
VA	n/a	n/a	n/a	n/a
Boys	n/a	n/a	n/a	n/a
Girls	n/a	n/a	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: *National Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Virginia is investing 44% of CDC's Best Practices lower estimated recommended funding and 16% of the upper estimated recommended funding. Virginia spent \$240 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$313,000,000 or \$479.69 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory \$597,000,000 Hospital \$406,000,000 Nursing Home \$353,000,000 Prescription Drugs \$150,000,000 Other \$123,000,000 Annual Total \$1,629,000,000 Annual Per Capita \$240

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,080,000,000 \$303

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$126,823,184 Gross cigarette tax revenue collected in 2000 \$16,918,000

Cigarette tax per pack was \$0.025 in 2001

Cigarette sales were 96.7 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$15,807,984	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$15,807,984	
Federal—CDC Office on Smoking and Health	\$1,131,145	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$16,939,129
CDC Best Practices Recommended Annual Total (Lower E	\$38,866,000	
CDC Best Practices Recommended Annual Total (Upper E	\$106,854,000	
FY02 Per Capita Investment in Tobacco Control	\$2.36	




SMOKING-ATTRIBUTABLE DEATHS, 1999	THS, SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		, YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall 7,761 Men 4,689 Women 3,072 Death Rate 284.9/100,000 All States 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	90.1/100,000 90.2/100,000 52.3/100,000 59.7/100,000 76.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	133,952 8,848/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.

Adult Cigarette Use, 2000



Environmental Tobacco Smoke, 1998–99




Youth Tobacco Use

Grades 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
WA	n/a	n/a	n/a	n/a
Boys	n/a	n/a	n/a	n/a
Girls	n/a	n/a	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: *National Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Washington is investing 62% of CDC's Best Practices lower estimated recommended funding and 23% of the upper estimated recommended funding. Washington spent \$269 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$508,000,000 or \$359.58 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,503,000,000 \$261

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$127,351,202 Gross cigarette tax revenue collected in 2000 \$254,770,000

Cigarette tax per pack was \$0.825 in 2001 and will increase to \$1.425 effective July 2002


Cigarette sales were 52.8 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$17,496,000	7/01-6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$898,500	7/01-6/02
Subtotal: State Appropriation	\$18,394,500	
Federal—CDC Office on Smoking and Health	\$1,524,995	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$722,795	10/00-9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$2,247,790	
FY02 Total Investment in Tobacco Control		\$20,642,290
CDC Best Practices Recommended Annual Total (Lower	\$33,341,000	
CDC Best Practices Recommended Annual Total (Upper	\$89,381,000	
FY02 Per Capita Investment in Tobacco Control	\$3.44	




SMOKING-ATTRIBUTABLE DEATHS, 1999		HS, SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		S, YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
	3,838 2,348 1,492 354.1/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	115.2/100,000 90.2/100,000 77.4/100,000 59.7/100,000 86.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	42,537 10,571/100,000 8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

GRADES 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
WV [†]	18.1%	25.3%	38.5%	47.9%
Boys [†]	17.5%	28.8%	36.0%	52.2%
Girls [†]	18.8%	21.4%	40.6%	42.5%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †West Virginia Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





West Virginia is investing 56% of CDC's *Best Practices* lower estimated recommended funding and 23% of the upper estimated recommended funding. West Virginia spent \$298 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$179,000,000 or \$520.55 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$175,000,000 \$152,000,000 \$118,000,000 \$52,000,000 \$42,000,000 \$539,000,000
Annual Per Capita	\$298

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$850,000,000 \$470

State Revenue from Tobacco Sales and Settlement


Tobacco settlement revenue received in 2001 \$54,981,783 Gross cigarette tax revenue collected in 2000 \$33,766,000

Cigarette tax per pack was \$0.17 in 2001

Cigarette sales were 107.9 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$5,850,592	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,850,592	
Federal—CDC Office on Smoking and Health	\$1,145,612	6/01-5/02
Federal—SAMHSA	\$35,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$955,689	10/00-9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$7,986,893
CDC Best Practices Recommended Annual Total (Lower E	\$14,160,000	
CDC Best Practices Recommended Annual Total (Upper E	\$35,365,000	
FY02 Per Capita Investment in Tobacco Control	\$4.41	

^{*}An additional \$500,000 was appropriated for the Tobacco/Alcohol Education Program. The appropriation language also states that to the extent permitted by law, eight classified exempt positions shall be provided for the educator-inspector positions to be used in the education and enforcement of activities relating to underage tobacco and alcohol use and sales.




Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Pr to Die from 1999–	SMOKING,
Overall Men	7,850 4,625	State Lung Cancer All States	79.6/100,000 90.2/100,000	Overall Projected Death	136,373
Women Death Rate	3,225 278.2/100,000	State CHD* All States	58.8/100,000 59.7/100,000	Rate Projected All	9,963/100,000
All States	295.5/100,000	State COPD [†] All States	61.8/100,000 59.7/100,000	Štates	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

	GRADI	ES 6-8	GRADES	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
WI [†]	12.2%	16.1%	32.9%	39.4%
Boys [†]	11.9%	17.7%	31.0%	42.1%
Girls [†]	12.7%	14.4%	35.0%	36.2%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; † Wisconsin Youth Tobacco Survey, 2000.

Disparities Among Adult Population Groups, 2000*





Wisconsin is investing 24% of CDC's *Best Practices* lower estimated recommended funding and 9% of the upper estimated recommended funding. Wisconsin spent \$303 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$375,000,000 or \$722.80 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total Annual Per Capita	\$570,000,000 \$322,000,000 \$475,000,000 \$121,000,000 \$92,000,000 \$1,580,000,000 \$303
Annual Per Capita	\$303

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,413,000,000 \$269

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$123,248,769 Gross cigarette tax revenue collected in 2000 \$251,616,000

Cigarette tax per pack was \$0.77 in 2001

Cigarette sales were 80.1 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$5,532,300	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,532,300	
Federal—CDC Office on Smoking and Health	\$1,298,146	6/01-5/02
Federal—SAMHSA	\$76,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$77,330	10/00-9/01
Non-Government Source—RWJF/AMA	\$485,819	3/01–2/02
Subtotal: Federal/National Sources	\$1,937,295	
FY02 Total Investment in Tobacco Control		\$7,469,595
CDC Best Practices Recommended Annual Total (Lower E	\$31,158,000	
CDC Best Practices Recommended Annual Total (Upper E	\$82,381,000	
FY02 Per Capita Investment in Tobacco Control	\$1.38	

^{*}In FY02, \$500,000 was appropriated to the Medical College of Wisconsin for tobacco research. An additional \$9,304,000 was carried over from FY01 to be used for tobacco control in FY02.




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBU 1999 DISEASE-S	9	TO DIE FRO	Projected om Smoking, D-2000
Overall Men Women	729 434 295	State Lung Cancer All States State CHD*	82.2/100,000 90.2/100,000 50.3/100,000	Overall Projected Deatl Rate	12,974 n 10,067/100,000
Death Rate All States	318.6/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 114.2/100,000 59.7/100,000	Projected All States	8,830/100,000

^{*}Coronary heart disease; †chronic obstructive pulmonary disease.



Environmental Tobacco Smoke, 1998-99




Youth Tobacco Use

	GRADI	ES 6–8	GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
WY	14.8%†	21.7%†	35.2%††	43.9%††
Boys	15.6%†	25.8%†	34.6%††	49.6%††
Girls	13.4%†	16.9%†	35.9%††	38.0%††

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: *National Youth Tobacco Survey, 2000; †Wyoming Youth Tobacco Survey, 2000; †Wyoming Youth Risk Behavior Survey, 1999.

Disparities Among Adult Population Groups, 2000*





Wyoming is investing 28% of CDC's *Best Practices* lower estimated recommended funding and 14% of the upper estimated recommended funding. Wyoming spent \$221 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$29,000,000 or \$626.16 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$33,000,000
Hospital	\$29,000,000
Nursing Home	\$26,000,000
Prescription Drugs	\$10,000,000
Other	\$8,000,000
Annual Total	\$106,000,000
Annual Per Capita	\$221

Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$136,000,000 \$284

State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$14,772,038 Gross cigarette tax revenue collected in 2000 \$5,704,000

Cigarette tax per pack was \$0.12 in 2001

Cigarette sales were 90.5 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$900,000	7/01-6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$900,000	
Federal—CDC Office on Smoking and Health	\$973,302	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01-5/02
Subtotal: Federal/National Sources	\$1,173,302	
FY02 Total Investment in Tobacco Control	\$2,073,302	
CDC Best Practices Recommended Annual Total (Lower F	\$7,381,000	
CDC Best Practices Recommended Annual Total (Upper F	\$14,397,000	
FY02 Per Capita Investment in Tobacco Control	\$4.16	