National Youth Risk Behavior Survey: 1991-2003

## Trends in the Prevalence of Tobacco Use

## What Is the National Youth Risk Behavior Survey (YRBS)?

The national YRBS monitors priority health risk behaviors that contribute to the leading causes of death, disability, and social problems among youth and adults in the United States. The national YRBS is conducted every two years during the spring semester and provides data representative of 9th through 12th grade students in public and private schools throughout the United States.

1991	1993	1995	1997	1999	2001	2003	Changes from 1991 – 2003¹	Change from 2001 – 2003 <sup>2</sup>
		obacco use cco, snuff, o	r dip on on	e or more o	of the 30 da	ys preceding	the survey.)	
NA <sup>3</sup>	NA	11.4 ( <u>+</u> 1.7) <sup>4</sup>	9.3 ( <u>+</u> 2.2)	7.8 ( <u>+</u> 2.2)	8.2 ( <u>+</u> 1.5)	6.7 ( <u>+</u> 1.5)	Decreased, 1995 - 2003	No change
		ise on scho			of the 30 da	ys preceding	the survey.)	
NA	NA	6.3 ( <u>+</u> 1.0)	5.1 ( <u>+</u> 1.4)	4.2 ( <u>+</u> 1.8)	5.0 ( <u>+</u> 1.2)	5.9 ( <u>+</u> 3.0)	No change, 1995 - 2003	No change
Current o		rillos, or little	e cigars on	one or mor	e of the 30	days precedi	ng the survey.)	
NA	NA	NA	22.0 ( <u>+</u> 2.1)	17.7 ( <u>+</u> 1.7)	15.2 ( <u>+</u> 1.2)	14.8 ( <u>+</u> 1.7)	Decreased, 1997 - 2003	No change
	obacco use cigarettes o		ısed chewir	ng tobacco,	snuff, or di	p on one or r	more of the 30 days preceding	the survey.)
NA	NA	NA	43.4 ( <u>+</u> 2.4)	40.2 ( <u>+</u> 2.7)	33.9 ( <u>+</u> 2.1)	27.5 ( <u>+</u> 2.4)	No change, 1997 - 1999 Decreased, 1999 - 2003	Decreased

Based on linear and quadratic trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade.

## Where can I find more information?

More information about the YRBSS is available at <a href="https://www.cdc.gov/yrbss">www.cdc.gov/yrbss</a> or call (888)231-6405.



<sup>&</sup>lt;sup>2</sup> Based on T-test analyses.

<sup>&</sup>lt;sup>3</sup> Not available.

<sup>&</sup>lt;sup>4</sup> 95% confidence interval.