

Summary

Staging

Guide

**Cancer Surveillance
Epidemiology and
End Results Reporting**

**SEER Program
April 1977**

SUMMARY STAGING GUIDE

Edited by

Evelyn N. Shambaugh, M.A.
SEER Program
Demographic Analysis Section
Surveillance and Operations Research Branch
National Cancer Institute
Bethesda, Maryland

Mildred A. Weiss, B.A.
UCLA Tumor Registry
University of California at Los Angeles
Los Angeles, California

Medical Consultants

Robert F. Ryan, M.D., Chairman
Department of Surgery
Tulane University School of Medicine
New Orleans, Louisiana

Charles E. Platz, M.D.
Department of Pathology
College of Medicine
University of Iowa
Iowa City, Iowa

FOREWARD

For ease in coding, the SEER Program favors the expanded extent of disease schemes. They act as a guide to complete reporting, and they indicate the usual progression of disease patterns. Often, however, comparisons are necessary with historical or other series for which only general staging categories are available. Therefore, this staging guide was developed which summarizes the expanded extent of disease categories into three general staging groups (i.e., localized, regional, and distant). Sometimes a series is so small that only general categories produce enough cases for a meaningful analysis. Many conditions may be grouped together which may seem incongruous, but such groupings are unavoidable if the extent of disease codes are to be summarized into three or four staging categories. Stage categories are based on a combination of clinical observations and operative-pathological evaluations. The priority order is pathologic, operative, clinical.

In order to make the staging groups, insofar as we are able, consistent with categories developed by the American Joint Committee for Cancer Staging and End Results Reporting (AJC) and by the International Union Against Cancer (UICC) and to give consideration to the prognostic significance (survival probability) of various factors, a grouping subdividing the localized, regional, and distant categories was necessary. The subdivisions can always be summarized into the three familiar

stage categories--localized, regional, and distant, or other alternative groupings. For example, for colon and rectum "confined to the mucosa" is coded L₁. These cases can then be included with either the "in situ" or the "localized" stage of disease depending on the rules of the series with which the comparison is being made. For melanoma, localized cases are subdivided into four groups (L₁ through L₄) in order to indicate Clark's levels of invasion. Sites considered to be distant involvement by direct extension are coded D₁ as distinct from D₂ or D₃ which are reserved for metastatic involvement.

TABLE OF CONTENTS

FOR

SUMMARY STAGING GUIDE

	Page
Organization of Summary Staging Guide	
Site-Specific Staging Guides for:	
Upper lip (400, 403)	9
Lower lip (401, 404)	11
Commissure of lip (406)	13
Base of tongue (410)	15
Anterior 2/3 tongue (411-414, 416)	17
Parotid gland (420)	19
Submandibular gland (421)	21
Upper gum (430)	23
Lower gum and retromolar trigone (431, 436)	25
Floor of mouth (440, 441, 448, 449)	27
Cheek mucosa and vestibule of mouth (450, 451)	31
Hard palate (452)	33
Soft palate and uvula (453, 454)	35
Oropharynx (460-469)	39
Nasopharynx (470-473, 478-479)	43
Hypopharynx (480-483, 488-489)	45
Cervical or upper esophagus (500, 503)	47
Thoracic or middle esophagus (501, 504)	49

	Page
Abdominal or lower esophagus (502, 505)	51
Stomach, (excluding cardioesophageal junction) (510-516, 518-519)	53
Duodenum (520)	57
Jejunum and ileum (521, 522)	61
Cecum (534)	63
Ascending colon (536)	65
Transverse colon, including hepatic and splenic flexures (530, 531, 537)	59
Descending colon (532)	73
Sigmoid colon (533)	75
Rectosigmoid junction (540)	79
Rectum (541)	83
Anal canal and anus, NOS (542, 543)	87
Liver and intrahepatic bile duct (550, 551)	88
Gallbladder (560)	91
Extrahepatic bile duct (561, 562)	92
Head of pancreas (570)	93
Body and tail of pancreas (571, 572)	94
Larynx (610-612)	97
Bronchus and lung, excluding carina (622-625, 628-529)	99
Bone (700-709)	102
Malignant melanoma of skin (730-737, 841-844, 871-872, 874; histology 872 thru 879)	105
Skin other than melanoma (730-737)	107
Breast (740-749, 759)	111
Cervix uteri (800-801, 808-809)	115
Corpus uteri (820-821, 828)	117
Ovary (830)	119

TABLE OF CONTENTS

	Page
Fallopian tube (832)	120
Vagina (840)	121
Vulva, including skin of vulva (841-844)	123
Prostate (859)	125
Testis (860, 869)	129
Penis, including skin of penis (871, 872, 874)	130
Bladder (880-886, 888-889)	133
Kidney (renal) parenchyma (890)	135
Renal (kidney) pelvis (891)	136
Ureter (892)	138
Thyroid gland (939)	139
Lymph nodes and lymphoid tissue (960-969, 416, 460, 471, 491, 640, 692; histology 959 thru 969, 975)	141

ORGANIZATION OF SUMMARY STAGING GUIDE

I. Summary staging definitions

IN_SITU: Intraepithelial, noninvasive, noninfiltrating

LOCALIZED: Within organ

a. Invasive cancer confined to the organ

of origin

b. Intraluminal extension where specified

For example, intraluminal extension to

immediately contiguous segments of the

large bowel is coded L unless the

3

invaded segment has an identifiably

different pattern of lymph node drain-

age.

REGIONAL: Beyond the organ of origin

a. By direct extension to adjacent organs/tissues

b. To regional lymph nodes

c. (a) and (b)

DISTANT: Direct extension or metastasis

a. Direct continuity to organs other than above

b. Discontinuous metastasis

c. To distant lymph nodes

II. Site definitions

The International Classification of Diseases for Oncology (ICD-O) codes are included at the top of the page for each scheme, however, the first digit (1) has been dropped in compliance with the practice of the SEER program.

III. Summary staging guide

A. Site-Specific staging schemes (See pages 8 through 141)

B. Nonspecific staging scheme

In situ
Localized
Regional, direct extension only
Regional, nodes only
Regional, direct extension and regional nodes
Regional, NOS
Non-localized, NOS
Distant
Unstaged

This nonspecific staging scheme applies to the following primary sites:

ICD-O Number	<u>PRIMARY SITE CODE</u>
405	Mucosa of lips, NOS
408	Two or more categories of lip
409	Lips, NOS
415	Junctional zone of tongue
418	Two or more categories of tongue
419	Tongue, NOS
422	Sublingual gland
428	Two or more categories of major salivary glands
429	Major salivary gland, NOS
438	Two or more categories of gum
439	Gum, NOS
455	Palate, NOS
458	Two or more categories of other parts of mouth
459	Oral cavity

490 Pharynx, NOS
491 Waldeyer's ring, NOS
498 Neoplasms of lip, oral cavity and pharynx whose point of origin cannot be assigned to any one of the categories 40 through 48
499 Ill-defined sites in lip, oral cavity, and pharynx

508 Junctions of esophagus
509 Esophagus, NOS

510 Cardioesophageal junction (excluding cardia of stomach)

523 Meckel's diverticulum
528 Two or more categories of the small intestine
529 Small intestine, NOS

535 Appendix
538 Two or more categories of colon
539 Colon, NOS

548 Other parts of rectum

568 Two or more categories of gallbladder and extrahepatic bile ducts
569 Biliary tract, NOS

573 Pancreatic duct
574 Islets of Langerhans
578 Two or more categories of pancreas
579 Pancreas, NOS

580 Retroperitoneum
588 Specified parts of peritoneum
589 Peritoneum, NOS

590 Intestinal tract
598 Neoplasms of digestive organs and peritoneum whose points of origin cannot be assigned to any one of the categories 50- through 58-
599 Gastrointestinal tract, NOS

600-605, 608, 609
Nasal cavities, accessory sinuses, middle ear, inner ear

- 613 Laryngeal cartilage
618 Two or more categories of larynx
619 Larynx, NOS
- 620 Trachea
622 Carina (excluding main bronchus)
- 630 Parietal pleura
631 Visceral pleura
638 Two or more categories of pleura
639 Pleura, NOS
- 640-643, 648, 649
Thymus and mediastinum (excluding histology 959 thru 969)
- 650, 658, 659
Other and ill-defined sites within respiratory system and intrathoracic organs
- 690-691, 699
Blood, bone marrow (hematopoietic system)
- 692 Spleen (excluding histology 959 thru 969, 975)
- 693 Reticuloendothelial system, NOS
- 710, 712-719
Connective tissue and other soft tissue
- 738 Two or more categories of skin (melanotic and nonmelanotic)
739 Skin, NCS (melanotic and nonmelanotic)
- 799 Uterus, NOS
- 819 Placenta
- 833 Broad ligament
834 Parametrium
835 Round ligament
838 Other uterine adnexa
839 Uterine adnexa
- 848 Two or more categories of other and unspecified female genital organs
849 Female genital tract, NOS
- 873 Body of penis
875 Epididymis
876 Spermatic cord
877 Scrotum, NOS
878 Other parts of male genital organs
879 Male genital tract, NOS

887 Urachus

893 Urethra

894 Parau'rethral gland

898 Two or more categories of other urinary organs

899 Urinary system, NOS

900-909
Eye and lacrimal gland

910-919
Brain

920-923, 928, 929
Other and unspecified parts of nervous system

940, 941, 943-946, 948, 949
Other endocrine glands

950-955, 958 Other ill-defined sites

999 Unknown primary site

IV. Definition of Anatomic Sites Within The Oral Cavity
According To The American Joint Committee on Cancer Staging

LIPS, upper and lower, form the upper and lower anterior wall of the oral cavity. They consist of an exposed surface of modified epidermis commonly referred to as the vermillion surface, which extends from commissure to commissure and the mucous membrane lining the inner surface of the lips.

COMMISSURE OF LIP is the point of union of upper and lower lips (corner of mouth).

POSTERIOR ONE-THIRD OF TONGUE (base of tongue) consists of the less mobile portion of the tongue which extends inferiorly from the line of circumvallate papillae to the base of the epiglottis, the pharyngoepiglottic and glossoepiglottic folds (which bound the vallecula).

ANTERIOR TWO-THIRDS OF TONGUE consists of the freely movable portion of the tongue which extends anteriorly from the line of circumvallate papillae to the root of the tongue at the junction of the floor of the mouth. It is composed of four areas: (1) Tip, (2) Lateral borders, (3) Dorsum, and (4) Undersurface (non-villous surface).

FLOOR OF MOUTH consists of a semilunar shaped space over the mylohyoid and hypoglossus muscles, extending from the inner surface of the lower alveolar ridge to the root of the tongue. Its posterior boundary is the base of the anterior pillar of the tonsil. It is divided into two sides by the frenulum of the tongue and contains the ostia of the submaxillary and lingual salivary glands.

LOWER GINGIVA includes the alveolar process of the mandible and its covering mucosa, which extends from the line of attachment of mucosa in the buccal gutter to the line of free mucosa of the floor of the mouth. Posteriorly it extends to the ascending ramus of the mandible (retromolar trigone).

UPPER GINGIVA is the covering mucosa of the upper alveolar ridge, extending from the line of attachment of mucosa in the upper gingival buccal gutter to the junction with the hard palate. Its posterior margin is the upper end of the pterygopalatine arch.

BUCCAL MUCOSA includes all the mucous membrane lining the inner surface of the cheek.

HARD PALATE consists of the semilunar area between the upper alveolar ridges and the mucous membrane covering the palatine process of maxillary palatine bones. It extends from the inner surface of the superior alveolar ridge to the posterior edge of the palatine bone.

SOFT PALATE consists of mucosa covering the oral cavity side of the palatine muscles and extends from the posterior edge of the hard palate to the free border of the soft palate and includes the uvula. Its superior lateral margin is the pterygomandibular raphe. The inferior lateral margin completes the faucial arch (glossopalatine arch) and includes the anterior surface of the anterior tonsillar pillar.

NASOPHARYNX

Posterior Superior Wall (vault) extends from the superior border of the choana to the level of the free border of the soft palate. The lateral limit is the groove between the lateral wall and the base of the skull.

Lateral Wall extends from the base of the skull on each side to the level of the free border of the soft palate. It includes Rosenmuller's fossae (pharyngeal recesses).

OROPHARYNX

Posterior Wall extends from the free borders of the soft palate to the tip of the epiglottis.

Lateral Wall includes the tonsillar pillars, tonsillar fossae, and tonsils.

Anterior Wall consists of the lingual (anterior) surface of the epiglottis, and the pharyngoepiglottic and glossoepiglottic folds which bound the vallecula.


HYPOPHARYNX

Pyriform Sinus extends from the pharyngoepiglottic fold to the upper edge of the esophagus, between the inner surface of the thyroid cartilage and the posterior lateral surface of the arytenoid and cricoid cartilages.

Post-Cricoid Area extends from the posterior surface of the arytenoid cartilages and their connecting folds to the inferior surface of the cricoid. The lateral margin is the anterior part of the pyriform sinus.



Posterior Pharyngeal Wall extends from the level of the tip of the epiglottis to the inferior margin of the cricoid cartilage, and laterally to the posterior margins of the pyriform sinus.

VERMILION SURFACE OF UPPER LIP



REGIONAL NODES

1. Facial
2. Parotid
3. Submental
4. Submaxillary



IN-SITU; noninvasive

LOCALIZED

Vermilion surface
Skin of lip
Labial mucosa (inner lip)
Musculature
Multiple foci

Localized, NOS

REGIONAL, Direct Extension

R₁ if:
Commissure(s) of lips
Lower lip

R₂ if:
Buccal mucosa (inner cheek)
Gingiva, upper

R₃ if:
Maxilla
Nose

REGIONAL, Lymph Nodes

Facial: buccinator
Parotid: infra-auricular, preauricular
Submental
Submandibular (submaxillary)

DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes



Internal jugular
Upper cervical (including cervical, NOS)
Supraclavicular (transverse cervical)
Other distant nodes

LOWER LIP



REGIONAL NODES

1. Facial
2. Submental
3. Submandibular



LOWER LIP Revised April, 1982
401, 404

IN-SITU; noninvasive

LOCALIZED

Vermilion surface
Skin of lip
Labial mucosa (inner lip)
Musculature
Multiple foci

Localized, NOS

REGIONAL, Direct Extension

R₁ if:
Commissure(s) of lips
Upper lip

R₂ if:
Buccal mucosa (inner cheek)
Gingiva, lower

R₃ if:
Mandible

REGIONAL, Lymph Nodes

Facial: mandibular
Submental
Submandibular (submaxillary)

DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes




Internal jugular
Upper cervical (including cervical, NOS)
Supraclavicular (transverse cervical)
Other distant nodes

COMMISSURE OF LIPS



REGIONAL NODES

1. Facial
2. Parotid
3. Submental
4. Submandibular



COMMISSURE OF LIP Revised April, 1977
40¢

IN-SITU; noninvasive

LOCALIZED

Vermilion surface
Skin of lip
Labial mucosa (inner lip)
Musculature

Localized, NOS

REGIONAL, Direct Extension

R₁ if:
Both lips

R₂ if:
Buccal mucosa (inner cheek)
Gingiva

R₃ if:
Maxilla
Mandible
Nose

REGIONAL, Lymph Nodes



Facial: mandibular
Parotid: infra-auricular, presuricular
Submental
Submandibular (submaxillary)

DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes



Internal jugular
Upper cervical (including cervical, NOS)
Supraclavicular (transverse cervical)
Other distant nodes

BASE OF TONGUE



REGIONAL NODES

1. Submandibular
2. Upper cervical
3. Internal jugular



IN SITU: noninvasive

LOCALIZED

L if:

1

Confined to posterior 1/3 of tongue on one side

L if:

2

Midline tumor; tumor has crossed midline

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Anterior 2/3 of tongue

Gingiva, lower

Sublingual gland

R if:

2

Vallate, including pharyngoepiglottic and glossoepiglottic folds

Epiglottis, lingual (pharyngeal) surface

Floor of mouth

Lateral pharyngeal wall (tonsillar pillars, tonsillar fossae, tonsils)

REGIONAL, Lymph Nodes

Submandibular (submaxillary)

Internal jugular; subdigastric

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Mandible

Larynx

Hypopharynx

Soft palate, including uvula

D if:

2


Other distant involvement

DISTANT, Lymph Nodes

SuprACLAVICULAR (transverse cervical)




Other distant nodes

ANTERIOR $\frac{2}{3}$ OF TONGUE



REGIONAL NODES

1. Submandibular
2. Submental
3. Upper cervical
4. Internal jugular
5. Sublingual



IN SITU: noninvasive

LOCALIZED

L if:

1

Confined to anterior 1/3 of tongue on one side
with or without invasion of musculature

L if:

2

Midline tumor; tumor has crossed midline

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Floor of mouth

Base of tongue

Sublingual gland

R if:

2

Gingiva, lower

R if:

3

Mandible

REGIONAL, Lymph Nodes

Submandibular (submaxillary)

Submental

Internal jugular: subdigastric, supraomohyoid

Upper cervical (including cervical, NOS)

Sublingual

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Lateral pharyngeal wall (tonsillar pillars, tonsillar fossae, tonsils)

Soft palate, including uvula

Maxilla

D if:

2



Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)

Other distant nodes

PAROTID GLAND



IN SITU; noninvasive

LOCALIZED

Entirely within benign tumor capsule

Substance of parotid gland invaded

Multiple foci but confined to substance of parotid gland

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Periglandular soft tissue

Nerve(s): facial, auricular, spinal accessory

Skeletal muscle(s): digastric, sternocleidomastoid, masseter,
pterygoid, styloid

Periosteum of mandible

Pharyngeal mucosa

Submandibular (submaxillary) gland

R if:

2

Skin

R if:

3

Mandible

Major blood vessel(s): carotid artery, facial artery or vein,
maxillary artery, Jugular vein

Mastoid process

External auditory meatus

Skull

REGIONAL, Lymph Nodes

Parotid: intra-parotid, infra-auricular, preauricular

Submandibular (submaxillary)

DISTANT, Direct Extension or Metastasis


DISTANT, Lymph Nodes

Upper cervical (including cervical, NOS)




SuprACLAVICULAR (transverse cervical)

Other distant nodes

SUBMANDIBULAR GLAND



MUSCLES



REGIONAL NODES

1. Submandibular
2. Submental
3. Cervical, NOS
4. Upper jugular

IN SITU; noninvasive

LOCALIZED

Entirely within benign tumor capsule
Substance of parotid gland invaded
Multiple foci but confined to substance of submandibular gland

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Periglandular soft tissue

Skeletal muscle(s): digastric, mylohyoid, stylohyoid, hyoglossus,
styloglossus; pterygoid

Perosteum of mandible

Parotid gland

Sublingual gland

R if:

2

Nerve(s)

Major blood vessel(s): facial artery or vein, maxillary artery

Mandible

REGIONAL, Lymph Nodes

Submental

Submandibular (submaxillary)

Internal jugular (subdigastric)

Upper cervical (including cervical, NOS)



DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)


Other distant nodes

UPPER GUM (GINGIVA)



REGIONAL NODES

1. Facial
2. Submandibular
3. Retropharyngeal
4. Internal jugular
5. Upper cervical



IN SITU: noninvasive

LOCALIZED

L if:

1

Confined to mucosa

L if invading:

2

Lamina propria (mucoperiosteum)

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Maxilla

Hard and/or soft palate

Buccal mucosa (inner cheek)

Labial mucosa, upper lip

R if:

2

Lateral pharyngeal wall (tonsillar pillars, tonsillar fossae, tonsils)

Soft tissue of face

REGIONAL, Lymph Nodes

Facial: mandibular

Submandibular (submaxillary)

Retropharyngeal

Internal jugular

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Skin

Nasal cavity

Maxillary antrum (sinus)

Skull, including floor of orbit

D if:

2



Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)


Other distant nodes

LOWER GUM (GINGIVA)



REGIONAL NODES

1. Facial
2. Submandibular
3. Submental
4. Internal jugular
5. Upper cervical



LOWER GUM (GINGIVA) April, 1977
AND RETROMOLAR TRIGONE
431,456

IN SITU: noninvasive

LOCALIZED

L if:

1

Confined to mucosa

L if invading:

2

Lamina propria (mucoperiosteum)

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Mandible; periosteum of mandible

Floor of mouth

Buccal mucosa (inner cheek)

Labial mucosa, lower lip

Tongue

R if:

2

Lateral pharyngeal wall (tonsillar pillars, tonsillar fossae, tonsils)

Soft palate, including uvula

Soft tissue of face

REGIONAL, Lymph Nodes

Facial: mandibular

Submandibular (submaxillary)

Submental

Internal jugular: subdiaphragm, supraomohyoid

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Skin

Skull

D if:

2




Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)


Other distant nodes

FLOOR OF MOUTH



REGIONAL NODES

1. Submental
2. Submandibular
3. Sublingual
4. Internal jugular
5. Cervical, NOS



IN-SITU: noninvasive

LOCALIZED

L if:

1

Confined to mucosa on one side

L if:

2

Confined to one side:

Submucosa invaded

Musculature invaded

L if:

3

Midline tumor; tumor has crossed midline

L if:

x

Localized, NOS

|REGIONAL, Direct Extension

R if:

1

Gingiva, lower

Anterior 2/3 of tongue

Submandibular (submaxillary) gland(s)

Sublingual gland

Periosteum of mandible

R if:

2

Mandible

Base of tongue

Vallecula, including pharyngoeiglottic and glossoepiglottic folds

R if:

3

Epiglottis, pharyngeal (lingual) surface

Lateral pharyngeal wall (tonsillar pillars, tonsillar fossae, tonsils)

Underlying soft tissues

R if:

4

Skin

REGIONAL, Lymph Nodes





Submandibular (submaxillary)
Submental
Sublingual
Internal Jugular: subdigastric, supraomohyoid
Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)
Other distant nodes

CHEEK (BUCCAL) MUCOSA



REGIONAL NODES

- | | |
|------------------|---------------------|
| 1. Facial | 4. Submental |
| 2. Parotid | 5. Internal jugular |
| 3. Submandibular | 6. Upper cervical |

CHEEK (Buccal) NUCOSA April, 1977
VESTIBULE OF MOUTH
450,451

IN SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa
Submucosa invaded

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Soft tissue of cheek (including muscle)

Gingiva

Lip(s)

Lateral pharyngeal wall (tonsillar pillars, tonsillar fossae, tonsils)

R if:

2

Skin

REGIONAL, Lymph Nodes

Facial: buccinator, mandibular

Parotid: preauricular, infra-auricular

Submandibular (submaxillary)

Internal jugular: subdigastric

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Base or anterior 2/3 of tongue

Hard or soft palate

Bone: maxilla, mandible, skull

D if:

2




Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)


Other distant nodes

HARD PALATE



REGIONAL NODES

1. Submandibular
2. Retropharyngeal
3. Internal jugular
4. Cervical, NOS



IN SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa on one side

L if:

2

Midline tumor; tumor has crossed midline

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Soft palate, including uvula

Gingiva, upper

Palatine bone

Maxilla

R if:

2

Buccal mucosa (inner cheek)

REGIONAL, Lymph Nodes

Submandibular (submaxillary)

Retropharyngeal

Internal Jugular: subdiaphragm

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Nasal cavity; floor of nose

Maxillary antrum (sinus)

Nasopharynx

D if:

2




Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)


Other distant nodes

SOFT PALATE AND UVULA



REGIONAL NODES

1. Submandibular
2. Retropharyngeal
3. Internal jugular
4. Upper cervical



SOFT PALATE April, 1977
AND UVULA
453,454

IN SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa on one side

L if:

2

Submucosa and/or musculature invaded on one side

L if:

3

Midline tumor; tumor has crossed midline

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Hard palate, mucosa

Gingiva, upper

Lateral pharyngeal wall (tonsillar pillars, tonsillar fossae, tonsils)

R if:

2

Buccal mucosa (inner cheek)

Nasal cavity floor

REGIONAL, Lymph Nodes

Submandibular (submaxillary)

Retropharyngeal

Internal Jugular: subdigastric

Upper cervical (including cervical, NOS)

**SOFT PALATE April, 1977
AND UVULA
453,454**

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Tongue
Nasopharynx
Palatine bone
Maxilla
Maxillary antrum (sinus)
Mandible



D if:
2

Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)
Other distant nodes

OROPHARYNX



IN SITU; noninvasive

LOCALIZED (Tumor is not fixed)

L if confined to:

1

Posterior wall

One lateral wall

Anterior wall (including laryngeal (anterior) surface of epiglottis, vallecula epiglottis, and junctional region of oropharynx)

L if tumor involves:

2

Lateral wall(s) and posterior (or anterior) wall

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Tumor is not fixed, but extends into:

Soft tissue of neck

Prevertebral fascia

Base of tongue

Larynx

Pyriform sinus

Hypopharynx, NOS

Soft palate, including uvula

Nasopharynx

Floor of mouth

Gum (gingiva), posterior

Buccal mucosa (inner cheek)

R if:

2

Tumor is described as "fixed to adjacent tissues"

REGIONAL, Lymph Nodes

Retropharyngeal

Internal jugular: subdigastric, supraomohyoide

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Both lateral walls involved via soft palate or base of tongue

Anterior 2/3 of tongue

Hard palate

Mandible

Parotid gland

D if:

2

Other distant involvement



DISTANT, Lymph Nodes

Submandibular

Supraclavicular (transverse cervical)

Other distant nodes

NASOPHARYNX



IN SITU: noninvasive

LOCALIZED (Tumor is not fixed)

L if confined to:

1

Posterior superior wall (vault)

One lateral wall (including aryepiglottic fold, NOS)

L if tumor involves:

2

Posterior superior wall (vault) and lateral wall(s)

Lateral wall into eustachian tube/ middle ear

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Tumor is not fixed, but extends into:

Oropharynx; nasal cavity

Skull, including floor of orbit

Pterygopalatine fossa

Soft palate, including uvula

R if:

2

Tumor is described as "fixed to adjacent tissues"

REGIONAL, Lymph Nodes

Retropharyngeal

Internal jugular

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Brain, including cranial nerves

Accessory sinus: maxillary, sphenoid, ethmoid, frontal

Hard palate

Hypopharynx

Soft tissues of neck

D if:

2

Other distant involvement



DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)


Submandibular

Other distant nodes

HYPOPHARYNX



PHARYNX (posterior view)



IN_SITU: noninvasive

LOCALIZED (Tumor is not fixed)

L if confined to:

1

Pyriform sinus

Postcricoid area

Posterior pharyngeal wall

L if tumor involves:

2

Pyriform sinus and postcricoid area

Pyriform sinus and posterior pharyngeal wall

Postcricoid area and posterior pharyngeal wall

Pyriform sinus, postcricoid area and posterior pharyngeal wall

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Tumor is not fixed, but extends into:

Oropharynx; larynx

Soft tissues of neck

Prevertebral muscle(s)

Upper esophagus

R if:

2

Tumor is described as "fixed to adjacent tissues"

REGIONAL, Lymph Nodes

Retropharyngeal

Internal jugular: subdigastric, supraomohyoid

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Nasopharynx

Base of tongue

Floor of mouth

D if:

2






Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)

Other distant nodes

UPPER OR CERVICAL ESOPHAGUS



CERVICAL OR UPPER April, 1977
ESOPHAGUS
500, 503

IN-SITU: noninvasive

LOCALIZED

L if confined to:

1

Mucosa of upper esophagus

Mucosa but extends to middle esophagus

L if tumor invades:

2

Muscularis

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Adventitia and/or soft tissues of neck

Major blood vessel(s): carotid artery, subclavian artery,
Jugular vein

Thyroid gland

Esophagus is described as "fixed"

R if:

2

Hypopharynx; larynx

Trachea, including carina

Cervical vertebral(e)

REGIONAL, Lymph Nodes

Paraeosophageal

Internal Jugular

Anterior deep cervical: laterotracheal (recurrent)

Upper cervical (including cervical, NOS)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Main stem bronchus

Lung and/or pleura

D if:

2

Other distant involvement







DISTANT, Lymph Nodes

Posterior mediastinal

SuprACLAVICULAR (transverse cervical)

Other distant nodes

MIDDLE OR THORACIC ESOPHAGUS



THORACIC OR MIDDLE April, '1977
ESOPHAGUS
501, 504

IN SITU; noninvasive

LOCALIZED

L if confined to:

1

Mucosa of middle esophagus

Mucosa but extends to upper and/or lower esophagus

L if tumor involves:

2

Muscularis

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Adventitia and/or soft tissues

Major blood vessel(s): aorta, pulmonary artery or vein, vena cava

Trachea

Carina

Main stem bronchus

Esophagus is described as "fixed"

R if:

2

Lung via bronchus

Pleura

Pericardium

Mediastinal structure(s) NOS

Rib(s)

Thoracic vertebra(e)

Diaphragm

REGIONAL, Lymph Nodes

Paraesophageal

Tracheobronchial: peritracheal, carinal (bifurcation), hilar
(pulmonary roots)

Posterior mediastinal

Internal jugular

Left gastric: cardiac, lesser curvature

Upper cervical (including cervical, NOS)





DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes

Supraclavicular (transverse cervical)

Other distant nodes

LOWER OR DISTAL ESOPHAGUS



ABDOMINAL OR LOWER April, 1977
ESOPHAGUS
502, 505

IN SITU; noninvasive

LOCALIZED

L if confined to:

1

Mucosa of lower esophagus

Mucosa but extends to middle esophagus

L if tumor involves:

2

Muscularis

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Adventitia and/or soft tissues

Esophagus is described as "fixed"

R if:

2

Diaphragm

Cardia of stomach

Major blood vessel(s): aorta, gastric artery or vein, vena cava

REGIONAL, Lymph Nodes

Paraesophageal

Left gastric: cardiac, lesser curvature, perigastric, NOS

Posterior mediastinal

DISTANT, Direct Extension or Metastasis

D if:

1

"Diaphragm is fixed" (indicates phrenic nerve involved by tumor)

D if:

2

Other distant involvement






DISTANT, Lymph Nodes

Celiac

Para-aortic

Other distant nodes

STOMACH



STOMACH (excluding April, 1977
cardioesophageal junction)
510-516, 518-519

IN-SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa (lamina propria or muscularis mucosae; intranucosal)

L if:

2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:

3

Muscularis propria invaded

Invasion through muscularis propria

Subserosal tissue invaded (including extension through the wall, NOS)

L if:

4

Implants inside the stomach

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Perigastric fat

Lesser omentum

Ligaments: gastrocolic, gastrohepatic, gastrosplenic

Gastric artery

R if:

2

Invasion of (through) serosa*

Diffuse involvement of entire thickness of stomach wall
(linitis plastica)

R if extension to:

3

Esophagus (intraluminal)

Duodenum (intraluminal)

R if extension to:

4

Spleen

Omentum (greater)

Transverse colon (including hepatic and splenic flexures)

Diaphragm

*Invasion of serosa may be considered localized in historical comparisons.

STOMACH (excluding April, 1977
cardioesophageal junction)
510-516, 518-519

R if extension to:

5

Esophagus via serosa
Duodenum via serosa
Liver
Pancreas
Jejunum, ileum

REGIONAL. Lymph Nodes

Inferior gastric:

Gastrocolic
Gastroepiploic, right or NOS
Greater curvature
Greater omentum
Infrapyloric
Pyloric
Subpyloric

Superior gastric:

Cardiac
Cardioesophageal
Gastrohepatic
Left gastric
Lesser curvature
Lesser omentum
Paracardial

Splenic hilar:

Left gastroepiploic
Pancreaticolienal
Peripancreatic
Splenic

Perigastric, NOS

Nodule(s) in perigastric fat

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Left Kidney
Adrenal gland(s)
Retroperitoneum
Abdominal wall
Ovary (Krukenberg tumor)

D if:


2

Other distant involvement

DISTANT, Lymph Nodes

Celiac
Hepatic
Mesenteric, superior or inferior
Para-aortic
Portal
Retroperitoneal
Other distant nodes

DUODENUM



IN SITU; noninvasive

LOCALIZED

L if:

1

Invasive cancer confined to a polyp
Confined to submucosa

L if:

2

Muscularis and/or serosa invaded

L if:

3

Intraluminal to jejunum

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Periduodenal tissue
Mesentery, including mesenteric fat
Stomach
Extrahepatic bile duct(s), including ampulla of Vater
Pancreas, including pancreatic duct

R if:

2

Greater omentum

Major blood vessel(s): aorta, superior mesenteric artery or vein,
vena cava, portal vein, renal vein, gastro-
duodenal artery

R if:

3

Small intestine via serosa
Transverse colon, including hepatic flexure
Right and/or quadrate lobe of liver
Gallbladder
Right kidney
Right ureter
Diaphragm
Abdominal wall
Retroperitoneum

REGIONAL, Lymph Nodes





Hepatic: pancreaticoduodenal, infrapyloric, gastroduodenal

DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes

Superior mesenteric
Other distant nodes

JEJUNUM & ILEUM



IN SITU; noninvasive

LOCALIZED

L if:

1

Invasive cancer confined to a polyp
Confined to submucosa

L if:

2

Muscularis and/or serosa invaded

L if:

3

Intraluminal to ileocecal valve or cecum from ileum
Intraluminal to duodenum from jejunum

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Mesentery, including mesenteric fat

R if:

2

Abdominal wall

Retroperitoneum

Small intestine via serosa

Large intestine, including appendix

REGIONAL, Lymph Nodes

Posterior cecal (terminal ileum only)

Ileocolic (terminal ileum only)

Superior mesenteric

DISTANT, Direct Extension or Metastasis

Bladder

Uterus




Ovary

Fallopian tube

Other distant involvement

DISTANT, Lymph Nodes

CECUM & ASCENDING COLON



CECUM April, 1977
534

IN SITU: noninvasive

LOCALIZED

L if:

1

Confined to mucosa (lamina propria or muscularis mucosae;
intramucosal)

L if:

2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:

3

Muscularis propria invaded

Invasion through muscularis propria

Subserosal tissue invaded (including extension through the wall, NOS)

L if:

4

Intraluminal to appendix, cecum or ileocecal valve, ileum, ascending colon

Implants inside the cecum

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Mesentery, including mesenteric fat

Pericolic (pericecal) fat

Adjacent tissue(s), NOS

R if:

2

Invasion of (through) serosa*

R if extension to:

3

Greater omentum

Retroperitoneum

Abdominal wall

Small intestine, other than ileum

*Invasion of serosa may be considered localized in historical comparisons.

REGIONAL, Lymph Nodes

Epicolic
Paracolic
Cecal
Ileocolic
Right colic (including colic, NOS)
Mesenteric, superior or NOS

Nodule(s) in pericolic fat

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Uterus
Ovary
Fallopian tube
Urinary bladder
Gallbladder
Right kidney or ureter
Liver
Other segment of colon via serosa

D if:
2

Other distant involvement

DISTANT, Lymph Nodes

Inferior mesenteric
Para-aortic
Retroperitoneal
Other distant nodes

IN SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa (lamina propria or muscularis mucosae;
intramucosal)

L if:

2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:

3

Muscularis propria invaded

Invasion through muscularis propria

Subserosal tissue invaded (including extension through the
wall, NOS)

L if:

4

Intraluminal to cecum, appendix, ileocecal valve, transverse colon

Implants inside the ascending colon

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Pericolic fat

Retroperitoneal fat

Adjacent tissue(s), NOS

R if:

2

Invasion of (through) serosa*

R if extension to:

3

Greater omentum

Retroperitoneum

Abdominal wall

Small intestine

Right ureter

Right kidney

Liver, right lobe

*Invasion of serosa may be considered localized in historical comparisons.

REGIONAL, Lymph Nodes

Epicolic
Paracolic
Ileocolic
Right colic (including colic, NOS)
Middle colic
Mesenteric, superior or NOS

Nodule(s) in pericolic fat

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Uterus
Ovary
Fallopian tube
Urinary bladder
Gallbladder
Other segment of colon via serosa





D if:
2

Other distant involvement

DISTANT, Lymph Nodes

Inferior mesenteric
Para-aortic
Retroperitoneal
Other distant nodes

**HEPATIC FLEXURE,
TRANSVERSE COLON &
SPLENIC FLEXURE**



TRANSVERSE COLON April, 1977
(Including flexures)
530, 531, 537

IN SITU: noninvasive

LOCALIZED

L if:

1

Confined to mucosa (lamina propria or muscularis mucosae;
intramucosal)

L if:

2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:

3

Muscularis propria invaded

Invasion through muscularis propria

Subserosal tissue invaded (including extension through the
wall, NOS)

L if:

4

Intraluminal to ascending or descending colon

L if:

X

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Mesentery (including mesenteric fat); mesocolon

Pericolic fat

Greater omentum; gastrocolic ligament

Adjacent tissue(s), NOS

R if:

2

Invasion of (through) serosa*

R if extension to:

3

Stomach

Small intestine

Liver

Spleen

Pancreas

Retropertitoneum

Gallbladder/ bile ducts

Kidney

Abdominal wall

*Invasion of serosa may be considered localized in historical comparisons.

REGIONAL, Lymph Nodes

Epicolic

Paracolic

Right colic for hepatic flexure only

Middle colic

Colic, NOS

Left colic for splenic flexure only

Inferior mesenteric for splenic flexure only

Superior mesenteric for hepatic flexure and transverse
colon only

Mesenteric, NOS

Nodule(s) in pericolic fat

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Other segment of colon via serosa

Diaphragm

Ureter

Adrenal gland

Ovary

D if:

2





Other distant involvement

DISTANT, Lymph Nodes

Para-aortic or retroperitoneal

Inferior mesenteric for hepatic flexure and transverse colon only
Superior mesenteric for splenic flexure only

**DESCENDING COLON &
SIGMOID COLON**



IN-SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa (lamina propria or muscularis mucosae)

L if:

2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:

3

Muscularis propria invaded

Invasion through muscularis propria

Subserosal tissue invaded (including extension through the wall, NOS)

L if:

4

Intraluminal to splenic flexure, transverse colon, sigmoid colon

L if:

X

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Pericolic fat, NOS

Retroperitoneal fat

Adjacent tissue(s), NOS

R if:

2

Invasion of (through) serosa*

R if extension to:

3

Small intestine

Retroperitoneum

Greater omentum

Abdominal or pelvic wall

Left ureter

Left kidney

Spleen

*Invasion of serosa may be considered localized in historical comparisons.

REGIONAL, Lymph Nodes

Epicolic

Paracolic

Left colic (including colic, NOS)

Mesenteric, inferior or NOS

Nodule(s) in pericolic fat

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Uterus

Ovary

Fallopian tube

Other segment of colon via serosa

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

Para-aortic

Retroperitoneal

Superior mesenteric

Other distant nodes

IN SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa (lamina propria or muscularis mucosae;
intramucosal)

L if:

2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:

3

Muscularis propria invaded

Invasion through muscularis propria

Subserosal tissue invaded (including extension through the
wall, NOS)

L if:

4

Intraluminal to descending colon, rectosigmoid, or rectum

L if:

X

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Mesentery (including mesenteric fat); mesosigmoid

Pericolic fat

Adjacent tissue(s), NOS

R if:

2

Invasion of (through) serosa*

R if extension to:

3

Greater omentum

Abdominal or pelvic wall

Small intestine

*Invasion of serosa may be considered localized in historical comparisons.

REGIONAL, Lymph Nodes

Epicolic
Paracolic
Colic, NOS
Sigmoidal
Superior hemorrhoidal
Superior rectal
Mesenteric, inferior or NOS

Nodule(s) in pericolic fat

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Uterus
Cul de sac (rectouterine pouch)
Ovary
Fallopian tube
Ureter
Urinary bladder
Other segment of colon via serosa





D if:
2

Other distant involvement

DISTANT, Lymph Nodes

Para-aortic
Retroperitoneal
Superior mesenteric
Other distant nodes

RECTOSIGMOID



IN SITU: noninvasive

LOCALIZED

L if:
1

Confined to mucosa (lamina propria or muscularis mucosae;
intramucosal)

L if:
2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:
3

Muscularis propria invaded

Invasion through muscularis propria

Subserosal tissue invaded (including through the wall, NOS)

L if:
4

Intraluminal to sigmoid colon or rectum

L if:
X

Localized, NOS

REGIONAL, Direct Extension

R if extension to:
1

Mesentery, including mesenteric fat

Pericolic (perirectal) fat

Adjacent tissue(s), NOS

R if:
2

Invasion of (through) serosa*

R if extension to:
3

Small intestine

Cul de sac (rectouterine pouch)

Pelvic wall/ pelvic plexuses

*Invasion of serosa may be considered localized in historical comparisons.

REGIONAL, Lymph Nodes

Paracolic (including colic, NOS)
Pararectal
Hemorrhoidal, superior or middle
Sigmoidal
Internal iliac (hypogastric)
Mesenteric, inferior or NOS

Nodule(s) in pericolic fat

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Uterus
Vagina
Urinary bladder and/or ureter
Prostate
Skeletal muscles of pelvic floor
Fallopian tube
Ovary
Other segment of colon via serosa







D if:
2

Other distant involvement

DISTANT, Lymph Nodes

Para-aortic
Retroperitoneal
Superior mesenteric
Other distant nodes

RECTUM



IN SITU; noninvasive

LOCALIZED

L if:

1

Confined to mucosa (lamina propria or muscularis mucosae)

L if:

2

Submucosa invaded (thru muscularis mucosae)

Stalk invaded (if polyp)

Superficial invasion

L if:

3

Muscularis propria invaded

Invasion through muscularis propria (including extension
through wall, NOS)

L if:

X

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Perirectal fat

Rectovaginal septum

Adjacent tissue(s), NOS

R if:

2

Invasion of (through) serosa*

R if extension to:

3

Intraluminal to rectosigmoid or anus

R if extension to:

4

Colon, anus (except intraluminal)

Vagina

Cul de sac (rectouterine pouch)

*Invasion of serosa may be considered localized in historical comparisons.

R if extension to:

5

Urinary bladder/rectovesical fascia, male
Prostate
Seminal vesicle
Ductus deferens
Skeletal muscle of pelvic floor
Pelvic wall

REGIONAL, Lymph Nodes

Pararectal
Hemorrhoidal, superior or middle
Sacral
Sigmoidal
Mesenteric, inferior or NOS
Internal iliac (hypogastric)

Nodule(s) in perirectal fat

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Uterus
Urinary bladder, female
Sacrum
Sacral plexus
Bones of pelvis
Ovary
Urethra
Perineum; perianal skin

D if:

2







Other distant involvement.

DISTANT LYMPH NODES

Para-aortic
Retroperitoneal
Superior mesenteric

Inguinal
Other distant nodes

**ANUS, ANAL CANAL &
ANAL MUCOSA**



ANAL CANAL April, 1977
ANUS, NOS
542, 543

IN SITU; noninvasive

LOCALIZED

L if:

1

Incidental finding of malignancy in hemorrhoid
Invasive cancer confined to submucosa

L if:

2

Muscularis (internal sphincter)

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Rectal mucosa or submucosa

Perianal skin

Skeletal muscle(s): anal sphincter (external), levator ani

Ischiorectal fat/tissue

R if:

2

Perineum

Vulva

REGIONAL, Lymph Nodes

Anorectal; pararectal

Internal iliac (hypogastric) for anal canal only

Lateral sacral for anal canal only

Superficial inguinal for anus only

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Prostate

Peritoneum of pelvic floor

Bladder

Urethra

Vagina

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

NOTE: Melanoma of the anus is classified according to the staging scheme for melanoma.

LIVER April, 1977
INTRAHEPATIC BILE DUCTS
550, 551

LOCALIZED

Confined to one lobe

Satellite nodule(s) confined to one lobe

Localized, NOS

REGIONAL, Direct Extension

R if:

1

More than one lobe involved by contiguous growth

Gallbladder from right lobe of liver

R if:

2

Extrahepatic blood vessel(s): hepatic artery, vena cava, portal vein
Extrahepatic bile duct(s)

Diaphragm

Peritoneum

Ligament(s): falciform, coronary, triangular, hepatogastric,
hepatoduodenal

Lesser omentum

REGIONAL, Lymph Nodes

Cardiac

Diaphragmatic: pericardial

Posterior mediastinal

Hepatic: hepatic pedicle, inferior vena cava, hepatic artery

Lateral aortic (retroperitoneal): coronary, renal artery

DISTANT, Direct Extension or Metastasis

D if:

1

Satellite nodules in more than one lobe of liver, surface or parenchymal

D if extension to:

2

Pleura

Pancreas

Stomach

D if:

3

Other distant involvement

DISTANT, Lymph Nodes

LYMPHATICS OF THE GALLBLADDER




Fig. 504. Schematic representation of the lymphatics of the gallbladder showing the pathways to, 1, cystic node; 2, node of anterior border of foramen of Winslow; and, 3, superior retropancreatic duodenal node.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

IN SITU; noninvasive

LOCALIZED

L if:
1

Invasive cancer confined to submucosa

L if:
2

Muscularis and/or serosa invaded

L if:
x

Localized, NOS

REGIONAL, Direct Extension

R if:
1

Right lobe of liver

Gallbladder is replaced by tumor (indicate extension to liver)

Extrahepatic bile duct(s), including ampulla of Vater

R if:
2

Blood vessels: cystic artery/vein, hepatic artery, portal vein

Pancreas via extrahepatic bile ducts

Greater omentum

Lesser omentum

Duodenum

REGIONAL, Lymph Nodes

Cystic (node of the neck of the gallbladder)

Node of the foramen of Winslow

Hepatic: periportal, pancreaticoduodenal

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Pancreas (other than above)

Large intestine

Stomach

D if:
2

Other distant involvement

DISTANT, Lymph Nodes

Mesenteric

Para-aortic

Other distant nodes

EXTRAHEPATIC BILE DUCT April, 1977
AMPULLA OF VATER
561, 562

IN SITU: noninvasive

LOCALIZED

Confined to bile duct(s): cystic, hepatic, common, ampulla of Vater

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Duodenum

Gallbladder

Pancreas

Liver, porta hepatis

Lesser omentum

R if:

2

Transverse colon, including hepatic flexure

Distal stomach

Blood vessels: portal vein, hepatic artery

REGIONAL, Lymph Nodes

Cystic (node of the neck of the gallbladder)

Node of the foramen of Winslow

Hepatic: periportal, pancreaticoduodenal

DISTANT, Direct Extension or Metastasis

DISTANT, Lymph Nodes

Mesenteric

Para-aortic

Other distant nodes

LOCALIZED

Confined to head of pancreas
Body of pancreas involved
With obstruction, but no invasion, of extrahepatic bile duct(s)

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Extrahepatic bile duct(s), including ampulla of Vater

Duodenum

Stomach adjacent to head of pancreas, including stomach NOS

R if:

2

Liver

Major blood vessel(s): hepatic, pancreaticoduodenal and/or gastro-duodenal arteries, superior mesenteric artery or vein, portal vein

Transverse colon, including hepatic flexure

Peritoneum

Mesentery, mesocolon, mesenteric fat

Greater and/or lesser omentum

Gallbladder

R if:

3

Tumor is described as "fixed to adjacent tissues"

REGIONAL, Lymph Nodes

Peripancreatic

Hepatic: pancreaticoduodenal, infrapyloric (subpyloric)

Superior mesenteric

Lateral aortic (retroperitoneal)

Celiac

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Body of stomach

Kidney

Ureter

Adrenal gland

Retroperitoneum

Jejunum

Ileum

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

BODY AND TAIL
OF PANCREAS April, 1977
571, 572

LOCALIZED

Confined to body and/or tail of pancreas
Head of pancreas involved

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Left kidney

Left ureter

Left adrenal gland

Retroperitoneal soft tissues (retroperitoneal space)

R if:

2

Spleen

Stomach

Liver, porta hepatis

Gallbladder

Small intestine

Splenic flexure

Peritoneum

Mesentery, mesocolon, mesenteric fat

Major blood vessel(s): aorta, celiac artery, hepatic artery, splenic artery or vein, superior mesenteric artery or vein, portal vein

REGIONAL, Lymph Nodes

Splenic: suprapancreatic, splenic hilum, pancreaticolicinal

Superior mesenteric

Lateral aortic (retroperitoneal)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Right kidney

Right ureter

Right adrenal gland

Diaphragm

Large intestine (other than splenic flexure)

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

LYMPH NODES OF THE PELVIS AND ABDOMEN




FIG. 5. PRINCIPAL LYMPH NODE GROUPS OF THE PELVIS AND ABDOMEN

Source: Taylor, G. W., Lymph Node Metastases, p. 44

**DEFINITION OF ANATOMICAL LIMITS OF THE LARYNX ACCORDING
TO THE AMERICAN JOINT COMMITTEE ON CANCER STAGING**

Anterior Limit is constituted by the anterior or lingual surface of the suprathyroid epiglottis, thyrohyoid membrane, the anterior commissure, and the anterior wall of the subglottic region, which is composed of the thyroid cartilage, the cricothyroid membrane, and the anterior arch of the cricoid cartilage.

Posterior Lateral Limits include the aryepiglottic folds, the arytenoid region, the interarytenoid space, and the posterior surface of the subglottic space represented by the mucous membrane covering the cricoid cartilage.

Superior Lateral Limits are constituted by the tip and the lateral border of the epiglottis.

Inferior Limits are constituted by a plane passing through the inferior edge of the cricoid cartilage.

The larynx is divided into the following anatomical regions and sites:

Region	Site
Supraglottic	Ventricular bands (false cords), right and left Arytenoids, right and left Epiglottis (both lingual and laryngeal aspects) Suprathyroid epiglottis Infrathyroid epiglottis Aryepiglottic folds
Glottic	True vocal cords, right and left Anterior and posterior commissures
Subglottic	Right and left walls of the subglottis, exclusive of the undersurface of the cords

LARYNX April, 1977
(Excluding cartilage)
610-612, 618, 619

IN SITU; noninvasive

LOCALIZED

L if:

1

Tumor limited to one area within a region

Supraglottic region

Laryngeal (posterior) surface of epiglottis

Arytenoid

Aryepiglottic fold

Ventricular band (false cord, vestibular fold)

Ventricular cavity

Glottic region (normal mobility)

Vocal cord, one side

Commissure

Subglottic region on one side

L if:

2

Tumor extends to adjacent area(s) within a region

Supraglottic region

More than one of the above areas

Glottic region (normal mobility)

Cord and commissure

Both vocal cords

Subglottic region on both sides

L if:

3

Glottic region: Fixation of cord(s)

L if:

4

Tumor involves adjacent region(s)

Supraglottic region

Glottic region (with or without fixation)

Subglottic region

Involves intrinsic muscle(s): aryepiglottic, arytenoid, cricoarytenoid,
cricothyroid, thyroepiglottic, thyroarytenoid,
vocalis

L if:

x

Localized, NOS

LARYNX April, 1977
(Excluding cartilage)
610-612, 618, 619

REGIONAL, Direct Extension

R if:

1

Pyriform sinus

Posterior laryngeal area

Hypopharynx NOS

Vallate fold, including pharyngoepiglottic and glossoepiglottic folds

Base of tongue from laryngeal surface of epiglottis

R if:

2

Extends into cricoid and/or thyroid cartilage

REGIONAL, Lymph Nodes

Internal jugular: subdigastric

Anterior deep cervical: prelaryngeal, pretracheal, laterotracheal
(recurrent)

Cervical, NOS

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Extrinsic muscle(s): omohyoid, sternohyoid, sternothyroid, thyrohyoid
(strap muscles)

Soft tissues of neck

Thyroid

Skin

Trachea

Upper esophagus

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

Supraclavicular

Submandibular

Other distant nodes

BRONCHUS AND LUNG April, 1977
(Excluding carina)
622-625, 628-629

IN SITU; noninvasive

LOCALIZED

Single tumor \geq 2 cm. from carina and confined to one lung
and/or main stem bronchus

L if:

1

Tumor size is 3.0 cm. or less

L if:

2

Tumor size $>$ 3.0 cm.

L if:

3

Tumor size is unknown

L if:

4

Single tumor of any size $<$ 2 cm. from carina and confined to one
lung and/or main stem bronchus

L if:

5

Multiple masses confined to one lung and/or main stem bronchus

L if:

x

Localized, NOS

BRONCHUS AND LUNG April, 1977
(Excluding carina)
622-625, 628-629

REGIONAL, Direct Extension

R if extension to:

1

Pleura, visceral/NOS

Pericardium, parietal/NOS

Pulmonary ligament

R if tumor involves:

2

Carina

Trachea

Esophagus

Nerve(s):

Recurrent laryngeal

Vagus

Phrenic

Cervical sympathetic (Horner's syndrome)

Major blood vessel(s):

Pulmonary artery or vein

Azygos vein

Superior vena cava

R if:

x

Extrapulmonary mediastinal extension, NOS

REGIONAL, Lymph Nodes

Intrapulmonary

Hilar (bronchial; parabronchial; pulmonary root)

Subcarinal; carinal

Mediastinal (paratracheobronchial; parastracheal; pericardial;
para-esophageal; para-aortic-above diaphragm)

BRONCHUS AND LUNG April, 1977
(Excluding carina)
622-625, 628-629

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Brachial plexus from superior sulcus or Pancoast tumor
Lung and/or main stem bronchus, contralateral
Pericardium, visceral
Heart
Pleura, parietal

D if extension to:

2

Rib, sternum, vertebra
Chest (thoracic) wall
Skeletal muscle
Skin of chest
Diaphragm
Abdominal organs

D if:

3

Other distant involvement

DISTANT, Lymph Nodes

Contralateral hilar or mediastinal (including bilateral)
Supraclavicular (transverse cervical)
Scalene
Cervical, NOS
Other distant nodes

LOCALIZED

Confined to bone
Tumor has broken through periosteum but not beyond
Abnormal configuration of bone

Localized, NOS

REGIONAL, Direct Extension

Surrounding tissues, including skeletal muscle(s)
Adjacent bone

REGIONAL, Lymph Nodes

First chain of nodes involved in the area of the tumor

DISTANT, Direct Extension or Metastasis

Skin
Other distant involvement

DISTANT, Lymph Nodes

LYMPHATICS OF THE SKIN




Fig. 100 Lymphatics of the skin of the face and upper and lower extremities.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

MALIGNANT MELANOMA OF SKIN Revised July, 1983
730-737, 841-844, 871-872, 874
HISTOLOGY: 872 thru 879

IN SITU; intraepidermal (Clark's Level 1)

LOCALIZED

L₁ if tumor invades:
Papillary dermis (Clark's Level 2)
OR
Thickness/Depth of invasion ≤ 0.75 mm

L₂ if tumor invades:
Papillary-reticular dermal interface (Clark's Level 3)
OR
Thickness/Depth of invasion 0.76 - 1.50 mm

L₃ if tumor invades:
Reticular dermis (Clark's Level 4)
OR
Thickness/Depth of invasion >1.50 mm

L₄ if:
Subcutaneous tissue (through entire dermis) (Clark's Level 5)

L_X if:
Localized, NOS; confined to skin/dermis, NOS

REGIONAL

R₁ if:
Satellite nodule(s) within immediate area (\leq 2 cm from the primary lesion)

R₂ if:
Intransit metastasis directed toward regional lymph nodes
(including satellite nodule(s) $>$ 2 cm from the primary lesion)

NOTE 1: Skin ulceration does not alter the classification.

NOTE 2: Clark's level takes precedence over thickness/depth of invasion in case of discrepancy

MALIGNANT MELANOMA OF SKIN Revised July, 1983
730-737, 841-844, 871-872, 874
HISTOLOGY: 872 thru 879

REGIONAL, Lymph Nodes (by primary site)

Parotid: preauricular, infra-auricular	Axillary
Forehead	Arm
Temporal region	Hand
Malar region	Shoulder
Lateral half of eyelids	Chest wall, anterior and posterior
Outer canthus	Scapula (upper back), below transverse line
Anterior half of ear	
Submandibular (submaxillary)	
Midline of forehead	Epitrochlear
Medial half of eyelids	Hand
Inner canthus	Forearm
Nose	
Cheeks, Nose	
Cervical	
Occipital scalp, posterior ear	Superficial inguinal
Head and neck tumors, any location	Lumbar region (lower back)
Scapula, above transverse line	Abdominal wall, anterior and posterior
Supraclavicular (transverse cervical)	Perineum
Chest wall, anterior and posterior	Perianal region
Neck	
Popliteal	
Heel	Femoral
Posterior leg	Lower extremities (excluding heel)
	Perineum
	Perianal region

DISTANT, Direct Extension or Metastasis

D₁ if:

Underlying cartilage, bone, muscle

D₂ if:

Metastatic (generalized) skin lesions

D₃ if:

Other distant involvement

DISTANT, Lymph Nodes

Other than above

SKIN OTHER THAN MELANOMA Revised July, 1983
730-737

**NOTE: For skin of vulva, use 841-844 schemes;
for penis, use 871, 872, 874 schemes.**

IN SITU; intraepidermal

LOCALIZED

**Single lesion confined to dermis
Skin ulceration
Subcutaneous tissues (through entire dermis)
Localized, NOS**

REGIONAL. Lymph Nodes (by primary site)

Parotid: preauricular, infra-auricular Axillary

Forehead	Arm
Temporal region	Hand
Malar region	Shoulder
Lateral half of eyelids	Chest wall, anterior and posterior
Outer canthus	Scapula (upper back), below transverse line
Anterior half of ear	

Submandibular (submaxillary) Epitrochlear

Midline of forehead	Hand
Medial half of eyelids	Forearm
Inner canthus	
Nose	Superficial inguinal
Lips	
Cheeks	Lumbar region (lower back)

Cervical

Occipital scalp, posterior ear	Abdominal wall, anterior and posterior
Head and neck tumors, any location	Perineum
Scapula, above transverse line	Perianal region

SuprACLAVICULAR (transverse cervical)

Chest wall, anterior and posterior	Lower extremities (excluding heel)
Neck	Perineum
	Perianal region

**Popliteal
Heel
Posterior leg**

SKIN OTHER THAN MELANOMA Revised July, 1983
730-737

DISTANT. Direct Extension or Metastasis

D₁ if:

Underlying cartilage, bone, and muscle

D₂ if:

Metastatic (generalized) skin lesions

D₃ if:

Other distant involvement

DISTANT, Lymph Nodes

Other than above

LYMPHATICS OF THE BREAST




Fig. 669. Lymphatics of the breast leading to, 1, axillary nodes which are distributed over a large area from the lateral aspects of the breast proper to the axillary vessels; 2, interpectoral chain leading to interpectoral node (circle detail) and to high nodes in the axilla; 3, chain of the internal mammary leading frequently to node in second interspace and to supraclavicular and cervical nodes. The lymphatics of the breast may empty into the opposite axillary nodes.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

BREAST Revised July, 1983
740-749 Female
759 Male

IN SITU (including noninfiltrating, intraductal without infiltration)

LOCALIZED

Confined to breast, including nipple and/or areola:

- L₁ if tumor size is < 2 cm
- L₂ if tumor size is 2-4.9 cm
- L₃ if tumor size is ≥ 5 cm
- L₄ if tumor size is unknown
- L_x Localized, NOS

Note: Skin changes such as dimpling, tethering, attachment, fixation, induration and thickening or Paget's disease of nipple do not alter the classification.

REGIONAL. Direct Extension

- R₁ if tumor of any size with:

- Invasion of subcutaneous tissue
- Skin infiltration of primary breast
- Skin edema, peau d'orange, "pisskin"
- En curraise, lenticular nodules
- Inflammation of skin, erythema
- Ulceration of skin of breast
- Satellite nodules in skin of primary breast
- Pectoral fascia or pectoral muscle involvement

- R₂ if:

- Invasion of (or fixation to) chest wall, ribs, intercostal or serratus anterior muscles

BREAST Revised July, 1983
740-749 Female
759 Male

REGIONAL. Lymph Nodes

Axillary (low, adjacent to tail of breast;
mid, central, interpectoral, Rotter's node;
high, subclavicular, axillary vein nodes, apical)
Internal mammary (parasternal)

Nodules in axillary fat

DISTANT, Direct Extension or Metastasis

D₁ if
Skin over sternum, upper abdomen, axilla or opposite breast
Satellite nodule(s) in skin other than primary breast


D₂ if tumor involves:
Breast, contralateral

D₃ if:
Other distant involvement

DISTANT, Lymph Nodes

Infraclavicular
Supraclavicular (transverse cervical)
Cervical, NOS
Axillary and/or internal mammary, contralateral
Other distant nodes

LYMPHATICS OF THE UTERUS



V.S. ACKERMAN

Fig. 619. Schematic representation of the lymphatics of the uterus showing, 1, the utero-ovarian pedicle; 2, the external iliac pedicle; and, 3, the round ligament pedicle leading to the inguinal lymph nodes.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

IN-SITU

Non-invasive, pre-invasive, intraepithelial
League of Nations Stage 0

LOCALIZED

L if:

1

Minimal stromal invasion; "microinvasion"

L if:

2

Invasive cancer confined to cervix

L if:

x

League of Nations Stage I

REGIONAL. Direct Extension

R if extension to:

1

Corpus uteri (body of uterus)

R if extension to:

2

Upper 2/3 of vaginal wall (including fornices and vagina, NOS)
Parametrium

Ligaments: broad, uterosacral, cardinal

League of Nations Stage II

R if extension to:

3

Lower 1/3 of vaginal wall

Pelvic wall(s)

League of Nations Stage III

R if extension to:

4

Rectal and/or bladder wall (excluding mucosa)
Bullous edema of bladder mucosa
Cul de sac (rectouterine pouch)

REGIONAL, Lymph Nodes

Hypogastric
Iliac (common, internal, external)
Obturator
Paracervical
Parametrial/pelvic, NOS
Sacral (laterosacral, presacral, sacral promontory, uterosacral)

DISTANT, Direct Extension or Metastasis

D if extension to:

Bladder mucosa
Rectal mucosa
Ureter
Urethra
Sigmoid colon
Small intestine
Vulva
Ovary and/or fallopian tube

D if:

2

"Frozen pelvis"

D if:

3

Other distant involvement

D if:

x

League of Nations Stage IV

DISTANT, Lymph Nodes

Aortic (para-aortic, periaortic, lumbar)
Inguinal
Other distant nodes

IN-SITU, pre-invasive, non-invasive

LOCALIZED

L if:

1

Invasive cancer confined to endometrium

L if:

2

Myometrium/serosa (perimetrium) invaded

**Invasive cancer confined to corpus, clinically:
(Depth of invasion unknown)**

L if:

3

Sounding of uterine cavity is \leq 8 cm. from cervical os

L if:

4

Sounding of uterine cavity is $>$ 8 cm. from cervical os

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if extension to:

1

Cervix uteri, including endocervix

R if extension to:

2

Parametrium

Ligaments: broad, round, uterosacral

R if extension to:

3

Pelvic wall(s)

Ovary and/or fallopian tube(s)

R if extension to:

4

Rectal and/or bladder wall (excluding mucosa)

REGIONAL, Lymph Nodes

Hypogastric

Iliac (common, internal, external)

Obturator

Paracervical

Parametrial/pelvic, NOS

Sacral (lateral sacral, sacral promontory, presacral, uterosacral)

Superficial inguinal

Lateral aortic, preaortic

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Vagina

Vulva

Cul de sac (rectouterine pouch)

Rectum or bladder mucosa

Ureter

Sigmoid colon

Small intestine

Serosa of abdominal organs

D if:

2

"Frozen pelvis"

D if:

3

Other distant involvement

DISTANT, Lymph Nodes

OVARY April, 1977
830

LOCALIZED

Confined to ovarian tissue--one ovary or, if not specified to be metastatic, both ovaries

Localized, NOS

REGIONAL, Direct Extension

Peritoneum (pelvic; immediately adjacent, not implants)

Broad ligament, ipsilateral

Mesovarium, ipsilateral

Fallopian tube, ipsilateral

Adnexa, ipsilateral

REGIONAL, Lymph Nodes

Aortic (lateral and preaortic)

Hypogastric

Iliac (common, internal, external)

Obturator

Retroperitoneal/pelvic, NOS

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Sigmoid

Omentum

Cul de sac (rectouterine pouch)

Uterus

Rectosigmoid, rectum

Small intestine

Bladder, ureter

D if:

2

Implants on ovary, fallopian tube, cul de sac (rectouterine pouch),
peritoneum, omentum

Metastatic to contralateral ovary and/or fallopian tube

D if:

3

Other distant involvement

DISTANT, Lymph Nodes

Inguinal

Other distant nodes

LOCALIZED

Confined to fallopian tube(s) (specify bilateral involvement)

Localized, NOS

REGIONAL, Direct Extension

Peritoneum

Broad ligament, ipsilateral

Mesosalpinx, ipsilateral

Ovary, ipsilateral

Uterus (endometrium), ipsilateral

REGIONAL, Lymph Nodes

Aortic (lateral and preaortic)

Hypogastric

Iliac (common, internal, external)

Obturator

Retroperitoneal/pelvic, NOS

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Omentum

Cul de sac (rectouterine pouch)

Sigmoid

Rectosigmoid

Ovary, contralateral

Small intestine

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

Inguinal

Other distant nodes

IN SITU; noninvasive

LOCALIZED

L if:
1

Invasive cancer confined to submucosa (stroma)

L if:
2

Musculature invaded

L IF:
X

Localized, NOS

REGIONAL, Direct Extension

Cervix
Vulva
Cul de sac (rectouterine pouch)
Vesicovaginal septum (paracystium)
Rectovaginal septum

REGIONAL, Lymph Nodes

External iliac
Internal iliac (hypogastric)
Common iliac (sacral promontory)

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Urethra
Bladder
Rectum

D if:
2
"Frozen pelvis"

DISTANT, Lymph Nodes

Inguinal
Periaortic
Other distant lymph nodes

LYMPHATICS OF THE VULVA




Fig. 662. Lymphatics of the vulva leading to, 1, the superficial inguinal lymph nodes; 2, the superficial femoral nodes; 3, the deep inguinal nodes through the main clearing station; 4, the node of Cloquet; 5, the external iliac nodes.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

VULVA April, 1977
(including skin of vulva*)
841-844

IN SITU; noninvasive; Bowen's disease

LOCALIZED

Invasive cancer confined to submucosa
Confined to skin of vulva
Musculature invaded

Localized, NOS

REGIONAL, Direct Extension

Vaginal wall or orifice
Urethral orifice
Perineum
Perianal skin
Anus

REGIONAL, Lymph Nodes

Superficial inguinal
Deep inguinal: Rosenmüller's or Cloquet's node
External iliac

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Perineal body
Rectal mucosa

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

*Except melanomas which are classified according to the staging scheme for melanoma.

LYMPHATICS OF THE PROSTATE




Fig. 542. Schematic representation of lymphatics of the prostate showing, 1, external iliac pedicle; 2, hypogastric pedicle; and 3, posterior pedicle. The inferior pedicle which follows a downward direction and ends in hypogastric nodes is not illustrated here.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

IN_SITU; noninvasive

LOCALIZED

L if:

1

Invasive cancer confined to prostatic capsule
(intra-capsular)

L if:

2

Invasion of prostatic capsule

L if:

3

Prostatic urethra involved

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if extension:

1

Periprostatic tissues

Seminal vesicle(s)

R if:

2

Through prostatic capsule, including "fixation"

R if extension to:

3

Rectovesical (Denonvillier's) fascia

Bladder

Rectum

Extraprostatic urethra (membranous urethra)

REGIONAL, Lymph Nodes

Hypogastric
Iliac (common, internal, external)
Obturator
Periprostatic/pelvic, NOS
Sacral (lateral sacral, sacral promontory, presacral)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Skeletal muscles: levator ani
Pelvic bone
Pelvic wall
Ureter
Sigmoid colon
Penis

D if:

2

"Frozen pelvis"

D if:

3

Other distant involvement

DISTANT, Lymph Nodes

Aortic (para-aortic, periaortic, lumbar)
Inguinal
Other distant nodes

LYMPHATICS OF THE TESTIS




Fig. 562. Schematic representation of the lymphatics of the testis showing main drainage in the para-aortic lymphatics and further lymphatic extension by way of the thoracic duct. The left suprACLAVIcular node is fairly frequently involved. The lymphatics of the epididymis drain into the external iliac lymph nodes.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

LOCALIZED

Confined to tunica albuginea (encapsulated tumor)
Tunica vaginalis involved

Localized, NOS

REGIONAL, Direct Extension

Epididymis
Scrotum, ipsilateral
Spermatic cord, ipsilateral
Vas deferens

REGIONAL, Lymph Nodes

Aortic, below level of renal arteries
External iliac
Retroperitoneal/pelvic, NOS

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Ulceration of scrotum
Scrotum, contralateral
Testis, bilateral
Penis

Kidney

Adrenal gland
Retroperitoneum

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

Inguinal
Other distant nodes

IN-SITU; noninvasive (Bowen's Disease)

LOCALIZED

Invasive cancer confined to skin of penis, prepuce and/or glans

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Corpus cavernosum

R if:

2

Urethra

Satellite nodule(s) on prepuce or glans

Skin: pubic, scrotal, abdominal, perineum

REGIONAL, Lymph Nodes

External iliac

Internal iliac (hypogastric)

Superficial inguinal

Deep inguinal: Rosenmuller's or Cloquet's node

DISTANT, Direct Extension or Metastasis

Testis

Other distant involvement

DISTANT LYMPH NODE

*Except melanomas which are classified according to the staging scheme for melanoma.

LYMPHATICS OF THE BLADDER




Fig. 534. Anatomic sketch of the lymphatics of the bladder drained mainly by, 1, the external iliac nodes but also by, 2, hypogastric and, 3, common iliac nodes.

Reproduced with permission from del Regato, Juan A., and Spjut, Harlan J.: Ackerman and del Regato's Cancer, ed. 5, St. Louis, 1977, The C. V. Mosby Co.

IN SITU; noninvasive; intraepithelial

LOCALIZED

L if:

1

Confined to mucosa

L if:

2

Submucosa (subepithelial connective tissue; tunica propria; lamina propria) invaded

L if:

3

Superficial muscle (less than one half way through the muscle coat)

L if:

4

Deep muscle (half-way or more through the muscle coat)

L if:

x

Localized, NOS; no detailed information of above

REGIONAL, Direct Extension

R if:

1

Invasion of perivesical fat

Invasion of (through) serosa; peritoneum

Surrounding connective tissue (including periprostatic tissue); adjacent tissue, NOS

R if extension to:

2

Prostate, including prostatic urethra

Ureter

Vas deferens

Seminal vesicle

Rectovesical (Denonvillier's) fascia

R if extension to:

3

Rectum, male

Parametrium and uterus, female

Bladder is "fixed"

Vagina

Pubic Bone

Urethra, female

REGIONAL, Lymph Nodes

Perivesical
Hypogastric
Iliac (common, internal, external)
Obturator
Sacral (lateral sacral, presacral, sacral promontory)
Pelvic, NOS; regional, NOS

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Tumor fixed to (invading) pelvic wall
Abdominal wall
Rectum, female
Bones, excluding pubic bone
Sigmoid

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

Aortic (para-aortic, periaortic, lumbar)
Inguinal
Other distant node(s)

LOCALIZED

L if tumor confined to:

1

Kidney cortex

Kidney medulla

L if:

2

Renal pelvis or calyces involved

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Perirenal tissue (fat)

Renal (Gerota's) fascia

Retroperitoneal soft tissues (retroperitoneal space)

Blood vessels: perirenal veins, extrarenal portion of renal vein, aorta, renal artery, hilar blood vessels, vena cava

Adrenal gland, ipsilateral

Ureter, including implant(s), ipsilateral

R if:

2

Peritoneum

Diaphragm

Tail of pancreas

Ascending colon from right kidney

Descending colon from left kidney

Duodenum from right kidney

REGIONAL, Lymph Nodes

Hilar (small nodes at renal pelvis)

Lateral aortic (retroperitoneal)

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Kidney, bilateral
Ureter, contralateral
Adrenal gland, contralateral
Ribs
Stomach
Spleen
Liver

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

RENAL (KIDNEY)
PELVIS April, 1977
891

IN SITU; noninvasive; intraepithelial

LOCALIZED

Invasive cancer confined to:
Submucosa
Musculature

Localized, NOS

REGIONAL, Direct Extension

Peripelvic tissue
Retroperitoneal soft tissue (retroperitoneal space)
Major blood vessel(s): aorta, renal artery or vein, vena cava
Ureter, including implants
Kidney parenchyma
Adrenal gland
Duodenum from right renal pelvis

REGIONAL, Lymph Nodes

Hilar (renal hilus)
Lateral aortic (retroperitoneal)

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Bladder
Spleen
Pancreas
Liver
Descending colon

D if:
2

Other distant involvement

DISTANT, Lymph Nodes

IN SITU; noninvasive; intraepithelial
LOCALIZED

Invasive cancer confined to:
Submucosa
Musculature

Localized, NOS

REGIONAL, Direct Extension

Periurethral tissue
Retroperitoneal soft tissue (retroperitoneal space)
Psoas muscle

Implant(s) distal in ureter
Bladder
Kidney, ipsilateral

Duodenum from right ureter
Ascending colon from right ureter
Descending colon from left ureter

REGIONAL, Lymph Nodes

Periureteral
Hypogastric
Iliac (common, internal, external)
Lateral aortic
Retroperitoneal/pelvic, NOS

DISTANT, Direct Extension or Metastasis

D if extension to:
1

Uterus
Pancreas
Implants in bladder
Prostate

D if:
2

Other distant involvement

DISTANT, Lymph Nodes

IN SITU; noninvasive

LOCALIZED

L if:

1

Confined to one lobe and/or isthmus

L if:

2

Both lobes involved

Thyroid gland capsule involved

Multiple foci but confined to thyroid gland

L if:

3

Through capsule of gland, but not beyond

L if:

x

Localized, NOS

REGIONAL, Direct Extension

R if:

1

Pericapsular tissues

Strap muscle(s): sternothyroid, omohyoid, sternohyoid

Nerve(s): recurrent laryngeal, vagus

R if:

2

Major blood vessel(s): carotid artery, thyroid artery or vein,
jugular vein

Soft tissues of neck

Esophagus

Larynx, including thyroid and cricoid cartilages

Sternocleidomastoid muscle

Tumor is described as "fixed to adjacent tissues"

REGIONAL, Lymph Nodes

Anterior deep cervical: prelaryngeal, pretracheal, laterotracheal
(recurrent)

Internal jugular: subdigastric

Retropharyngeal

Cervical, NOS

DISTANT, Direct Extension or Metastasis

D if extension to:

1

Trachea

Mediastinal tissues

Skeletal muscle, other than strap muscles and sternocleidomastoid

Bone

D if:

2

Other distant involvement

DISTANT, Lymph Nodes

Submandibular (submaxillary)

Submental

Other distant nodes

LYMPH NODES AND LYMPHOID TISSUE Revised July, 1983
960-969, 416, 460
471, 491, 640, 692
Histology: 959 thru 969, 975

Stage I (Localized)

Confined to one lymphatic region above or below
the diaphragm

Stage II (Regional)

Involvement of more than one lymphatic region on only
one side of the diaphragm

Stage III (Distant 1)

Involvement of lymphatic regions on both sides of the
diaphragm

Stage IV (Distant 2)

Bone
Bone marrow
Lung and/or pleura
Liver
Kidney
Gastrointestinal tract (but not primary G.I.)
Skin lesions or subcutaneous nodules (but not primary skin)

SYSTEMIC SYMPTOMS

Night sweats
Unexplained fever
Pruritis
Unexplained weight loss

NOTE: Lymphoid tissue includes spleen, lingual and palatine tonsils
adenoids (pharyngeal tonsils), thymus and Waldeyer's ring, NOS.

HODGKIN'S DISEASE and NON-HODGKIN'S LYMPHOMA** Revised July, 1983
Histology: 959 thru 969, 975
(Applicable to all primary site codes)

Stage I (Localized)

Confined to one lymphatic region above or below the diaphragm

OR

Confined to a single extranodal organ or site

Stage II (Regional)

Involvement of more than one lymphatic region on only one side of the diaphragm

OR

Involvement of an extranodal organ or site with

- (1) direct extension to adjacent organs or tissues, or
- (2) involvement of one or more lymphatic regions on the same side of the diaphragm, or (3) both (1) and (2)

Stage III (Distant)¹

Involvement of lymphatic regions on both sides of the diaphragm

OR

Involvement of an extranodal organ or site with involvement

of lymphatic regions on opposite or both sides of the diaphragm

Stage IV (Distant)²

Diffuse or disseminated involvement of one or more metastatic sites with or without associated lymph node enlargement

Bone	Gastrointestinal tract
Bone marrow	Skin lesions or subcutaneous nodules
Lung and/or pleura	Brain
Liver	Eye
Kidney	

SYSTEMIC SYMPTOMS

Night sweats

Pruritis

Unexplained fever

Unexplained weight loss

**An alternative scheme for ONLY those hospitals wishing to stage lymphoma of extranodal sites. This scheme will not be used in the Certification Examination.