

Table 3b. Ranked Saltwater Genus Mean Acute Values with Species Mean Acute-Chronic Ratios

GMAV Rank	GMAV (µg/L)	Species	SMAV (µg/L)	ACR
44	6,448	Common rangia, <i>Rangia cuneata</i>	6,448	
43	4,743	Atlantic croaker, <i>Micropogon undulatus</i>	4,743	
42	4,648	Striped bass, <i>Morone saxatilis</i>	4,648	
41	2,305	Pinfish, <i>Langodon rhomboides</i>	2,305	
40	1,690	Killifish, <i>Fundulus heteroclitus</i>	1,690	
39	1,419	Mangrove rivulus, <i>Rivulus marmoratus</i>	1,419	
38	955.3	Sheepshead, <i>Archosargus probatocephalus</i>	955.3	
37	816.3	Sand shrimp, <i>Crangon spp.</i>	816.3	
36	546.3	Coho salmon, <i>Oncorhynchus kisutch</i>	546.3	
35	502.8	Green crab, <i>Carcinus maenas</i>	502.8	
34	502.8	Amphipod, <i>Corophium insidiosum</i>	502.8	
33	380.0	Shiner perch, <i>Cymatogaster aggregata</i>	380.0	
32	345.0	Florida pompano, <i>Trachinotus carolinus</i>	345.0	
31	334.5	Sheepshead minnow, <i>Cyprinodon variegatus</i>	334.5	1.48
30	318.3	Polychaete worm, <i>Hediste diversicolor</i>	318.3	
29	280.9	Squid, <i>Loligo opalescens</i>	280.9	
28	234.6	Spot, <i>Leiostomus xanthurus</i>	234.6	
27	220.9	Topsmelt, <i>Atherinops affinis</i>	220.9	
26	217.9	Nematode, <i>Caenorhabditis elegans</i>	217.9	
25	209.5	Amphipod, <i>Elasmopus bampo</i>	209.5	
24	197.3	Copepod, <i>Pseudodiaptomus coronatus</i>	197.3	
23	186.9	Copepod, <i>Tigriopus furcata</i>	178.0	
		Copepod, <i>Tigriopus californicus</i>	196.2	
22	145.2	Mysid, <i>Americamysis bahia</i>	164.5	
		Mysid, <i>Mysidopsis bigelowi</i>	128.2	
21	136.9	Polychaete worm, <i>Neanthes arenaceodentata</i>	136.9	
20	124.9	Winter flounder, <i>Pseudopleuronectes americanus</i>	124.9	
19	123.4	Mysid, <i>Neomysis mercedis</i>	123.4	
18	117.1	Tidewater silverside, <i>Menidia peninsulae</i>	117.3	
		Atlantic silverside, <i>Menidia menidia</i>	123.3	
		Inland silverside, <i>Menidia beryllina</i>	111.1	
17	100.6	Polychaete worm, <i>Phyllodoce maculata</i>	100.6	
16	86.4	Cabazon, <i>Scorpaenichthys marmoratus</i>	86.36	
15	54.98	Black abalone, <i>Haliotis cracherodii</i>	41.90	
		Red abalone, <i>Haliotis rufescens</i>	72.14	
14	42.74	Surf clam, <i>Spisula solidissima</i>	42.74	
13	41.06	Dungeness crab, <i>Cancer magister</i>	41.06	
12	40.22	American lobster, <i>Homarus americanus</i>	40.22	
11	32.68	Soft-shell clam, <i>Mya arenaria</i>	32.68	
10	32.45	Copepod, <i>Acartia tonsa</i>	25.74	

Table 3b. Ranked Saltwater Genus Mean Acute Values with Species Mean Acute-Chronic Ratios

GMAV Rank	GMAV (µg/L)	Species	SMAV (µg/L)	ACR
		Copepod, <i>Acartia clausi</i>	40.89	
9	25.83	Copepod, <i>Eurytemora affinis</i>	25.83	
8	24.30	Bay scallop, <i>Argopecten irradians</i>	24.30	
7	21.40	Sea urchin, <i>Arbacia punctulata</i>	21.40	
6	17.69	Coot clam, <i>Mulina lateralis</i>	17.69	
5	15.45	Mysid, <i>Holmesimysis costata</i>	15.45	
4	12.81	Sea urchin, <i>Strongylocentrotus purpuratus</i>	12.81	
3	12.66	Summer flounder, <i>Paralichthys dentatus</i>	12.66	
2	12.60	Eastern oyster, <i>Crassostrea virginica</i>	14.49	
		Pacific oyster, <i>Crassostrea gigas</i>	10.96	
1	11.53	Blue mussel, <i>Mytilus edulis</i>	21.50	
		Mussel, <i>Mytilus sp.</i>	6.19	